

A MATHEMATICAL SYN-ECOLOGICAL MODEL COMPRISING  
OF PREY-PREDATOR, HOST-COMMENSAL, MUTUALISM AND NEUTRAL PAIRS

R. Srilatha<sup>1</sup>, B. Ravindra Reddy<sup>2\*</sup> and N. Ch. Pattabhiramacharyulu<sup>3</sup>

<sup>1</sup>Sree Chaitanya Inst. of Tech. & Sci., Thimmapur, Karimnagar-505527, India

<sup>2</sup>JNTUH College of Engineering, Nachupally, Karimnagar-505501, India

<sup>3</sup>Professor (Retd.) of Mathematics, NIT, Warangal – 506004, India

E-mail: [bsrilatha82@gmail.com](mailto:bsrilatha82@gmail.com)

(Received on: 09-04-12; Accepted on: 25-04-12)

ABSTRACT

This investigation deals with a mathematical model of a four species ( $S_1, S_2, S_3$  and  $S_4$ ) Syn-Ecological system.  $S_2$  is a predator surviving on the prey  $S_1$ . The predator  $S_2$  is a commensal to the host  $S_3$ . The pairs  $S_2$  and  $S_4, S_1$  and  $S_3$  are neutral. The mathematical model equations characterizing the syn-ecosystem constitute a set of four first order non-linear coupled differential equations. There are in all sixteen equilibrium points. Criteria for the asymptotic stability of the sixteen equilibrium points are established. The linearised equations for the perturbations over the equilibrium point are analyzed to establish the criteria for stability. Trajectories of the perturbations have been illustrated. Also global stability is discussed by Lyapunov's function. Further analytical stability criteria are supported by numerical simulations using Mat lab.

**Key words:** Equilibrium state, stability, Mutualism, Co-Existent State.

1. INTRODUCTION:

Mathematical modeling is an important interdisciplinary activity which involves the study of some aspects of diverse disciplines. Biology, Epidemiology, Physiology, Ecology, Immunology, Bio-economics, Genetics, Pharmacokinetics are some of those disciplines. This mathematical modeling has raised to the zenith in recent years and spread to all branches of life and drew the attention of every one. Mathematical modeling of ecosystems was initiated by Lotka [6] and by Volterra [12] followed by several mathematicians and ecologists. They contributed their might to the growth of this area of knowledge as reported in the treatises of Meyer [7], Paul Colinvaux [8], Freedman [2], Kapur [3, 4]. The ecological interactions can be broadly classified as prey-predation, competition, mutualism and so on. N.C. Srinivas [11] studied the competitive eco-systems of two species and three species with regard to limited and unlimited resources. Later, Lakshmi Narayan [5] has investigated the two species prey-predator models and stability analysis of competitive species was investigated by Archana [1]. Further Local stability analysis for a two-species ecological mutualism model has been investigated by B. Ravindra Reddy et al. [9, 10]. In this connection here we constructed a four species ( $S_1, S_2, S_3$  and  $S_4$ ) mathematical model based on the system of non-linear equations.  $S_2$  is a predator surviving on the prey  $S_1$ . The predator  $S_2$  is a commensal to the host  $S_3$ . The pairs  $S_2$  and  $S_4, S_1$  and  $S_3$  are neutral. Equilibrium points of the system are identified and the stability analysis is carried out. Example for  $S_1, S_2, S_3$  and  $S_4$  are Insects, Insectivorous Plants (nephantis, drosera etc.), VAM associated with the plant roots, Soil bacteria respectively.

2. BASIC EQUATIONS:

The model equations for a four species multi-system are given by a set of four non-linear ordinary differential equations as

- (i) For  $S_1$ : The Prey of  $S_1$  and Neutral to  $S_3$

$$\frac{dN_1}{dt} = a_1N_1 - a_{11}N_1^2 - a_{12}N_1N_2 \quad (2.1)$$

\*Corresponding author: B. Ravindra Reddy<sup>2\*</sup>, \*E-mail: [rbollareddy@gmail.com](mailto:rbollareddy@gmail.com)

(ii) For  $S_2$ : The Predator surviving on  $S_1$  and Commensal to  $S_3$

$$\frac{dN_2}{dt} = a_2 N_2 - a_{22} N_2^2 + a_{21} N_2 N_1 + a_{23} N_2 N_3 \quad (2.2)$$

(iii) For  $S_3$ : The Host of  $S_2$  and Mutual to  $S_4$

$$\frac{dN_3}{dt} = a_3 N_3 - a_{33} N_3^2 + a_{34} N_3 N_4 \quad (2.3)$$

(iv) For  $S_4$ : Mutual to  $S_3$  and Neutral to  $S_2$

$$\frac{dN_4}{dt} = a_4 N_4 - a_{44} N_4^2 + a_{43} N_4 N_3 \quad (2.4)$$

with the following notation.

$N_i(t)$ : Population strengths of the species  $S_i$  at time  $t$ ,  $i=1, 2, 3, 4$ .

$a_i$  : The natural growth rates of  $S_i$ ,  $i = 1,2,3,4$

$a_{12}, a_{21}$  : Interaction (Prey-Predator) coefficients of  $S_1$  due to  $S_2$  and  $S_2$  due to  $S_1$

$a_{13}$  : Coefficient for commensal for  $S_1$  due to the Host  $S_3$

$a_{34}, a_{43}$  : Mutually interaction between  $S_3$  and  $S_4$

$K_i = \frac{a_i}{a_{ii}}$  : Carrying capacities of  $S_i$ ,  $i=1, 2, 3, 4$ .

Further the variables  $N_1, N_2, N_3, N_4$  are non-negative and the model parameters  $a_1, a_2, a_3, a_4; a_{11}, a_{22}, a_{33}, a_{44}; a_{12}, a_{21}, a_{13}, a_{24}$  are assumed to be non-negative constants.

## 1. EQUILIBRIUM STATES:

The system under investigation has sixteen equilibrium states defined by

$$\frac{dN_i}{dt} = 0, \quad i = 1, 2, 3, 4 \quad (3.1)$$

are given in the following table.

**I.** Fully washed out state:

$$E_1: \quad \overline{N}_1 = 0, \overline{N}_2 = 0, \overline{N}_3 = 0, \overline{N}_4 = 0$$

**II.** States in which three of the four species are washed out and fourth is surviving

$$E_2: \quad \overline{N}_1 = 0, \overline{N}_2 = 0, \overline{N}_3 = 0, \overline{N}_4 = \frac{a_4}{a_{44}}$$

$$E_3: \quad \overline{N}_1 = 0, \overline{N}_2 = 0, \overline{N}_3 = \frac{a_3}{a_{33}}, \overline{N}_4 = 0$$

$$E_4: \quad \overline{N}_1 = 0, \overline{N}_2 = \frac{a_2}{a_{22}}, \overline{N}_3 = 0, \overline{N}_4 = 0$$

$$E_5: \quad \overline{N}_1 = \frac{a_1}{a_{11}}, \overline{N}_2 = 0, \overline{N}_3 = 0, \overline{N}_4 = 0$$

**III.** States in which two of the four species are washed out while the other two are surviving

$$E_6: \quad \overline{N}_1 = 0, \overline{N}_2 = 0, \overline{N}_3 = \frac{a_4 a_{34} + a_3 a_{44}}{a_{33} a_{44} - a_{34} a_{43}}, \overline{N}_4 = \frac{a_3 a_{43} + a_4 a_{33}}{a_{33} a_{44} - a_{34} a_{43}}$$

This state exists only when  $a_{33} a_{44} - a_{34} a_{43} > 0$

$$E_7: \quad \overline{N}_1 = 0, \overline{N}_2 = \frac{a_2}{a_{22}}, \overline{N}_3 = 0, \overline{N}_4 = \frac{a_4}{a_{44}}$$

$$\begin{aligned}
 E_8: \quad & \overline{N}_1 = 0, \overline{N}_2 = \frac{a_3 a_{23}}{a_{22} a_{33}} + \frac{a_2}{a_{22}}, \overline{N}_3 = \frac{a_3}{a_{33}}, \overline{N}_4 = 0 \\
 E_9: \quad & \overline{N}_1 = \frac{a_1}{a_{11}}, \overline{N}_2 = 0, \overline{N}_3 = 0, \overline{N}_4 = \frac{a_4}{a_{44}} \\
 E_{10}: \quad & \overline{N}_1 = \frac{a_1}{a_{11}}, \overline{N}_2 = 0, \overline{N}_3 = \frac{a_3}{a_{33}}, \overline{N}_4 = 0 \\
 E_{11}: \quad & \overline{N}_1 = \frac{a_1 a_{22} - a_2 a_{12}}{a_{11} a_{22} + a_{12} a_{21}}, \overline{N}_2 = \frac{a_1 a_{21} + a_2 a_{11}}{a_{11} a_{22} + a_{12} a_{21}}, \overline{N}_3 = 0, \overline{N}_4 = 0
 \end{aligned}$$

This state exists only when  $a_1 a_{22} - a_2 a_{12} > 0$

#### IV. States in which one of the four species is washed out while the other three are surviving

$$\begin{aligned}
 E_{12}: \quad & \overline{N}_1 = 0, \overline{N}_2 = \frac{a_{23}(a_4 a_{34} + a_3 a_{44})}{a_{22}(a_{33} a_{44} - a_{34} a_{43})} + \frac{a_2}{a_{22}}, \overline{N}_3 = \frac{a_4 a_{34} + a_3 a_{44}}{a_{33} a_{44} - a_{34} a_{43}}, \\
 & \overline{N}_4 = \frac{a_4 a_{33} + a_3 a_{43}}{a_{33} a_{44} - a_{34} a_{43}}
 \end{aligned}$$

This state exists only when  $a_{33} a_{44} - a_{34} a_{43} > 0$

$$E_{13}: \quad \overline{N}_1 = \frac{a_1}{a_{11}}, \overline{N}_2 = 0, \overline{N}_3 = \frac{a_4 a_{34} + a_3 a_{44}}{a_{33} a_{44} - a_{34} a_{43}}, \overline{N}_4 = \frac{a_4 a_{33} + a_3 a_{43}}{a_{33} a_{44} - a_{34} a_{43}}$$

This state exists only when  $(a_{33} a_{44} - a_{34} a_{43}) > 0$

$$E_{14}: \quad \overline{N}_1 = \frac{a_1 a_{22} - a_2 a_{12}}{a_{11} a_{22} + a_{12} a_{21}}, \overline{N}_2 = \frac{a_1 a_{21} + a_2 a_{11}}{a_{11} a_{22} + a_{12} a_{21}}, \overline{N}_3 = 0, \overline{N}_4 = \frac{a_4}{a_{44}}$$

This state exists only when  $a_1 a_{22} - a_2 a_{12} > 0$

$$E_{15}: \quad \overline{N}_1 = \frac{\beta_4}{\beta_1}, \overline{N}_2 = \frac{\beta_5}{\beta_1}, \overline{N}_3 = \frac{a_3}{a_{33}}, \overline{N}_4 = 0$$

Where

$$\beta_1 = a_{33}(a_{11} a_{22} + a_{12} a_{21})$$

$$\beta_4 = a_{33}(a_1 a_{22} - a_2 a_{12}) - a_3 a_{23} a_{12}$$

$$\beta_5 = a_{33}(a_1 a_{21} + a_2 a_{11}) + a_3 a_{23} a_{11}$$

This state exists only when  $\beta_4 > 0$

#### V. The co-existent state (or) Normal steady state

$$\begin{aligned}
 E_{16}: \quad & \overline{N}_1 = \frac{\gamma_1 + a_{12} a_{23} \gamma_2}{\gamma_3 (a_{33} a_{44} - a_{34} a_{43})}, \overline{N}_2 = \frac{\gamma_4 + a_{11} a_{23} \gamma_2}{\gamma_3 (a_{33} a_{44} - a_{34} a_{43})}, \\
 & \overline{N}_3 = \frac{a_4 a_{34} + a_3 a_{44}}{a_{33} a_{44} - a_{34} a_{43}}, \overline{N}_4 = \frac{a_4 a_{33} + a_3 a_{43}}{a_{33} a_{44} - a_{34} a_{43}}
 \end{aligned}$$

Where

$$\gamma_1 = (a_1 a_{22} + a_2 a_{12})(a_{33} a_{44} - a_{34} a_{43}), \gamma_2 = a_3 a_{44} + a_4 a_{34}$$

$$\gamma_3 = a_{11} a_{22} + a_{12} a_{21}, \gamma_4 = (a_1 a_{21} - a_2 a_{11})(a_{33} a_{44} - a_{34} a_{43})$$

This state exists only when  $(a_1 a_{21} - a_2 a_{11}) > 0$  and  $(a_{33} a_{44} - a_{34} a_{43}) > 0$ .

#### 4. STABILITY OF THE EQUILIBRIUM STATES:

$$\text{Let } N = (N_1, N_2, N_3, N_4) = \bar{N} + U \tag{4.1}$$

where  $U = (u_1, u_2, u_3, u_4)$  is a perturbation over the equilibrium state  $\bar{N} = (\bar{N}_1, \bar{N}_2, \bar{N}_3, \bar{N}_4)$ .

The basic equations (2.1), (2.2), (2.3), (2.4) are linearized to obtain the equations for the perturbed state as

$$\frac{dU}{dt} = AU \tag{4.2}$$

where

$$A = \begin{bmatrix} a_1 - 2a_{11}\bar{N}_1 - a_{12}\bar{N}_2 & -a_{12}\bar{N}_1 & 0 & 0 \\ a_{21}\bar{N}_2 & a_2 - 2a_{22}\bar{N}_2 + a_{21}\bar{N}_1 + a_{23}\bar{N}_3 & a_{23}\bar{N}_2 & 0 \\ 0 & 0 & a_3 - 2a_{33}\bar{N}_3 + a_{34}\bar{N}_4 & a_{34}\bar{N}_3 \\ 0 & 0 & a_{43}\bar{N}_4 & a_4 - 2a_{44}\bar{N}_4 + a_{43}\bar{N}_3 \end{bmatrix} \tag{4.3}$$

$$\text{The characteristic equation for the system is } \det[A - \lambda I] = 0 \tag{4.4}$$

The equilibrium state is stable if both the roots of the equation (4.4) are negative in case they are real or have negative real parts in case they are complex. The equilibrium states  $E_1, E_2, E_3, E_4, E_5, E_6, E_7, E_8, E_9, E_{10}, E_{11}, E_{13}, E_{14}, E_{15}$  are noticed to be unstable (The detailed investigation of these states is not included here for treatment). The stability criteria of states  $E_{12}, E_{16}$  are discussed below.

#### 4.1 Stability of Prey ( $S_1$ ) Washed out Equilibrium State: ( $E_{12}$ )

$$\bar{N}_1 = 0, \bar{N}_2 = \frac{a_{23}(a_4 a_{34} + a_3 a_{44})}{a_{22}(a_{33} a_{44} - a_{34} a_{43})} + \frac{a_2}{a_{22}}, \bar{N}_3 = \frac{a_4 a_{34} + a_3 a_{44}}{a_{33} a_{44} - a_{34} a_{43}}, \bar{N}_4 = \frac{a_4 a_{33} + a_3 a_{43}}{a_{33} a_{44} - a_{34} a_{43}} \tag{4.1.1}$$

Substituting (4.1.1) in (2.1), (2.2), (2.3), (2.4) and neglecting products and higher powers of  $u_1, u_2, u_3, u_4$ , we get

$$\frac{du_1}{dt} = w_1 u_1 \tag{4.1.2}$$

$$\frac{du_2}{dt} = a_{21} \bar{N}_2 u_1 - w_3 u_2 + a_{23} \bar{N}_2 u_3 \tag{4.1.3}$$

$$\frac{du_3}{dt} = -a_{33} \bar{N}_3 u_3 + a_{34} \bar{N}_3 u_4 \tag{4.1.4}$$

$$\frac{du_4}{dt} = a_{43} \bar{N}_4 u_3 - a_{44} \bar{N}_4 u_4 \tag{4.1.5}$$

$$\text{Here } w_1 = a_1 - a_{12} \bar{N}_2, w_3 = a_2 + a_{23} \bar{N}_3 \tag{4.1.6}$$

The characteristic equation of which is

$$(\lambda - w_1)(\lambda + w_3) \left[ \lambda^2 + (a_{33} \bar{N}_3 + a_{44} \bar{N}_4) \lambda + (a_{33} a_{44} - a_{34} a_{43}) \bar{N}_3 \bar{N}_4 \right] = 0 \tag{4.1.7}$$

The characteristic roots of (4.1.7) are

$$\lambda = w_1, \lambda = -w_3, \lambda = \frac{-(a_{33} \bar{N}_3 + a_{44} \bar{N}_4) \pm \sqrt{(a_{33} \bar{N}_3 - a_{44} \bar{N}_4)^2 + 4a_{34} a_{43} \bar{N}_3 \bar{N}_4}}{2}$$

Case (A): If  $w_1 < 0$  [i.e.  $a_1 < a_{12} \bar{N}_2$ ]

Here  $w_1, -w_3$  are negative and the other two roots are also negative.

Hence the equilibrium state is **stable**.

The solutions of the equations (4.1.2), (4.1.3), (4.1.4), (4.1.5) are

$$u_1 = u_{10} e^{w_1 t} \tag{4.1.8}$$

$$u_2 = \left[ u_{20} - \frac{a_{21} \bar{N}_2 u_{10}}{(w_1 + w_3)} - \frac{a_{23} \bar{N}_2 (P_1 + P_2)}{(\lambda_3 + w_3)} \right] e^{-w_3 t} + \frac{a_{21} \bar{N}_2 u_{10}}{(w_1 + w_3)} e^{w_1 t} + \frac{a_{23} \bar{N}_2 (P_1 e^{\lambda_3 t} + P_2 e^{\lambda_4 t})}{(\lambda_3 + w_3)} \tag{4.1.9}$$

$$u_3 = \left[ \frac{u_{30} (\lambda_3 + a_{44} \bar{N}_4) + u_{40} a_{34} \bar{N}_3}{\lambda_3 - \lambda_4} \right] e^{\lambda_3 t} + \left[ \frac{u_{30} (\lambda_4 + a_{44} \bar{N}_4) + u_{40} a_{34} \bar{N}_3}{\lambda_4 - \lambda_3} \right] e^{\lambda_4 t} \tag{4.1.10}$$

$$u_4 = \left[ \frac{u_{40} (\lambda_3 + a_{33} \bar{N}_3) + u_{30} a_{43} \bar{N}_4}{\lambda_3 - \lambda_4} \right] e^{\lambda_3 t} + \left[ \frac{u_{40} (\lambda_4 + a_{33} \bar{N}_3) + u_{30} a_{43} \bar{N}_4}{\lambda_4 - \lambda_3} \right] e^{\lambda_4 t} \tag{4.1.11}$$

Where  $P_1 = \frac{u_{30} (\lambda_3 + a_{44} \bar{N}_4) + u_{40} a_{34} \bar{N}_3}{\lambda_3 - \lambda_4}$ ,  $P_2 = \frac{u_{30} (\lambda_4 + a_{44} \bar{N}_4) + u_{40} a_{34} \bar{N}_3}{\lambda_4 - \lambda_3}$

where  $u_{10}, u_{20}, u_{30}, u_{40}$  are the initial values of  $u_1, u_2, u_3, u_4$  respectively.

There would arise in all 576 cases depending upon the ordering of the magnitudes of the growth rates  $a_1, a_2, a_3, a_4$  and the initial values of the perturbations  $u_{10}(t), u_{20}(t), u_{30}(t), u_{40}(t)$  of the species  $S_1, S_2, S_3, S_4$ . Of these 576 situations some typical variations are illustrated through respective solution curves that would facilitate to make some reasonable observations.

The solutions are illustrated in figures.

Case (i): If  $u_{30} < u_{20} < u_{10} < u_{40}$  and  $a_3 < a_1 < A_2 < a_4$

In this case initially  $S_4$  dominates the Prey ( $S_1$ ) and the Predator ( $S_2$ ) till the time instant  $t_{14}^*, t_{24}^*$  respectively and thereafter the dominance is reversed. And  $u_1, u_2, u_3, u_4$  are converging asymptotically to the equilibrium point. Hence the equilibrium point is **stable**.




Fig. 1

Case (ii): If  $u_{40} < u_{10} < u_{30} < u_{20}$  and  $A_2 < a_3 < a_1 < a_4$

In this case initially the Predator ( $S_2$ ) dominates the Prey ( $S_1$ ) till the time instant  $t_{12}^*$  and thereafter the dominance is reversed. Also the host ( $S_3$ ) of  $S_2$  dominates the Prey ( $S_1$ ) and  $S_4$  till the time instant  $t_{13}^*, t_{43}^*$  respectively and the dominance gets reversed thereafter.




Fig. 2

**Case (B):** If  $w_1 > 0$  [i.e.  $a_1 > a_{12}\bar{N}_2$ ]

Here the root  $w_1$  is positive and the other three roots are negative.

Hence the equilibrium state is **unstable** and the solutions in this case are same as in Case (A).

#### 4.2 Stability of Co-Existing State : (E<sub>16</sub>)

$$\bar{N}_1 = \frac{\gamma_1 + a_{12}a_{23}\gamma_2}{\gamma_3(a_{33}a_{44} - a_{34}a_{43})}, \bar{N}_2 = \frac{\gamma_4 + a_{11}a_{23}\gamma_2}{\gamma_3(a_{33}a_{44} - a_{34}a_{43})}, \bar{N}_3 = \frac{a_4a_{34} + a_3a_{44}}{a_{33}a_{44} - a_{34}a_{43}},$$

$$\bar{N}_4 = \frac{a_4a_{33} + a_3a_{43}}{a_{33}a_{44} - a_{34}a_{43}}$$
(4.2.1)

Where

$$\gamma_1 = (a_1a_{22} + a_2a_{12})(a_{33}a_{44} - a_{34}a_{43}), \gamma_2 = a_3a_{44} + a_4a_{34}$$

$$\gamma_3 = (a_{11}a_{22} + a_{12}a_{21}), \gamma_4 = (a_1a_{21} - a_2a_{11})(a_{33}a_{44} - a_{34}a_{43})$$

$$\text{This can exist only when } (a_1a_{21} - a_2a_{11}) > 0 \text{ and } (a_{33}a_{44} - a_{34}a_{43}) > 0$$
(4.2.2)

Let us consider small deviations  $u_1(t), u_2(t), u_3(t), u_4(t)$  from the steady state

$$\text{i.e. } N_i(t) = \bar{N}_i + u_i(t), i = 1, 2, 3, 4$$
(4.2.3)

Substituting (4.2.1) in (2.1), (2.2), (2.3), (2.4) and neglecting products and higher powers of  $u_1, u_2, u_3, u_4$ , we get

$$\frac{du_1}{dt} = -a_{11}\bar{N}_1u_1 - a_{12}\bar{N}_1u_2$$
(4.2.4)

$$\frac{du_2}{dt} = -a_{22}\bar{N}_2u_2 + a_{21}\bar{N}_2u_1 + a_{23}\bar{N}_2u_3$$
(4.2.5)

$$\frac{du_3}{dt} = -a_{33}\bar{N}_3u_3 + a_{34}\bar{N}_3u_4$$
(4.2.6)

$$\frac{du_4}{dt} = -a_{44}\bar{N}_4u_4 + a_{43}\bar{N}_4u_3$$
(4.2.7)

The characteristic equation of which is

$$\left[ \lambda^2 + (a_{11}\bar{N}_1 + a_{22}\bar{N}_2)\lambda + (a_{11}a_{22} + a_{12}a_{21})\bar{N}_1\bar{N}_2 \right] \times$$

$$\left[ \lambda^2 + (a_{33}\bar{N}_3 + a_{44}\bar{N}_4)\lambda + (a_{33}a_{44} - a_{34}a_{43})\bar{N}_3\bar{N}_4 \right] = 0$$
(4.2.8)

The characteristic roots of (4.2.8) are

$$\lambda = \frac{-(a_{11}\bar{N}_1 + a_{22}\bar{N}_2) \pm \sqrt{(a_{11}\bar{N}_1 - a_{22}\bar{N}_2)^2 - 4a_{12}a_{21}\bar{N}_1\bar{N}_2}}{2},$$

$$\lambda = \frac{-(a_{33}\bar{N}_3 + a_{44}\bar{N}_4) \pm \sqrt{(a_{33}\bar{N}_3 - a_{44}\bar{N}_4)^2 + 4a_{34}a_{43}\bar{N}_3\bar{N}_4}}{2}$$
(4.2.9)

$$\Rightarrow \lambda = \frac{-(a_{11}\bar{N}_1 + a_{22}\bar{N}_2) \pm \sqrt{\Delta_1}}{2}, \lambda = \frac{-(a_{33}\bar{N}_3 + a_{44}\bar{N}_4) \pm \sqrt{\Delta_2}}{2}$$
(4.2.10)

Where

$$\Delta_1 = (a_{11}\bar{N}_1 - a_{22}\bar{N}_2)^2 - 4a_{12}a_{21}\bar{N}_1\bar{N}_2, \Delta_2 = (a_{33}\bar{N}_3 - a_{44}\bar{N}_4)^2 + 4a_{34}a_{43}\bar{N}_3\bar{N}_4$$

**Case (i):** When  $\Delta_1 > 0$  and  $\Delta_2 > 0$

In this case the roots are real and negative.

Hence the equilibrium state is **stable**.

**Case (ii):** When  $\Delta_1 < 0$  and  $\Delta_2 < 0$

In this case the roots are complex with negative real parts.

Hence the equilibrium state is **stable**.

**Case (iii):** When  $\Delta_1 = 0$  and  $\Delta_2 = 0$

In this case the roots are repeated, which are negative.

Hence the equilibrium state is **stable**.

The trajectories are given by

$$u_1 = \left[ \frac{a_{12}\bar{N}_1(u_{10} - u_{20}) + \mu_1\lambda_2 + \mu_2a_{13}\bar{N}_1 - \mu_3}{\lambda_1 - \lambda_2} \right] e^{\lambda_1 t} + \left[ \frac{(\mu_1 - u_{10})\lambda_1 + u_{10}\lambda_2 + (a_{12}u_{10} - a_{12}u_{20} + \mu_2a_{13})\bar{N}_1 - \mu_3}{\lambda_2 - \lambda_1} \right] e^{\lambda_2 t} + \sigma_1 e^{\lambda_3 t} + \sigma_2 e^{\lambda_4 t} \quad (4.2.11)$$

$$u_2 = \left[ \frac{a_{12}\bar{N}_1(u_{10} - u_{20}) + \mu_1\lambda_2 + \mu_2a_{13}\bar{N}_1 - \mu_3}{\lambda_1 - \lambda_2} \right] \delta_1 e^{\lambda_1 t} + \left[ \frac{(\mu_1 - u_{10})\lambda_1 + u_{10}\lambda_2 + (a_{12}u_{10} - a_{12}u_{20} + \mu_2a_{13})\bar{N}_1 - \mu_3}{\lambda_2 - \lambda_1} \right] \delta_2 e^{\lambda_2 t} + \sigma_3 e^{\lambda_3 t} + \sigma_4 e^{\lambda_4 t} \quad (4.2.12)$$

$$u_3 = \left[ \frac{u_{30}(\lambda_3 + a_{44}\bar{N}_4) + u_{40}a_{34}\bar{N}_3}{\lambda_3 - \lambda_4} \right] e^{\lambda_3 t} + \left[ \frac{u_{30}(\lambda_4 + a_{44}\bar{N}_4) + u_{40}a_{34}\bar{N}_3}{\lambda_4 - \lambda_3} \right] e^{\lambda_4 t} \quad (4.2.13)$$

$$u_4 = \left[ \frac{u_{40}(\lambda_3 + a_{33}\bar{N}_3) + u_{30}a_{43}\bar{N}_4}{\lambda_3 - \lambda_4} \right] e^{\lambda_3 t} + \left[ \frac{u_{40}(\lambda_4 + a_{33}\bar{N}_3) + u_{30}a_{43}\bar{N}_4}{\lambda_4 - \lambda_3} \right] e^{\lambda_4 t} \quad (4.2.14)$$

Here

$$\mu_1 = \sigma_1 + \sigma_2; \mu_2 = p_1 + p_2; \mu_3 = \sigma_1\lambda_3 + \sigma_2\lambda_4;$$

$$\sigma_1 = \frac{\alpha_2}{\lambda_3^2 + (a_{11}\bar{N}_1 + a_{22}\bar{N}_2)\lambda_3 + \alpha_1}; \sigma_2 = \frac{\alpha_3}{\lambda_4^2 + (a_{11}\bar{N}_1 + a_{22}\bar{N}_2)\lambda_4 + \alpha_1};$$

$$\alpha_1 = (a_{11}a_{22} + a_{12}a_{21})\bar{N}_1\bar{N}_2; \alpha_2 = p_1a_{13}\bar{N}_1(\lambda_3 + a_{22}\bar{N}_2); \alpha_3 = p_2a_{13}\bar{N}_3(\lambda_4 + a_{22}\bar{N}_2);$$

$$p_1 = \frac{u_{30}(\lambda_3 + a_{44}\bar{N}_4) + u_{40}a_{34}\bar{N}_3}{\lambda_3 - \lambda_4}; p_2 = \frac{u_{30}(\lambda_4 + a_{44}\bar{N}_4) + u_{40}a_{34}\bar{N}_3}{\lambda_4 - \lambda_3}; \delta_1 = \frac{a_{11}}{a_{12}} - \frac{\lambda_1}{a_{12}\bar{N}_1};$$

$$\delta_2 = \frac{a_{11}}{a_{12}} - \frac{\lambda_2}{a_{12}\bar{N}_1}; \sigma_3 = \frac{a_{13}P_1}{a_{12}} + \left( a_{11} - \frac{\lambda_3}{\bar{N}_1} \right) \frac{\sigma_1}{a_{12}}; \sigma_4 = \frac{a_{13}P_2}{a_{12}} + \left( a_{11} - \frac{\lambda_4}{\bar{N}_1} \right) \frac{\sigma_2}{a_{12}}$$

and  $u_{10}, u_{20}, u_{30}, u_{40}$  are the initial values of  $u_1, u_2, u_3, u_4$  respectively.

There would arise in all 576 cases depending upon the ordering of the magnitudes of the growth rates  $a_1, a_2, a_3, a_4$  and the initial values of the perturbations  $u_{10}(t), u_{20}(t), u_{30}(t), u_{40}(t)$  of the species  $S_1, S_2, S_3, S_4$ . Of these 576 situations some typical variations are illustrated through respective solution curves that would facilitate to make some reasonable observations.

The solutions are illustrated in figures.

**Case (i):** If  $u_{20} < u_{30} < u_{40} < u_{10}$  and  $a_3 < a_4 < a_1 < a_2$

In this case initially  $S_4$  dominates the Prey ( $S_1$ ) till the time instant  $t_{14}^*$  and the dominance gets reversed thereafter. And  $u_1, u_2, u_3, u_4$  are converging asymptotically to the equilibrium point.

Hence the equilibrium point is **stable**.




Fig. 3

**Case (ii):** If  $u_{40} < u_{30} < u_{20} < u_{10}$  and  $a_2 < a_3 < a_1 < a_4$

In this case initially  $S_4$  dominates the Prey ( $S_1$ ) and the Predator ( $S_2$ ) till the time instant  $t_{14}^*, t_{24}^*$  respectively and thereafter the dominance is reversed. Also the host ( $S_3$ ) of  $S_2$  dominates the Predator ( $S_2$ ) and the Prey ( $S_1$ ) till the time instant  $t_{23}^*, t_{13}^*$  respectively and the dominance gets reversed thereafter. And the Predator ( $S_2$ ) dominates the Prey ( $S_1$ ) till the time instant  $t_{12}^*$  and thereafter the dominance is reversed.




Fig. 4

## 5. LIAPUNOV'S FUNCTION FOR GLOBAL STABILITY

We discussed the local stability of the state of co-existence. We now examine the global stability of the dynamical system (2.1), (2.2), (2.3) and (2.3). We have already noted that this system has a unique, stable non-trivial co-existent

$$\text{equilibrium state at } \bar{N}_1 = \frac{\gamma_1 + a_{13}a_{22}\gamma_2}{\gamma_3}, \bar{N}_2 = \frac{\gamma_4 + a_{13}a_{21}\gamma_2}{\gamma_3}, \bar{N}_3 = \frac{a_4a_{34} + a_3a_{44}}{a_{33}a_{44} - a_{34}a_{43}}, \bar{N}_4 = \frac{a_4a_{33} + a_3a_{43}}{a_{33}a_{44} - a_{34}a_{43}}$$

We define a Liapunov function

$$V(N_1, N_2, N_3, N_4) = N_1 - \bar{N}_1 - \bar{N}_1 \log\left(\frac{N_1}{\bar{N}_1}\right) + l_1 \left\{ N_2 - \bar{N}_2 - \bar{N}_2 \log\left(\frac{N_2}{\bar{N}_2}\right) \right. \\ \left. + l_2 \left\{ N_3 - \bar{N}_3 - \bar{N}_3 \log\left(\frac{N_3}{\bar{N}_3}\right) \right\} + l_3 \left\{ N_4 - \bar{N}_4 - \bar{N}_4 \log\left(\frac{N_4}{\bar{N}_4}\right) \right\} \right\} \quad (5.1)$$

where  $l_1, l_2$  and  $l_3$  are suitable constants to be determined in the subsequent steps.



Now, the time derivative of  $V$  along the solution of (2.1), (2.2), (2.3) and (2.4) is

$$\frac{dV}{dt} = \left( \frac{N_1 - \bar{N}_1}{N_1} \right) \frac{dN_1}{dt} + l_1 \left( \frac{N_2 - \bar{N}_2}{N_2} \right) \frac{dN_2}{dt} + l_2 \left( \frac{N_3 - \bar{N}_3}{N_3} \right) \frac{dN_3}{dt} + l_3 \left( \frac{N_4 - \bar{N}_4}{N_4} \right) \frac{dN_4}{dt} \quad (5.2)$$

$$\begin{aligned} \frac{dV}{dt} = & \left( \frac{N_1 - \bar{N}_1}{N_1} \right) N_1 \{a_1 - a_{11}N_1 - a_{12}N_2\} + l_1 \left( \frac{N_2 - \bar{N}_2}{N_2} \right) N_2 \{a_2 - a_{22}N_2 + a_{21}N_1 + a_{23}N_3\} \\ & + l_2 \left( \frac{N_3 - \bar{N}_3}{N_3} \right) N_3 \{a_3 - a_{33}N_3 + a_{34}N_4\} + l_3 \left( \frac{N_4 - \bar{N}_4}{N_4} \right) N_4 \{a_4 - a_{44}N_4 + a_{43}N_3\} \end{aligned} \quad (5.3)$$

$$\begin{aligned} = & (N_1 - \bar{N}_1) \{a_1 - a_{11}N_1 - a_{12}N_2\} + l_1 (N_2 - \bar{N}_2) \{a_2 - a_{22}N_2 + a_{21}N_1 + a_{23}N_3\} \\ & + l_2 (N_3 - \bar{N}_3) \{a_3 - a_{33}N_3 + a_{34}N_4\} + l_3 (N_4 - \bar{N}_4) \{a_4 - a_{44}N_4 + a_{43}N_3\} \end{aligned} \quad (5.4)$$

$$\begin{aligned} \frac{dV}{dt} = & (N_1 - \bar{N}_1) \{a_{11}\bar{N}_1 + a_{12}\bar{N}_2 - a_{11}N_1 - a_{12}N_2\} + l_1 (N_2 - \bar{N}_2) \{a_{22}\bar{N}_2 - a_{21}\bar{N}_1 - a_{23}\bar{N}_3 - a_{22}N_2 + a_{21}N_1 + a_{23}N_3\} \\ & + l_2 (N_3 - \bar{N}_3) \{a_{33}\bar{N}_3 - a_{34}\bar{N}_4 - a_{33}N_3 + a_{34}N_4\} + l_3 (N_4 - \bar{N}_4) \{a_{44}\bar{N}_4 - a_{43}\bar{N}_3 - a_{44}N_4 + a_{43}N_3\} \end{aligned} \quad (5.5)$$

$$\begin{aligned} = & (N_1 - \bar{N}_1) \{-a_{11}(N_1 - \bar{N}_1) - a_{12}(N_2 - \bar{N}_2)\} + l_1 (N_2 - \bar{N}_2) \{-a_{22}(N_2 - \bar{N}_2) + a_{21}(N_1 - \bar{N}_1) + a_{23}(N_3 - \bar{N}_3)\} \\ & + l_2 (N_3 - \bar{N}_3) \{-a_{33}(N_3 - \bar{N}_3) + a_{34}(N_4 - \bar{N}_4)\} + l_3 (N_4 - \bar{N}_4) \{-a_{44}(N_4 - \bar{N}_4) + a_{43}(N_3 - \bar{N}_3)\} \end{aligned} \quad (5.6)$$

$$\begin{aligned} \frac{dV}{dt} = & -a_{11}(N_1 - \bar{N}_1)^2 - a_{12}(N_1 - \bar{N}_1)(N_2 - \bar{N}_2) \\ & + l_1 \left\{ (-a_{22})(N_2 - \bar{N}_2)^2 + a_{21}(N_1 - \bar{N}_1)(N_2 - \bar{N}_2) + a_{23}(N_2 - \bar{N}_2)(N_3 - \bar{N}_3) \right\} \\ & + l_2 \left\{ (-a_{33})(N_3 - \bar{N}_3)^2 + a_{34}(N_3 - \bar{N}_3)(N_4 - \bar{N}_4) \right\} \\ & + l_3 \left\{ (-a_{44})(N_4 - \bar{N}_4)^2 + a_{43}(N_3 - \bar{N}_3)(N_4 - \bar{N}_4) \right\} \end{aligned} \quad (5.7)$$

Choosing  $l_1 = \frac{a_{12}}{a_{21}}$ ,  $l_2$  and  $l_3$  are any positive constants, (5.7) becomes,

$$\frac{dV}{dt} = -a_{11}(N_1 - \bar{N}_1)^2 - \frac{a_{12}a_{22}}{a_{21}}(N_2 - \bar{N}_2)^2 + \frac{a_{12}a_{23}}{a_{21}}(N_2 - \bar{N}_2)(N_3 - \bar{N}_3) - l_2 a_{33}(N_3 - \bar{N}_3)^2 \quad (5.8)$$

$$\begin{aligned} & + l_2 a_{34}(N_3 - \bar{N}_3)(N_4 - \bar{N}_4) - l_3 a_{44}(N_4 - \bar{N}_4)^2 + l_3 a_{43}(N_3 - \bar{N}_3)(N_4 - \bar{N}_4) \\ & < -a_{11}(N_1 - \bar{N}_1)^2 - \frac{a_{12}a_{22}}{a_{21}}(N_2 - \bar{N}_2)^2 + \frac{a_{12}a_{23}}{2a_{21}} \left\{ (N_2 - \bar{N}_2)^2 + (N_3 - \bar{N}_3)^2 \right\} \end{aligned} \quad (5.9)$$


$$\begin{aligned} & -a_{33}l_2(N_3 - \bar{N}_3)^2 - a_{44}l_3(N_4 - \bar{N}_4)^2 + \frac{(a_{34}l_2 + a_{43}l_3)}{2} \left\{ (N_3 - \bar{N}_3)^2 + (N_4 - \bar{N}_4)^2 \right\} \\ & < -a_{11}(N_1 - \bar{N}_1)^2 + \left[ \frac{a_{23}}{2} - a_{22} \right] \frac{a_{12}}{a_{21}}(N_2 - \bar{N}_2)^2 + \left[ \frac{(a_{34}l_2 + a_{43}l_3)}{2} - \frac{a_{12}a_{23}}{2a_{21}} - a_{33}l_2 \right] (N_3 - \bar{N}_3)^2 \end{aligned} \quad (5.10)$$

$$\begin{aligned} & + \left[ \frac{(a_{34}l_2 + a_{43}l_3)}{2} - a_{44}l_3 \right] (N_4 - \bar{N}_4)^2 \\ & < 0, \text{ Provided } \frac{a_{23}}{2} < a_{22}, \frac{(a_{34}l_2 + a_{43}l_3)}{2} < \frac{a_{12}a_{23}}{2a_{21}} + a_{33}l_2 \text{ and } \frac{(a_{34}l_2 + a_{43}l_3)}{2} < a_{44}l_3 \end{aligned}$$

Hence the co-existent is globally asymptotically stable.

**6. NUMERICAL EXAMPLES:**


(1) Let  $a_1=1, a_{11}=0.1, a_{12}=0.1, a_{23}=0.1, a_2=1, a_{22}=0.2, a_{21}=0.2, a_3=1, a_{33}=0.3, a_{34}=0.2, a_4=1, a_{44}=0.3, a_{43}=0.3, N_1=20, N_2=30, N_3=40, N_4=50$



**Fig.1**

**Fig.1:** Graph Represents Variations in the growth rate of the populations against time.

(2) Let  $a_1=2, a_{11}=0.1, a_{12}=0.02, a_{23}=0.1, a_2=2, a_{22}=0.4, a_{21}=0.3, a_3=1, a_{33}=0.3, a_{34}=0.001, a_4=2, a_{44}=0.2, a_{43}=0.001, N_1=40, N_2=30, N_3=20, N_4=50$



**Fig.2**

**Fig. 2:** Graph Represents Variations in the growth rate of the populations against time.

**REFERENCES:**

- [1] R. Archana Reddy, on the stability of some mathematical models in biosciences- interacting species, Ph. D thesis, 2009, JNTU.
- [2] H.I. Freedman, Deterministic Mathematical Models in Population Ecology, Marcel – Decker, New York, 1980.
- [3] J.N. Kapur, Mathematical Modeling, Wiley – Eastern, 1988.
- [4] J.N. Kapur, Mathematical Models in Biology and Medicine Affiliated East – West, 1985.
- [5] K. Lakshmi Narayan, A Mathematical study of Prey-Predator Ecological Models with a partial covers for the prey and alternative food for the predator, Ph. D thesis, 2004, J. N. T. University.

- [6] A.j. Lotka, Elements of Physical biology, Williams and Wilkins, Baltimore, 1925.
- [7] W.J. Meyer, Concepts of Mathematical Modeling, Mc Graw – Hill, 1985.
- [8] Paul Colinvaux, Ecology, John Wiley and Sons Inc., New York, 1986.
- [9] B. Ravindra Reddy, K. Lakshmi Narayan and N.Ch. Pattabhiramacharyulu, A model of two mutually interacting species with limited resources and a time delay, Advances in Theoretical and Applied Mathematics, Vol.5, No.2 (2010), pp. 121-132.
- [10] B. Ravindra Reddy, K. Lakshmi Narayan and N.Ch. Pattabhiramacharyulu, A model of two mutually interacting species with limited resources and harvesting of both the species at a constant rate, International J. of Math. Sci.& Engg. Appls. (IJMSEA), Vol. 4, No. III (August, 2010), pp. 97-106.
- [11] N.C. Srinivas, Some Mathematical aspects of modeling in Bio Medical Sciences, Ph. D thesis, 1991, Kakatiya University.
- [12] V. Volterra, Leconsen la theorie mathematique de la leitte pou lavie, Gauthier – Villars, Paris, 1931.

\*\*\*\*\*