

ANTI Q-FUZZY FINITE DIMENSIONAL G – MODULAR DISTRIBUTIVE LATTICES

Dr. R. NAGARAJAN*

Associate Professor & Head, Department of Mathematics
J.J. College of Engineering & Technology, Tiruchirappalli- 620009, Tamilnadu, India
E-mail: nagalogesh@yahoo.co.in

(Received on: 30-03-12; Accepted on: 19-04-12)

ABSTRACT

We study the characterization of fuzzy G-modular distributive lattice such as ring sums, lattice homomorphism and upper level sets and characterization of these lattices which is similar to a well known result of lattice theory. The main goal of this paper is to study the finite groups whose lattices of fuzzy G-modular are distributive.

Key Words: Fuzzy lattices, level set, G-modular, anti Q-Fuzzy G-modular lattices, fuzzy filter, fuzzy ideal, lattice homomorphism.

SECTION-1: INTRODUCTION

The theory of fuzzy sets which was introduced by L. A. Zadeh [14] is applied to many mathematical branches. Rosenfeld [14] inspired the fuzzification of algebraic structures and introduced the notion of fuzzy sub groups. P. Das [6] characterized fuzzy sub groups by their level sub groups. A.Solairaju and R.Nagarajan introduced the notion of Q-fuzzy groups [15]. In [11] Liu applied the concept of fuzzy sets to the theory of rings and introduced and examined the notion of a fuzzy ideal of a ring. The study of fuzzy sub module was introduced by pan and Golan in [7]. Also Pan [13] studied the fuzzy finitely generated modules and fuzzy quotient modules. Later Katsaras and D.B. Liu introduced the concept of fuzzy vector spaces and fuzzy topological vector spaces. The formation of a lattice of sub modules of a module is well known features in classical algebra. However the same has not been explored in fuzzy setting. In order to initiate such studies the concept of fuzzy sub-module generated by an arbitrary fuzzy set is formulate in this note. Using this concept S.K. Bambri and Pratibakumar [5] introduced Lattice of fuzzy sub modules and established an embedding of the lattice of all sub module of a module M into the lattice of fuzzy sub module.

Q. Zhang and Meng [17] considered the sub property to be an assumption and established the more general result that “the lattice $l_t(R)$ of all fuzzy ideals of a ring R with the same tip “t” is modular. On the other hand in [11] the authors have arrived at the false result that The lattice $l(R)$ is distributive. In fact the lattice $I(R)$ of ideals of a ring R is not distributive and has an obvious embedding in $l(R)$. The corrected result has already appeared in several papers [10, 16]. I Jahan in [8], by constructing and employing the technique of strong level subsets, he proved that the lattice (R) of all ideals of a ring R is modular. This proof of the modularity of $l(R)$ is different from the Ajmal’s proof of modularity of the lattice of fuzzy normal sub groups of group appeared in [1]. K. C. Gupta and S. Roy in introduce the indirect proof of the above result that the modularity of the quasi Hamiltonians fuzzy sub groups. Hence N. Ajmal and K.V. Thomas in [2] initiated a discussion on the aspect of modularity of the set of fuzzy normal sub groups. A special class of fuzzy normal sub groups of group has been shown to constitute a modular sub lattice of its fuzzy sub group lattice. They first established that the sub lattice (G) is modular. Moreover using the same technique in [3], they demonstrate that whenever the set (G) of all fuzzy quasi normal sub groups of a given group forms a lattice, the lattice is modular. B. Yuan and Wu. Wangning in [4], introduced fuzzy ideals on a distributive lattice. In this paper, we study the characterization of fuzzy G-modular distributive lattice such as ring sums, lattice homomorphism and upper level sets and characterization of these lattices which is similar to a well known result of lattice theory

SECTION- 2: PRELIMINARIES

Definition: 2.1 A non empty set L together with two binary operations \vee and \wedge on L is called a lattice if it satisfies the following identities.

L1: $x \vee y \approx y \vee x, x \wedge y \approx y \wedge x$

L2: $x \vee (y \vee z) \approx (x \vee y) \vee z, x \wedge (y \wedge z) \approx (x \wedge y) \wedge z$

L3: $x \vee x \approx x, x \wedge x \approx x$

Corresponding author: Dr. R. NAGARAJAN, *E-mail: nagalogesh@yahoo.co.in

L4: $x \approx x \vee (x \wedge y), x \approx x \wedge (x \vee y)$. The operation \wedge is called meet and the operation \vee is called join. Let L be the set of proposition and let \vee denote the connective “or” and \wedge denote the connective “and”. Then L1 to L4 are well known properties from propositional logic.

Definition: 2.2 A distributive lattice is a lattice which satisfies either of the distributive laws. D1. $x \wedge (y \vee z) \approx (x \wedge y) \vee (x \wedge z)$, D2. $x \vee (y \wedge z) \approx (x \vee y) \wedge (x \vee z)$. One can see a lattice L satisfies D1 if and only if satisfies D2.

Definition: 2.3 Let G be a group and L be a vector distributive lattice then L is called G-modular if for $a \in G$ and $l \in L$, there exist a product $a \cdot l \in L$ satisfies the following axioms.

- (i) $e \cdot l = l$, for all $l \in L$
- (ii) $(a \cdot h) \cdot l = a \cdot (h \cdot l)$
- (iii) $a \cdot (k_1 m_1 + k_2 m_2) = k_1 (a m_1) + k_2 (a m_2)$ for all $k_1, k_2 \in K$ and $m_1, m_2 \in M$.

Definition: 2.4 A mapping $\mu: X \times Q \rightarrow [0, 1]$ where X is an arbitrary non-empty set and is called a Q-fuzzy set in X.

Definition: 2.5 Let X be a Q-fuzzy set and $\mu: X \times Q \rightarrow G$ be a lattice ordered group of X. Then μ is called Q-fuzzy lattice ordered group (QFLG) if

- (i) $\mu(x + y, q) \geq \min \{ \mu(x, q), \mu(y, q) \}$
- (ii) $\mu(-x, q) \geq \mu(x, q)$
- (iii) $\mu(0, q) = 1$, for all $x, y \in G$.

Definition: 2.6 Let μ be a Q-fuzzy lattice ordered group of G and $\mu: X \times Q \rightarrow G$. Then μ is called Q-fuzzy lattice if

- (i) $\mu(x + y, q) \geq \min \{ \mu(x, q), \mu(y, q) \}$
- (ii) $\mu(-x, q) \geq \mu(x, q)$
- (iii) $\mu(x \vee y) \geq \min \{ \mu(x), \mu(y) \}$
- (iv) $\mu(x \wedge y) \geq \min \{ \mu(x), \mu(y) \}$, for all $x, y \in G$.

Definition: 2.7 Let X be any group and L be a vector distributive lattice extended real valued functions on X. If μ is called a anti Q-fuzzy G-modular lattice on L then it satisfies the following conditions.

- (i) $\mu(ax + by, q) \leq \max \{ \mu(x), \mu(y) \}$
- (ii) $\mu(gx, q) \leq \mu(x)$
- (iii) $\mu(x \vee y)_q \wedge \mu(x \wedge y)_q \leq \max \{ \mu(x, q), \mu(y, q) \}$, for all $x, y \in L$.

Definition: 2.8 A Q-fuzzy sub set μ is called monotonic if

- (i) $\mu(x, q) \leq \mu(y, q)$ whenever $x \leq y$

Definition: 2.9 Let μ be any Q-fuzzy sub set of X. Then the set

$\mu_t = \{ x \in X / \mu(x, q) \geq t, t \in [0, 1] \}$ is called a level sub set of μ .

Definition: 2.10

- (i) A monotonic fuzzy G-modular lattice is called a fuzzy filter of L.
- (ii) A anti-monotonic fuzzy G-modular lattice is called a fuzzy ideal of L.
- (iii) A Q-fuzzy sub set μ is called a Q-fuzzy filter if and only if $\mu(x \wedge y)_q = \mu(x, q) \wedge \mu(y, q)$, for all $x, y \in L$.
- (iv) A Q-fuzzy subset μ is called a Q-fuzzy ideal if and only if $\mu(x \vee y)_q = \mu(x, q) \wedge \mu(y, q)$, for all $x, y \in L$.

Definition: 2.11

- (i) A Q-fuzzy filter is prime if $\mu(x \vee y)_q \leq S \{ \mu(x, q), \mu(y, q) \}$

- (ii) A Q-fuzzy ideal is prime if $\mu(x \wedge y)_q \leq S \{ \mu(x, q), \mu(y, q) \}$, holds for all $x, y \in L$.

Definition: 2.12 A Q-fuzzy G-modular lattice A is modular if and only if

$A(x, z) \geq 0 \Rightarrow x \vee (y \wedge z) = (x \vee y) \wedge z$. for all x, y and $z \in X$.

Definition: 2.13 Let μ be a anti Q-fuzzy G-modular lattice. If μ is distributive then

- (i) $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$ and
- (ii) $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$ for all x, y and $Z \in X$.

Definition: 2.14 Let L and L' be lattices. A mapping $f: L \rightarrow L'$ is said to be lattice homomorphism if

- (i) $f(x + y) = f(x) + f(y)$ and
- (ii) $f(xy) = f(x) \cdot f(y)$, for all $x, y \in L$.

Definition: 2.15 We say that a Anti Q-fuzzy G-modular distributive lattice $M = (L, \mu_M)$ in L satisfies the imaginable property if

$$I_m(\mu_M) \subseteq \Delta T$$

Definition: 2.16 Let λ and μ be two anti Q-fuzzy G-modular distributive lattice of L then the sum of λ and μ are denoted by $\lambda + \mu$ and is defined as

$$(\lambda + \mu)(z, q) = \inf_{z = x + y} \{ \max \{ \lambda(x, q), \mu(y, q) \} \} \quad \text{where } x, y \in L.$$

Clearly $\lambda + \mu$ is a anti Q-fuzzy G modular distributive lattice in L .

Definition: 2.17 Let λ and μ be two anti Q-fuzzy G-modular distributive lattice of L , Then the ring sum of λ and μ is defined as

$$(\lambda \oplus \mu)(z, q) = \inf_{Z \in \lambda \oplus \mu} \{ (\mu_1 + \mu_2)(z, q) / x = f^{-1}(\lambda, q), y = f^{-1}(\mu, q) \} \text{ for all } x, y \text{ and } z \in L$$

SECTION -3: CHARACATERIZATION OF ANTI Q-FUZZY G-MODULAR DISTRUBUTIVE LATTICES

Throughout this paper G be a finite groups and $L = \langle L, +, \cdot \rangle$ denotes a lattice.

That is of maps from L into $\langle [0, 1]; V, \wedge \rangle$, where $[0, 1]$ is the set of real's between 0 and 1. and

- (i) $x \vee y = \max(x, y)$, if $y \leq x$ then $\mu(y) \geq \mu(x)$ and
- (ii) $x \wedge y = \min(x, y)$, if $x \leq y$ then $\mu(x) \geq \mu(y)$

Proposition: 3.1 Let M be a G-modular lattice and N be a anti Q-fuzzy G-sub modular lattice of M has a anti Q-fuzzy G-modular distributive lattice then M/N and N has anti Q-fuzzy G-modular distributive lattice.

Proof: Let μ be a fuzzy sub modular lattice of M then $V: \mu_N$ is anti Q-fuzzy G-modular lattice on N .

Define $\mu: \frac{M}{N} \times Q \rightarrow [0, 1]$ by $\mu(x + N, q) = \mu(x, q)$, for all $x \in G$ and $n \in \frac{M}{N}$

- (1) $\mu(a(x + N) + b(y + N), q) = \mu((ax + by) + N, q)$
 $= \mu(ax + by, q)$
 $\leq \max \{ \mu(x, q) + \mu(y, q) \}$
 $\leq \max \{ \mu(x + N, q) + \mu(y + N, q) \}$
- (2) $\mu(g(x + N, q)) = \mu(gx + N, q) = \mu(gx, q)$
 $\leq \mu(x, q)$
 $\leq \mu(x + N, q)$
- (3) $(\mu(x \vee y)q \wedge \mu(x \wedge y)q) + N = \max \{ \mu((x \vee y)q + N), \mu(x \wedge y)q + N \}$
 $\leq \max \{ \max \{ (\mu(x, q), \mu(y, q)) + N \}, \{ \max \{ (\mu(x, q), \mu(y, q)) + N \} \}$
 $\leq \max \{ \mu(x, q) + N, \mu(y, q) + N \}$
 $\leq \max \{ \mu(x + N, q), \mu(y + N, q) \}$
 $\leq \max \{ \mu(x, q), \mu(y, q) \}$, for all $x, y \in G$

Therefore $\frac{M}{N}$ and N are anti Q-fuzzy G-modular distributive lattice.

Proposition: 3.2 Let $f: L \rightarrow L^*$ be a G-modular distributive lattice homomorphism. Where L and L^* are G-modular distributive lattices. If V is anti Q-fuzzy G –modular distributive lattice on L^* then $f^{-1}(v)$ is anti Q- fuzzy G-modular distributive lattice on L.

Proof: Since V is anti Q-fuzzy G-modular distributive lattice on L^* and $f: L \rightarrow L^*$ be a G-modular lattice homomorphism. For any $a, b \in K$ and $x, y \in L$, we have

$$\begin{aligned}
 (1) \quad f^{-1}(V)(ax + by, q) &= V(f(ax + by, q)) \\
 &= V(f(ax, q) + f(by, q)) \\
 &\leq V\{(f(x, q) + f(y, q))\} \\
 &\leq \max\{V f(x, q), V f(y, q)\} \\
 &\leq \max\{f^{-1}(V)(x, q), f^{-1}(V)(y, q)\} \\
 (2) \quad f^{-1}(V)(gm, q) &= V f(gm, q) \\
 &\leq V f(m, q) \\
 &\leq f^{-1}(V)(m, q) \\
 (3) \quad f^{-1}(V)(x \vee y)q \wedge f^{-1}(V)(x \wedge y)q &= \max\{f^{-1}(V)((x \vee y)q), f^{-1}(V)(x \wedge y)q\} \\
 &= \max\{V f(x \vee y, q), V f(x \wedge y, q)\} \\
 &= \max\{V(f(x, q) + f(y, q)), V(f(x, q) \cdot f(y, q))\} \text{(since V is homomorphism)} \\
 &\leq \max\{\max\{V f(x, q), V f(y, q)\}, \{\max\{V f(x, q), V f(y, q)\}\} \\
 &\leq \max\{V f(x, q), V f(y, q)\} \\
 &\leq \max\{f^{-1}(v)(x, q), f^{-1}(v)(y, q)\}
 \end{aligned}$$

Therefore $f^{-1}(V)$ is anti Q- fuzzy G-modular distributive lattice on L.

Proposition: 3.3 Let S be a t-norm, then every imaginable anti Q-fuzzy G-modular distributive lattice of L is anti Q-fuzzy G-modular distributive lattice.

Proof: Let $M = (L, \mu_M)$ be an imaginable anti Q-fuzzy G-modular distributive lattice of L under S-norms. Then

- (i) $\mu_M(ax + by, q) \leq S\{\mu_M(x, q), \mu_M(y, q)\}$
- (ii) $\mu_M(gm, q) \leq \mu_M(m, q)$,
- (iii) $\mu_M(x \vee y)q \wedge \mu_M(x \wedge y)q \leq S\{\mu_M(x, q), \mu_M(y, q)\}$ for all $a, b \in G, x, y$ and $m \in M$.

We have

$$\begin{aligned}
 \max\{\mu_M(x, q), \mu_M(y, q)\} &= S\{\max\{\mu_M(x, q), \mu_M(y, q)\}, \max\{\mu_M(x, q), \mu_M(y, q)\}\} \\
 &\geq S\{\mu_M(x, q), \mu_M(y, q)\} \\
 &\geq \max\{\mu_M(x, q), \mu_M(y, q)\}
 \end{aligned}$$

It follows that $\mu_M(ax + by, q) \leq S\{\mu_M(x, q), \mu_M(y, q)\}$

Therefore $M = (L, \mu_M)$ is an imaginable anti Q-fuzzy G-modular distributive lattice of L.

Proposition: 3.4 If L is a complete lattice then the intersection of a family of anti Q- fuzzy G-modular distributive lattice is a fuzzy G-modular distributive lattice.

Proof: Let $\{M_j : j \in J\}$ be a family of anti Q-fuzzy G-modular distributive lattice and let $M = \bigcap M_j$

We have $\mu_M(x, q) = \sup \mu_{M_j}(x, q)$ and $\mu_M(y, q) = \sup \mu_{M_j}(y, q)$

$$\begin{aligned} (1) \quad \mu_M(ax + by, q) &= \sup \mu_{M_j}(ax + by, q) \\ &\leq \text{Sup} \{ \max \{ \mu_{M_j}(x, q), \mu_{M_j}(y, q) \} \} \\ &\leq \text{Max} \{ \sup \mu_{M_j}(x, q), \sup \mu_{M_j}(y, q) \} \\ &\leq \text{Max} \{ \mu_M(x, q), \mu_M(y, q) \} \end{aligned}$$

$$\begin{aligned} (2) \quad \mu_M(gx, q) &= \sup \mu_{M_j}(gx, q) \\ &\leq \text{Sup} \mu_{M_j}(x, q) \\ &\leq \mu_M(x, q) \end{aligned}$$

$$\begin{aligned} (3) \quad \mu_M(x \vee y)q \wedge \mu_M(x \wedge y)q &= \max \{ \mu_M((x \vee y), q), \mu_M(x \wedge y, q) \} \\ &= \max \{ \sup \mu_{M_j}(x \vee y, q), \sup \mu_{M_j}(x \wedge y, q) \} \\ &= \max \{ \sup \max \{ \mu_{M_j}(x, q), \mu_{M_j}(y, q) \}, \text{Sup} \max \{ \mu_{M_j}(x, q), \mu_{M_j}(y, q) \} \} \\ &\leq \max \{ \sup \mu_{M_j}(x, q), \sup \mu_{M_j}(y, q) \} \\ &\leq T \{ \mu_M(x, q), \mu_M(y, q) \}, \text{ for all } x, y \in L. \end{aligned}$$

Therefore $\bigcap \mu_{M_j}$ is a anti Q-fuzzy G-modular distributive lattice.

Proposition: 3.5 Any finite dimensional G-modular distributive lattice over a sub lattice L is anti Q- fuzzy G-modular distributive lattice over L.

Proof: Let $\mu: M \rightarrow [0, 1]$ be a map and M be a finite n-dimensional anti Q-fuzzy G-modular distributive lattice over L.

For all $\alpha, \beta \in L$ and $x, y \in M$

Such that $\mu^n(\alpha x, q) = \mu(n(\alpha x), q) = \mu^n(x, q)$

$$\begin{aligned} (1) \quad \mu^n(\alpha x + \beta y, q) &= \mu(n(\alpha x + \beta y), q) \\ &= n \mu(\alpha x + \beta y, q) \\ &\leq n \{ \max \{ \mu(\alpha x, q) + \mu(\beta y, q) \} \} \\ &\leq \max \{ n\mu(\alpha x, q), n\mu(\beta y, q) \} \\ &\leq \max \{ \mu^n(\alpha x, q), \mu^n(\beta y, q) \} \\ &\leq \max \{ \mu^n(x, q), \mu^n(y, q) \} \end{aligned}$$

$$\begin{aligned} (2) \quad \mu^n(gx, q) &= \mu(n(gx), q) \\ &= n\mu(gx, q) \\ &\leq n\mu(x, q) \\ &\leq \mu^n(x, q) \end{aligned}$$

$$\begin{aligned}
 (3) \mu^n(x \vee y)_q \wedge \mu^n(x \wedge y)_q &= \max \{ \mu^n((x \vee y, q), \mu^n(x \wedge y, q)) \} \\
 &= \max \{ \mu(n(x \vee y, q), \mu(n(x \wedge y, q)) \} \\
 &= \max \{ n\mu\{(x \vee y, q), n\mu(x \wedge y, q)\} \} \\
 &\leq \max \{ n\{\max \{ \mu(x, q), \mu(y, q)\}, n\{\max \{ \mu(x, q), \mu(y, q)\} \} \} \\
 &\leq \max \{ n\mu(x, q), n\mu(y, q) \} \\
 &\leq \max \{ \mu^n(x, q), \mu^n(y, q) \}, \text{ for all } x, y \in M.
 \end{aligned}$$

Therefore, n-dimensional anti Q-fuzzy G-modular distributive lattice is anti Q- fuzzy G-modular distributive lattice.

Proposition: 3.6 If M and N are anti Q-fuzzy G-modular distributive lattices over a fuzzy sub-lattice L then $L = M \oplus N$ is anti Q- fuzzy G-modular distributive lattice on L. Here \oplus is referred as a ring sum.

Proof: Let M and N be two anti Q-fuzzy G-modular distributive lattices of μ_1 and μ_2 respectively. Then we define the ring sum of M. and N as

$$(M \oplus N)(z, q) = \inf_{z \in M \oplus N} \{ (\mu_1 + \mu_2)(z, q) / x = f^{-1}(\mu_1, q), y = f^{-1}(\mu_2, q) \}$$

Now

$$\begin{aligned}
 (1) (M \oplus N)(\alpha x + \beta y, q) &= (\mu_1 + \mu_2)(\alpha x + \beta y, q) \\
 &\leq \max \{ ((\mu_1 + \mu_2)(x, q), (\mu_1 + \mu_2)(y, q)) \} \\
 &\leq \max \{ \inf_{z \in M \oplus N} \mu_1(z, q), \inf_{z \in M \oplus N} \mu_2(z, q) \} \\
 &\leq \max \{ (M \oplus N)(z, q), M \oplus N(z, q) \} \\
 &\leq \max \{ (M \oplus N)(x, q), M \oplus N(y, q) \}
 \end{aligned}$$

$$\begin{aligned}
 (2) (M \oplus N)(gz, q) &= (\mu_1 + \mu_2)(gz, q) \\
 &\leq (\mu_1 + \mu_2)(z, q) \\
 &\leq \inf \{ \mu_1 + \mu_2 \}(z, q) \\
 &z \in M \oplus N \\
 &\leq M \oplus N(z, q)
 \end{aligned}$$

$$\begin{aligned}
 (3) (M \oplus N)(x \vee y)_q \wedge (M \oplus N)(x \wedge y)_q &= \max \{ (M \oplus N)(x \vee y, q), (M \oplus N)(x \wedge y, q) \} \\
 &= \max \{ (\mu_1 + \mu_2)(x \vee y, q), (\mu_1 + \mu_2)(x \wedge y, q) \} \\
 &\leq \max \{ \max \{ (\mu_1 + \mu_2)(x, q), (\mu_1 + \mu_2)(y, q) \}, \max \{ (\mu_1 + \mu_2)(x, q), (\mu_1 + \mu_2)(y, q) \} \} \\
 &\leq \max \{ (\mu_1 + \mu_2)(x, q), (\mu_1 + \mu_2)(y, q) \} \\
 &\leq \max \{ \inf_{z \in M \oplus N} \mu_1(z, q), \inf_{z \in M \oplus N} \mu_2(z, q) \} \\
 &\leq \max \{ (M \oplus N)(x, q), M \oplus N(y, q) \}
 \end{aligned}$$

Therefore, $M \oplus N$ is anti Q- fuzzy G-modular distributive lattices.

Proposition: 3.7 Let M be a G -modular distributive lattices over a sub lattice L and $M = \sum_{i=1}^n M_i$, where M_i 's are G sub modular distributive lattice of M . If V_i are anti Q -fuzzy G -modular distributive lattice of M_i then $V : M \rightarrow [0, 1]$ is anti Q -fuzzy G -modular distributive lattice in L .

Proof: Since V_i is anti Q - fuzzy G -modular distributive lattice on M_i for every $x, y \in M$; $g \in L$ and $\alpha, \beta \in L$. We have

$$\begin{aligned} (1) \quad V(\alpha x + \beta y, q) &= V(\sum(\alpha M_i + \beta M_i', q)) \\ &= \wedge (V_i(\alpha M_i + \beta M_i', q)), \quad \text{where } i = 1, 2 \dots n. \\ &= V_j(\alpha M_j + \beta M_j', q), \text{ for some } j \\ &\leq \max(V_j(M_j, q), V_j(M_j', q)) \\ &\leq \max(V(x, q), V(y, q)) \end{aligned}$$

$$\begin{aligned} (2) \quad V(gx, q) &= V(\sum gM_i, q) \\ &= \wedge (V_i(gM_i, q), \quad \text{where } i = 1, 2 \dots n. \\ &= V_j(gM_j, q), \text{ for some } j \\ &\leq V_j(M_j, q) \\ &\leq V(x, q) \end{aligned}$$

$$\begin{aligned} (3) \quad V(x \vee y, q) \wedge V(x \wedge y, q) &= \max\{V((x \vee y, q), V(x \wedge y, q) \\ &= \max\{V(\sum(x \vee y, q)M_i, V(\sum(x \wedge y, q)M_i') \text{ for } i = 1, 2, \dots n \\ &= \max\{V(x \vee y, q)M_i, V(x \wedge y, q)M_i'\} \\ &\leq \max\{\max\{V(x, q), V(y, q)\}M_i, \{\max\{V(x, q), V(y, q)\}M_i'\} \\ &\quad V(y, q)M_i'\} \\ &\leq \max\{V_i M_j, V_j M_j'\}, \text{ for some } j. \\ &\leq \max\{V(x, q), V(y, q)\} \end{aligned}$$

Therefore V is anti Q - fuzzy G -modular distributive lattice on anti Q - fuzzy G -modular distributive lattice in L .

Proposition 3.8: Let λ and μ be anti Q -fuzzy G -modular distributive lattices of L , then $\lambda + \mu$ is the smallest anti Q -fuzzy G -modular distributive lattice of L .

Proof: For any $x, y, h, f \in L$, we have

$$\begin{aligned} (1) \quad \max\{(\lambda + \mu)_q x, (\lambda + \mu)_q y\} &= \max\{\inf_{x=a+b} \max\{(\lambda(a, q), \mu(b, q)), \inf_{y=c+d} \max\{(\lambda(c, q), \mu(d, q))\} \\ &= \max\{\inf_{x=a+b} (\max\{(\lambda(a, q), \mu(b, q)), \max\{(\lambda(c, q), \mu(d, q))\} \\ &= \max\{\inf_{x=a+b} \max\{(\lambda(a, q), \mu(b, q), \lambda(c, q), \mu(d, q))\} \\ &\geq \max\{\inf_{x=a+b} \max(\lambda(a+(b-c-b), q), \mu(b+(c-d-c), q))\} \end{aligned}$$

$$\geq \max \{ \inf \max \{ (\lambda(h, q), \lambda(f, q)) \} \\ x + y = h + f \\ \geq (\lambda + \mu)q(x + y)$$

Therefore $(\lambda + \mu)(x + y) \leq S \{ (\lambda + \mu)q(x), (\lambda + \mu)q(y) \}$

$$(2) (\lambda + \mu)(gx, q) = \inf \max \{ (\lambda(a, q), \mu(b, q)) \\ x = a + b \\ = \inf \max \{ (\lambda(a, q), \mu(b, q)) \\ g_x = g_a + g_b \\ \leq \inf \max \{ (\lambda(a, q), \mu(b, q)) \\ x = G_a + G_b \\ \leq (\lambda + \mu)(x, q)$$

$$(3) (\lambda + \mu)(x \vee y)q \wedge (\lambda + \mu)(x \wedge y)q = \max \{ (\lambda + \mu)(x \vee y)q, (\lambda + \mu)(x \wedge y)q \} \\ = S \{ \max \{ (\lambda + \mu)(x, q), (\lambda + \mu)(y, q) \}, \max \{ (\lambda + \mu)(x, q), (\lambda + \mu)(y, q) \} \} \\ = S \{ \max \{ \inf \max \{ \lambda(x, q), \mu(x, q) \} x = a + b \inf \max \{ \lambda(y, q), \mu(y, q) \}, \\ y = c + d \\ = \max \{ \inf \max \{ \lambda(x, q), \mu(x, q) \} x = a + b \inf \max \{ \lambda(y, q), \mu(y, q) \} \\ y = c + d \\ \leq S \{ \inf \max \{ \lambda(y, q), \mu(y, q) \}, \inf \max \{ \lambda(x, q), \mu(x, q) \} \\ y = c + d x = a + b \\ \leq S \{ \inf \max \{ \lambda(x, q), \mu(x, q) \}, \inf \max \{ \lambda(y, q), \mu(y, q) \} \\ x = a + b y = c + d \\ \leq S \{ (\lambda + \mu)(x, q), (\lambda + \mu)(y, q) \}$$

Therefore $(\lambda + \mu)(x \vee y)q \wedge (\lambda + \mu)(x \wedge y)q \leq S \{ (\lambda + \mu)(x, q), (\lambda + \mu)(y, q) \}$

Hence $\lambda + \mu$ is anti Q- fuzzy G-modular distributive lattice on L.

CONCLUSIONS:

I. Jahan introduce the concept of modularity of Ajmal for the lattices of fuzzy ideals of a ring. In this paper we study the characterization of anti Q-fuzzy G-modular distributive lattices. One can obtain the similar results in soft modules and soft lattices by using a suitable mathematical tool.

REFERENCES:

1. N. Ajmal, The "Lattice of fuzzy normal sub groups is modular", Inform. Sci. 83 (1995), pp. 199-209.
2. N. Ajmal, K.V. Thomas, "A complete study of the lattices of fuzzy congruences and fuzzy normal sub groups, Inform. Sci. 82 (1995), pp. 198-218.
3. S-Z. Bai, "Pre-semi compact L-sub sets in fuzzy lattices". International Journal of fuzzy systems, Vol.12, No.1, pp.59-65, 2010.
4. Y. Bo and Wu. Wangning, "Fuzzy ideals on a distributive lattice", Fuzzy sets and systems. Vol. 81 (1996), pp.383 – 393.
5. S.K. Bhambri, Pratibha Kumar, "Lattice of fuzzy sub modules", International mathematical forum, 4 (2009) No.13, pp.653-659.
6. P. Das, "Fuzzy groups and level sub groups", J. Math. Anal. Appl., Vol.85 (1981), pp.264-269.
7. Golan. J.S., "Making Modules fuzzy", Fuzzy sets and systems, 32 (1989), pp.91-94.
8. I. Jahan, "Modularity of Ajmal for the lattices of fuzzy ideals of a ring", Iranian Journal of fuzzy systems, Vol.5, No.2 (2008), pp. 71 – 78.

9. M. Marudai and V. Rajendran, "New structure of G- modular distributive lattice" IJFMS,1(2),181-196, 2011.
10. R. Kumar, "Non distributivity of the lattice of fuzzy ideals of a ring R", Fuzzy sets and systems, 97 (1998), pp.393-394.
11. W.J. Liu, "Fuzzy invariant sub groups and fuzzy ideals, "Fuzzy sets and systems, Vo.8 (1982), pp.133-139.
12. Majumdar, Sultana, "The lattice of fuzzy ideals of a ring R", Fuzzy sets and systems, 81 (1996), pp.271-273.
13. Pan, Fu-zheng, "Fuzzy finitely generated modules", Fuzzy sets and systems, 21 (1987), pp. 105-113.
14. A. Rosenfeld, "Fuzzy groups", J. Maths. Anal. Appl. Vol.35 (1971), pp.512-517.
15. A.Solairaju and R.Nagarajan, "A New structure and constructions of Q-fuzzy group" Advances in Fuzzy mathematics, Vol.4., No.1 (2009), PP 23-29.
16. L.A. Zadeh, "Fuzzy sets", Inform. Control. Vo.8 (1965), pp.338-353.
17. Q. Zhang, Meng, "On the lattice of fuzzy ideals of a ring", Fuzzy sets and systems, 112 (2000) pp. 349-353.
