

## ON EINSTEIN-SASAKIAN PROJECTIVE RECURRENT SPACES

A. K. Singh\*

*Department of Mathematics, H. N. B. Garhwal University (A Central University) Campus,  
 Badshahi Thaul, TEHRI GARHWAL-249 199, Uttarakhand (INDIA)*

*(Received on: 03-05-12; Revised & Accepted on: 27-05-12)*

### ABSTRACT

*In the present paper, we define and study Einstein-Sasakian Projective recurrent spaces and establish several theorems. We investigate the necessary and sufficient condition for an Einstein-Projective (E-P) space to be Sasakian recurrent also Einstein- Sasakian (E-S) space is an Einstein-Projective (E-P) space, iff the scalar curvature is equal to zero.*

**Key words:** Einstein, Kaehlerian, Sasakian space, Conharmonic and Recurrent.

**Subject classification:** 53C20, 53C25.

### 1. INTRODUCTION

An  $n$  –dimensional Sasakian space  $S_n$  (or, normal contact metric space) is an odd dimensional Riemannian space, which admits a unit killing vector field  $\eta^i$  satisfying [4]:

$$\eta_{i,j} = \eta_i g_{jk} - \eta_k g_{ij}, \quad (1.1)$$

where the comma (,) followed by an index denotes the operator of covariant differentiation w.r.t. the metric tensor  $g_{ij}$  of the Riemannian space. Also we have the relation:

$$F_i^h F_h^i = -\delta_i^j \quad (1.2)$$

The Riemannian curvature tensor field  $R_{ijk}^h$  is given by

$$R_{ijk}^h = \partial_j \left\{ \begin{matrix} h \\ i \quad k \end{matrix} \right\} - \partial_k \left\{ \begin{matrix} h \\ i \quad j \end{matrix} \right\} + \left\{ \begin{matrix} m \\ i \quad k \end{matrix} \right\} \left\{ \begin{matrix} h \\ m \quad j \end{matrix} \right\} - \left\{ \begin{matrix} m \\ i \quad j \end{matrix} \right\} \left\{ \begin{matrix} h \\ m \quad k \end{matrix} \right\}, \quad (1.3)$$

where  $\partial_j \equiv \frac{\partial}{\partial x^j}$ .

The Ricci-tensor and scalar curvature are respectively given by  $R_{ij} = R_{ijk}^k$  and  $R = R_{ij} g^{ij}$ .

It is well known that these tensors satisfy the following identities (Tachibana [3]):

$$F_i^a R_a^j = R_i^a F_a^j \quad (1.4)$$

and

$$F_i^a R_{aj} = -R_{ai} F_j^a.$$

In view of (1.1), the relation (1.2) gives

$$F_i^a R_a^b F_b^j = -R_i^j$$

Thus, both  $S_n$  and  $K_n$  are Riemannian spaces, satisfying all the properties of a Riemannian space.

The Sasakian Conharmonic curvature tensor  $T_{ijk}^h$  and the holomorphic Sasakian projective curvature tensor  $P_{ijk}^h$  are respectively given by [3]:

**Corresponding author: A. K. Singh\***

*Department of Mathematics, H. N. B. Garhwal University (A Central University) Campus,  
 Badshahi Thaul, TEHRI GARHWAL-249 199, Uttarakhand (INDIA)*

$$T_{ijk}^h = R_{ijk}^h + \frac{1}{n+4} (R_{ik} \delta_j^h - R_{jk} \delta_i^h + g_{ik} R_j^h - g_{jk} R_i^h + S_{ik} F_j^h - S_{jk} F_i^h + F_{ik} S_j^h - F_{jk} S_i^h + 2S_{ij} F_k^h + 2F_{ij} S_k^h) \quad (1.5)$$

and

$$P_{ijk}^h = R_{ijk}^h + \frac{1}{n+2} (R_{ik} \delta_j^h - R_{jk} \delta_i^h + S_{ik} F_j^h - S_{jk} F_i^h + 2S_{ij} F_k^h), \quad (1.6)$$

where

$$S_{ij} = F_i^a R_{aj}.$$

Let us suppose that a Sasakian space is an Einstein one, then Ricci tensor satisfies

$$R_{ij} = \frac{R}{n} g_{ij}, \quad R_{,a} = 0 \quad (1.7)$$

from which, we have

$$R_{ij,a} = 0, \quad S_{ij,a} = 0 \quad \text{and} \quad S_{ij} = \frac{R}{n} F_{ij}. \quad (1.8)$$

If a Sasakian space is an Einstein one, then the Sasakian Conharmonic curvature tensor and Sasakian projective curvature tensor respectively reduce to the forms:

$$E_{ijk}^h = R_{ijk}^h + \frac{2}{n(n+4)} (g_{ik} \delta_j^h - g_{jk} \delta_i^h + F_{ik} F_j^h - F_{jk} F_i^h + 2F_{ij} F_k^h) \quad (1.9)$$

and

$$*P_{ijk}^h = R_{ijk}^h + \frac{R}{n+2} (g_{ik} \delta_j^h - g_{jk} \delta_i^h + F_{ik} F_j^h - F_{jk} F_i^h + 2F_{ij} F_k^h) \quad (1.10)$$

In view of equations (1.9) and (1.10), we have

$$E_{ijk}^h = *P_{ijk}^h + \frac{R}{(n+2)(n+4)} (g_{ik} \delta_j^h - g_{jk} \delta_i^h + F_{ik} F_j^h - F_{jk} F_i^h + 2F_{ij} F_k^h) \quad (1.11)$$

Now, we shall use the following definition:

**Definition (1.1):** a Sasakian space is said to be recurrent if, we have [1]:

$$R_{ijk,a}^h - \lambda_a R_{ijk}^h = 0, \quad (1.12)$$

for some non-zero recurrence vector  $\lambda_a$  and is called Sasakian Ricci-recurrent, if it satisfies the relation:

$$R_{ij,a} - \lambda_a R_{ij} = 0. \quad (1.13)$$

Multiplying the above equation by  $g^{ij}$ , we get

$$R_{,a} - \lambda_a R = 0. \quad (1.14)$$

**Remark (1.1):** From (1.13), it follows that every Sasakian recurrent space is Sasakian Ricci-recurrent, but the converse is not necessarily true.

**Definition (1.2):** a Sasakian space satisfying the relation [6]:

$$E_{ijk,a}^h - \lambda_a E_{ijk}^h = 0, \quad (1.15)$$

where  $\lambda_a$  is a non-zero recurrence vector, will be called an Einstein-Sasakian conharmonic recurrent, or briefly an  $E - S^* - space$ .

**Definition (1.3):** a Sasakian space is called a space of constant holomorphic sectional curvature tensor [3], if the tensor  $U_{ijk}^h$  defined by

$$U_{ijk}^h = R_{ijk}^h + \frac{R}{(n+2)(n+4)} (g_{ik} \delta_j^h - g_{jk} \delta_i^h + F_{ik} F_j^h - F_{jk} F_i^h + 2F_{ij} F_k^h) \quad (1.16)$$

Vanishes identically.

**Remark (1.2):** In an Einstein-Sasakian space, the curvature tensor  $P_{ijk}^h$  coincides with constant holomorphic sectional curvature.

## 2. EINSTEIN-SASAKIAN PROJECTIVE RECURRENT SPACES

**Definition (2.1):** A Sasakian space satisfying the relation:

$$*P_{ijk,a}^h - \lambda_a *P_{ijk}^h = 0, \quad (2.1)$$

for some non-zero recurrence vector  $\lambda_a$  and is called an Einstein-Sasakian projective recurrent space, or briefly an  $E - P^* - space$ .

We, now, have the following theorem:

**Theorem (2.1):** A necessary and sufficient condition for an  $E - P$  space to be a Sasakian recurrent space is that the scalar curvature be equal to zero.

**Proof:** Suppose that an  $E - P^*$  space is Sasakian recurrent. Making use of equations (1.7), (1.8) and (1.10) in (2.1), we get

$$R_{ijk,a}^h = \lambda_a \left[ R_{ijk}^h + \frac{n}{n(n+2)} (g_{ik} \delta_j^h - g_{jk} \delta_i^h + F_{ik} F_j^h - F_{jk} F_i^h + 2F_{ij} F_k^h) \right]. \quad (2.2)$$

Since an  $E - P^*$  space is Sasakian recurrent, equation (2.2) reduces to

$$\frac{R}{n(n+4)} (g_{ik} \delta_j^h - g_{jk} \delta_i^h + F_{ik} F_j^h - F_{jk} F_i^h + 2F_{ij} F_k^h) = 0, \quad (2.3)$$

which gives  $R = 0$ , i.e., scalar curvature is zero.

Conversely, if an  $E - P^*$  space satisfies  $R = 0$ , then (2.2) reduces to

$$R_{ijk,a}^h - \lambda_a R_{ijk}^h = 0,$$

which shows that the space is Sasakian recurrent.

This completes the proof of the theorem.

Similarly, in view of Theorem (2.1) and equations (1.7), (1.8) and (1.11), we can prove the following theorem:

**Theorem (2.2):** A necessary and sufficient condition for an  $E - S^*$  space to be a Sasakian recurrent is that the scalar curvature be equal to zero.

**Theorem (2.3):** An  $E - P^*$  space is an  $E - S^*$  space, iff the scalar curvature  $R$  is equal to zero.

**Proof:** Suppose that an  $E - P^*$  space is an  $E - S^*$  space. Differentiating (1.11) covariantly with respect to  $x^n$  and using (1.7), we get

$$E_{ijk,a}^h = *P_{ijk,a}^h + \frac{R_a}{(n+2)(n+4)} (g_{ik} \delta_j^h - g_{jk} \delta_i^h + F_{ik} F_j^h - F_{jk} F_i^h + 2F_{ij} F_k^h). \quad (2.4)$$

Multiplying (1.11) by  $\lambda_a$  and subtracting the resulting equation from (2.4), we have

$$E_{ijk,a}^h - \lambda_a E_{ijk}^h = *P_{ijk,a}^h - \lambda_a *P_{ijk}^h + \frac{(R_a - \lambda_a R)}{(n+2)(n+4)} (g_{ik} \delta_j^h - g_{jk} \delta_i^h + F_{ik} F_j^h - F_{jk} F_i^h + 2F_{ij} F_k^h) \quad (2.5)$$

Now, making use of the above supposition and (1.7), equation (2.5) reduces to

$$\frac{\lambda_a R}{(n+2)(n+4)} (g_{ik} \delta_j^h - g_{jk} \delta_i^h + F_{ik} F_j^h - F_{jk} F_i^h + 2F_{ij} F_k^h) = 0, \text{ which implies } R = 0.$$

Conversely, let us suppose that in an  $E - P^*$  space, the scalar curvature  $R = 0$ .

Hence equation (2.5) reduces to

$$E_{ijk,a}^h - \lambda_a E_{ijk}^h = 0,$$

which shows that the space is an  $E - S^*$  space.

This completes the proof of the theorem.

**Theorem (2.4):** If a Sasakian space satisfies any two of the following properties:

- (i).the space is an  $E - S^*$  space,
- (ii).the space is an  $E - P^*$  space,
- (iii).the scalar curvature is equal to zero, then it must satisfy the third also.

**Proof:** An  $E - S^*$  space and an  $E - P^*$  space are characterized respectively by equations (1.15) and (2.1).

The statement of the above theorem follows in view of equations (1.15), (2.1) and (2.5).

We have the following Lemmas from Walkar [7] and Yano & Bochner [9].

**Lemma (2.1):** The curvature tensor  $R_{hijk}$  satisfies the identity:

$$R_{hijk,lm} - R_{hijk,ml} + R_{jklm,hi} - R_{lmhi,jk} + R_{lmhi,kj} = 0, \quad (2.6)$$

where  $R_{hijk,l,m} \stackrel{\text{def}}{=} R_{hijk,lm}$ .

**Lemma (2.2):** If  $\alpha_{\alpha\beta}, b_\gamma$  are quantities satisfying:

$$\alpha_{\alpha\beta} = \alpha_{\beta\alpha} \text{ and } \alpha_{\alpha\beta} b_\gamma + \alpha_{\beta\gamma} b_\alpha + \alpha_{\gamma\alpha} b_\beta = 0, \quad (2.7)$$

for  $\alpha, \beta, \gamma = 1, 2, \dots, N$ , then either all the  $\alpha_{\alpha\beta}$  are zero, or all the  $b_\gamma$  are zero.

Making use of the above Lemmas, we shall prove the following theorems:

**Theorem (2.5):** In an  $E - P^*$  space, either recurrence vector is gradient, or the space is of constant holomorphic sectional curvature.

**Proof:** Differentiating (2.2) covariantly with respect to  $x^b$  and using equations (1.4), (1.7), (1.8) and (1.10), we get

$$R_{ijk,ab}^h = (\lambda_{a,b} + \lambda_a \lambda_b) * P_{ijk}^h, \quad (2.8)$$

where  $R_{ijk,a,b}^h \stackrel{\text{def}}{=} R_{ijk,ab}^h$ .

Multiplying (2.8) by  $g_{hi}$ , we have

$$R_{hijk,ab} = (\lambda_{a,b} + \lambda_a \lambda_b) * P_{ijkl}. \quad (2.9)$$

From (2.9) and the identity (2.6), we get

$$\lambda_{ab} * P_{ijkh} + \lambda_{ij} * P_{khab} + \lambda_{kh} * P_{abij} = 0, \quad (2.10)$$

where  $\lambda_{ab} \stackrel{\text{def}}{=} \lambda_{b,a} - \lambda_{a,b}$ .

Equation (2.10) is of the form (2.7), since  $*P_{ijkh} = *P_{khij}$ . Thus, from Lemma (2.2), we prove the Theorem (2.5).

Now, from (1.8) and the Bianchi identity

$$R_{ijk,a}^h + R_{jka,i}^h + R_{kai,j}^h = 0 \quad (2.11)$$

we have

$$R_{ijk,l}^h = 0.$$

Thus, contracting (2.2) with respect to  $h$  and  $a$ , we get

$$\lambda_a R_{ijk}^l + \frac{R}{n(n+2)} (\lambda_j g_{ik} - \lambda_i g_{jk} + \lambda_l F_{ik} F_j^l - \lambda_l F_{jk} F_i^l + 2\lambda_l F_{ij} F_k^l) = 0, \quad (2.12)$$

Furthermore, substituting (2.2) in (2.11) and then transvecting with  $\lambda^a$ , we get

$$\begin{aligned} \lambda^l \lambda_l R_{ijk}^h + \lambda^l \lambda_l \frac{R}{n(n+2)} (g_{ik} \delta_j^h - g_{jk} \delta_i^h + F_{ik} F_j^h - F_{jk} F_i^h + 2F_{ij} F_k^h) \\ + \lambda_i \left\{ \lambda^l R_{jik}^h + \frac{R}{n(n+2)} (\lambda^h g_{jk} - \lambda_k \delta_j^h + \lambda^a F_{jk} - \lambda^a F_{kj} - \lambda^a F_{ak} F_j^h + 2\lambda^a F_{ja} F_k^h) \right\} \\ + \lambda_j \left\{ \lambda^l R_{ijk}^h + \frac{R}{n(n+2)} (\lambda_k \delta_i^h - \lambda^h g_{ik} + \lambda^a F_{ak} F_i^h - \lambda^a F_{ik} F_a^h + 2\lambda^a F_{ai} F_k^h) \right\} = 0. \end{aligned} \quad (2.13)$$

Which, in view of (2.12), gives

$$\lambda^l \lambda_l \left\{ R_{ijk}^h + \frac{R}{n(n+2)} (g_{ik} \delta_j^h - g_{jk} \delta_i^h + F_{ik} F_j^h - F_{jk} F_i^h + 2F_{ij} F_k^h) \right\} = 0. \quad (2.14)$$

Since  $\lambda_a$  is a non-zero vector, in view of Definition (1.3), we have the following:

**Theorem (2.6):** An  $E - P^*$  space is the space of constant holomorphic sectional curvature [8].

## REFERENCES

- [1] Lal K. B. and Singh S. S., On Kaehlerian spaces with recurrent Bochner curvature, *Acc. Naz. Deilincei, Rend.*, **51 (3-4)**, (1971), 213-220.
- [2] Mathai S., Kaehlerian recurrent spaces, *Ganita*, **20 (2)**, (1969), 121-133.
- [3] Sinha B. B., On H- curvature tensors in a Kaehlerian manifold, *Kyungpook Math, Jour.*, **13 (2)**, (1973), 185-189.
- [4] Singh U. P. and Singh A. K., On Kaehlerian Conharmonic recurrent and Kaehlerian Conharmonic symmetric spaces, *Acc. Naz. Deilincei, Rend.*, **62 (2)**, (1977), 173-179.
- [5] Singh S. S., On Kaehlerian space with recurrent holomorphic projective curvature tensor, *Acc. Naz. Deilincei, Rend.*, **55 (3-4)**, (1973), 214-218.
- [6] Singh A. K., On Einstein-Kaehlerian conharmonic recurrent spaces, *Indian J. Pure and Appl. Math.*, **10 (4)**, (1979), 486-492.
- [7] Walker A. G., On Ruse's space of recurrent curvature, *Proc. Lond, math. Soc.*, **52 (II series)**, (1950), 36-64.
- [8] Yano K., *Differential Geometry on Complex and Almost Complex Spaces*, Pergamon Press, London (1965).
- [9] Yano K. and Bochner S., Curvature and Betti numbers, *Ann. Math. Stud.*, **32**, (1953), 83-105.

**Source of support: Nil, Conflict of interest: None Declared**