

## SOME RESULTS BASED ON RELATIVE DEFECTS OF SPECIAL TYPE OF DIFFERENTIAL POLYNOMIALS

Sanjib Kumar Datta<sup>1\*</sup> & Sudipta Kumar Pal<sup>2</sup>

<sup>1</sup>Department of Mathematics, University of Kalyani, P. O. - Kalyani, Dist.-Nadia, PIN-741235,  
West Bengal, India

<sup>2</sup>Uttar Dalkhola High School, P.O.-Dalkhola, Dist. - Uttar Dinajpur, PIN-733201, West Bengal, India

(Received on: 16-06-12; Revised & Accepted on: 10-07-12)

### ABSTRACT

The aim of this paper is to compare the relative Valiron defect with the relative Nevanlinna defect of special type of differential polynomials generated by transcendental meromorphic functions.

**Mathematics Subject Classification (2010):** 30D35, 30D30.

**Keywords and phrases:** Meromorphic function, relative Nevanlinna defect, relative Valiron defect, special type of differential polynomial.

### 1. INTRODUCTION, DEFINITIONS AND NOTATIONS

Let  $f$  be a transcendental meromorphic function defined in the open complex plane  $\mathbb{C}$ . A monomial in  $f$  is an expression of the form  $M[f] = (f)^{n_0} (f^{(1)})^{n_1} \dots (f^{(k)})^{n_k}$  where  $n_0, n_1, n_2, \dots, n_k$  are non negative integers.  $\gamma_M = n_0 + n_1 + \dots + n_k$  and  $\Gamma_M = n_0 + 2n_1 + \dots + (k+1)n_k$  are respectively called the degree and weight of the monomial.

If  $M_1[f], M_2[f], \dots, M_n[f]$  denote monomials in  $f$ , then  $Q[f] = a_1 M_1[f] + a_2 M_2[f] + \dots + a_n M_n[f]$ , where  $a_i \neq 0 (i=1, 2, \dots, n)$  is called a differential polynomial generated by  $f$  of degree  $\gamma_Q = \max \{\gamma_{M_j} : 1 \leq j \leq n\}$  and weight  $\Gamma_Q = \max \{\Gamma_{M_j} : 1 \leq j \leq n\}$ .

Also we call numbers  $\underline{\gamma}_Q = \min_{1 \leq j \leq n} \gamma_{M_j}$  and  $k$  (the order of the highest derivative of  $f$ ) the lower degree and the order of  $Q[f]$  respectively. If  $\underline{\gamma}_Q = \gamma_Q$ ,  $Q[f]$  is called a homogeneous differential polynomial.

For  $a \in \mathbb{C} \cup \{\infty\}$ , the quantity

$$\delta(a; f) = 1 - \limsup_{r \rightarrow \infty} \frac{N(r, a; f)}{T(r, f)} = \liminf_{r \rightarrow \infty} \frac{m(r, a; f)}{T(r, f)}$$

is called the Nevanlinna's deficiency of the value 'a'. Similarly the Valiron defect of 'a' is defined as

$$\Delta(a; f) = 1 - \liminf_{r \rightarrow \infty} \frac{N(r, a; f)}{T(r, f)} = \limsup_{r \rightarrow \infty} \frac{m(r, a; f)}{T(r, f)}.$$

The term  $\delta_R^{(k)}(a; f) = 1 - \limsup_{r \rightarrow \infty} \frac{N(r, a; f^{(k)})}{T(r, f)}$  for  $k = 1, 2, 3, \dots$  is called the relative Nevanlinna's defect of 'a' with respect to  $f^{(k)}$ . In a like manner  $\Delta_R^{(k)}(a; f) = 1 - \liminf_{r \rightarrow \infty} \frac{N(r, a; f^{(k)})}{T(r, f)}$  for  $k = 1, 2, 3, \dots$  is called the relative Valiron defect of 'a' with respect to  $f^{(k)}$ . Xiong [3] has shown various relations between the usual defects and relative defects of meromorphic functions. Following Datta and Mondal [1], in the paper we consider  $F = f^n Q[f]$ ,  $Q[f]$  being a differential polynomial in  $f$  and  $n = 1, 2, 3, \dots$  and compare the relative Valiron defect with the relative Nevanlinna defect of  $F$ .

**Corresponding author: Sanjib Kumar Datta<sup>1\*</sup>**

<sup>1</sup>Department of Mathematics, University of Kalyani, P.O.-Kalyani, Dist.-Nadia, PIN-741235, West Bengal, India

The term  $S(r, f)$  denotes any quantity satisfying  $S(r, f) = o\{T(r, f)\}$  as  $r \rightarrow \infty$  through all values of  $r$  if  $f$  is of finite order and except possibly for a set of  $r$  of finite linear measure otherwise. We do not explain the standard definitions and notations of the value distribution and the Nevanlinna theory as those are available in [2].

## 2. LEMMAS

In this section we present some lemmas which will be needed in the sequel.

**Lemma 1:** Let  $k$  be any positive integer and  $\varphi = \sum_{i=0}^n a_i f^{(i)}$ , where  $a_i$  are meromorphic functions such that  $T(r, a_i) = S(r, f)$  for  $i = 0, 1, 2, \dots, k$ .

**Lemma 2:** Let  $F = f^n Q[f]$  where  $Q[f]$  is a differential polynomial in  $f$ . If  $n \geq 1$  then  $\lim_{r \rightarrow \infty} \frac{T(r, F)}{T(r, f)} = 1$ .

The proof is omitted.

**Lemma 3:** Let  $F = f^n Q[f]$  where  $Q[f]$  is a differential polynomial in  $f$ . If  $n \geq 1$  then for any

$$\alpha, \delta_R^F(\alpha; f) = \liminf_{r \rightarrow \infty} \frac{m(r, \alpha; F)}{T(r, f)} \text{ and } \Delta_R^F(\alpha; f) = \limsup_{r \rightarrow \infty} \frac{m(r, \alpha; F)}{T(r, f)}.$$

**Proof:** In view of Lemma 2 we get that

$$\begin{aligned} \delta_R^F(\alpha; f) &= 1 - \limsup_{r \rightarrow \infty} \frac{N(r, \alpha; F)}{T(r, f)} \\ &= 1 - \limsup_{r \rightarrow \infty} \frac{N(r, \alpha; F)}{T(r, F)} \cdot \lim_{r \rightarrow \infty} \frac{T(r, F)}{T(r, f)} \\ &= 1 - \limsup_{r \rightarrow \infty} \frac{N(r, \alpha; F)}{T(r, F)} \cdot 1 \\ &= 1 - \limsup_{r \rightarrow \infty} \frac{N(r, \alpha; F)}{T(r, F)} \\ &= \liminf_{r \rightarrow \infty} \frac{m(r, \alpha; F)}{T(r, F)} \\ &= \liminf_{r \rightarrow \infty} \frac{m(r, \alpha; F)}{T(r, f)} \cdot \lim_{r \rightarrow \infty} \frac{T(r, f)}{T(r, F)} \\ &= \liminf_{r \rightarrow \infty} \frac{m(r, \alpha; F)}{T(r, f)} \cdot 1 \\ &= \liminf_{r \rightarrow \infty} \frac{m(r, \alpha; F)}{T(r, f)}. \end{aligned}$$

This proves the first part of the lemma.

Similarly the second part of Lemma 3 follows.

## 3. THEOREMS.

In this section we present the main results of the paper.

**Theorem 1:** Let  $f$  be a transcendental meromorphic function of finite order  $\rho_f$  and satisfying the condition  $m(r, f) = S(r, f)$ . If  $a, b$  and  $c$  are three non zero finite complex numbers then

$$3\delta(a; f) + 2\delta(b; f) + \delta(c; f) + 5\Delta_R^F(\infty; F) \leq 5\Delta(\infty; f) + 5\Delta_R^F(\infty; F)$$

where  $F$  is a differential polynomial in  $f$  of the form  $F = f^n Q[f]$  with  $n \geq 1$ .

**Proof:** Let us consider the following identity

$$\frac{b-a}{f-a} = \left[ \frac{F}{f-a} \left\{ \frac{f-a}{F} - \frac{f-b}{F} \right\} - \frac{f-c}{F} \cdot \frac{F}{f-a} \cdot \frac{F}{f-a} \left\{ \frac{f-a}{F} - \frac{f-b}{F} \right\} \right] \cdot \frac{f}{c}$$

Since  $m\left(r, \frac{1}{f-a}\right) \leq m\left(r, \frac{b-a}{f-a}\right) + O(1)$  and  $m\left(r, \frac{f}{c}\right) \leq m(r, f) + O(1)$ , we get from the above identity in view of Lemma 1 that

$$m\left(r, \frac{b-a}{f-a}\right) \leq m\left(r, \frac{f-a}{F}\right) + m\left(r, \frac{f-b}{F}\right) + m\left(r, \frac{f-a}{F}\right) + m\left(r, \frac{f-b}{F}\right) + m\left(r, \frac{f-c}{F}\right) + m\left(r, \frac{f}{c}\right) + S(r, f)$$

$$i.e., m\left(r, \frac{1}{f-a}\right) \leq 2m\left(r, \frac{f-a}{F}\right) + 2m\left(r, \frac{f-b}{F}\right) + m\left(r, \frac{f-c}{F}\right) + m(r, f) + S(r, f) + O(1)$$

$$i.e., m\left(r, \frac{1}{f-a}\right) \leq 2T\left(r, \frac{f-a}{F}\right) - 2N\left(r, \frac{f-a}{F}\right) + 2T\left(r, \frac{f-b}{F}\right) - 2N\left(r, \frac{f-b}{F}\right) + T\left(r, \frac{f-c}{F}\right) - N\left(r, \frac{f-c}{F}\right) + m(r, f) + S(r, f) + O(1). \quad (1)$$

Now by the relation  $T\left(r, \frac{1}{f}\right) = T(r, f) + O(1)$  and in view of Lemma 1 it follows from (1) that

$$m\left(r, \frac{1}{f-a}\right) \leq 2T\left(r, \frac{F}{f-a}\right) - 2N\left(r, \frac{f-a}{F}\right) + 2T\left(r, \frac{F}{f-b}\right) - 2N\left(r, \frac{f-b}{F}\right) + T\left(r, \frac{F}{f-c}\right) - N\left(r, \frac{f-c}{F}\right) + m(r, f) + S(r, f) + O(1)$$

$$i.e., m\left(r, \frac{1}{f-a}\right) \leq 2\left\{N\left(r, \frac{F}{f-a}\right) - N\left(r, \frac{f-a}{F}\right)\right\} + 2\left\{N\left(r, \frac{F}{f-b}\right) - N\left(r, \frac{f-b}{F}\right)\right\} + N\left(r, \frac{F}{f-c}\right) - N\left(r, \frac{f-c}{F}\right) + m(r, f) + S(r, f) + O(1). \quad (2)$$

In view of [p.34, [2]] it follows from (2) that

$$m\left(r, \frac{1}{f-a}\right) \leq 2\left\{N(r, F) + N\left(r, \frac{1}{f-a}\right) - N(r, f-a) - N\left(r, \frac{1}{f}\right)\right\} + 2\left\{N(r, F) + N\left(r, \frac{1}{f-b}\right) - N(r, f-b) - N\left(r, \frac{1}{f}\right)\right\} + \left\{N(r, F) + N\left(r, \frac{1}{f-c}\right) - N(r, f-c) - N\left(r, \frac{1}{f}\right)\right\} + m(r, f) + S(r, f) + O(1). \quad (3)$$

Now applying the condition  $m(r, f) = S(r, f)$  it follows from (3) that

$$m\left(r, \frac{1}{f-a}\right) \leq 5N(r, F) - 5N(r, f) - 5N\left(r, \frac{1}{f}\right) + 2N\left(r, \frac{1}{f-a}\right) + 2N\left(r, \frac{1}{f-b}\right) + N\left(r, \frac{1}{f-c}\right) + S(r, f)$$

$$i.e., \liminf_{r \rightarrow \infty} \frac{m\left(r, \frac{1}{f-a}\right)}{T(r, f)} \leq 5 \liminf_{r \rightarrow \infty} \left\{ \frac{N(r, F)}{T(r, f)} - \frac{N(r, f)}{T(r, f)} - \frac{N\left(r, \frac{1}{f}\right)}{T(r, f)} \right\} + 2 \limsup_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{f-a}\right)}{T(r, f)} + 2 \limsup_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{f-b}\right)}{T(r, f)} + \limsup_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{f-c}\right)}{T(r, f)}$$

$$i.e., \liminf_{r \rightarrow \infty} \frac{m\left(r, \frac{1}{f-a}\right)}{T(r, f)} \leq 5 \liminf_{r \rightarrow \infty} \frac{N(r, F)}{T(r, f)} - 5 \liminf_{r \rightarrow \infty} \frac{N(r, f)}{T(r, f)} - 5 \liminf_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{f}\right)}{T(r, f)} + 2 \limsup_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{f-a}\right)}{T(r, f)} + 2 \limsup_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{f-b}\right)}{T(r, f)} + \limsup_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{f-c}\right)}{T(r, f)}$$

$$i.e., \delta(a; f) \leq 5\{1 - \Delta_R^F(\infty; f)\} - 5\{1 - \Delta(\infty; f)\} - 5\{1 - \Delta_R^F(0; f)\} + 2\{1 - \delta(a; f)\} + 2\{1 - \delta(b; f)\} + \{1 - \delta(c; f)\}$$

$$i.e., 3\delta(a; f) + 2\delta(b; f) + \delta(c; f) + 5\Delta_R^F(\infty; f) \leq 5\Delta(\infty; f) + 5\Delta_R^F(0; f).$$

This proves the theorem.

**Theorem 2:** Let  $f$  be a meromorphic function of finite order satisfying the condition  $m(r, f) = S(r, f)$  and also let If  $F = f^n Q[f]$  is a differential polynomial in  $f$  with  $n \geq 1$ . If  $a, b, c$  and  $d$  are any four distinct complex numbers then

$$\delta(d; f) + \delta_R^F(b; f) + \delta_R^F(c; f) \leq 2.$$

**Proof:** Let us consider the following identity

$$\frac{1}{f-d} = \left[ \frac{1}{a} \left\{ \frac{F}{f-a} - \frac{F-a}{f^n} \cdot \frac{f^n}{f-a} \right\} \cdot \left\{ \frac{F}{f-d} \cdot \frac{1}{F} \right\} \right] \cdot (f-a)$$

Since  $m(r, f-a) \leq m(r, f) + O(1)$ , we get from the above identity in view of Lemma 1 that

$$m\left(r, \frac{1}{f-d}\right) \leq m\left(r, \frac{1}{f}\right) + m(r, f) + S(r, f) + O(1)$$

$$i.e., m\left(r, \frac{1}{f-d}\right) \leq T\left(r, \frac{1}{F}\right) - N\left(r, \frac{1}{F}\right) + m(r, f) + S(r, f) + O(1)$$

Now by the relation  $T\left(r, \frac{1}{F}\right) = T(r, f) + O(1)$  we get from the above

$$m\left(r, \frac{1}{f-d}\right) \leq T(r, F) - N\left(r, \frac{1}{F}\right) + m(r, f) + S(r, f) + O(1). \quad (4)$$

Now by Nevanlinna's second fundamental theorem it follows from (4) that

$$m\left(r, \frac{1}{f-d}\right) \leq \bar{N}\left(r, \frac{1}{F}\right) + \bar{N}\left(r, \frac{1}{F-b}\right) + \bar{N}\left(r, \frac{1}{F-c}\right) - N\left(r, \frac{1}{F}\right) + m(r, f) + S(r, f) + O(1). \quad (5)$$

As  $\bar{N}\left(r, \frac{1}{F}\right) - N\left(r, \frac{1}{F}\right) \leq 0$  and applying the condition  $m(r, f) = S(r, f)$  it follows from (5) that

$$m\left(r, \frac{1}{f-d}\right) \leq \bar{N}\left(r, \frac{1}{F-b}\right) + \bar{N}\left(r, \frac{1}{F-c}\right) + S(r, f)$$

$$i.e., m\left(r, \frac{1}{f-d}\right) \leq N\left(r, \frac{1}{F-b}\right) + N\left(r, \frac{1}{F-c}\right) + S(r, f)$$

$$i.e., m\left(r, \frac{1}{f-d}\right) \leq T\left(r, \frac{1}{F-b}\right) - m\left(r, \frac{1}{F-b}\right) + T\left(r, \frac{1}{F-c}\right) - m\left(r, \frac{1}{F-c}\right) + S(r, f)$$

$$i.e., m\left(r, \frac{1}{f-d}\right) \leq T(r, F) - m\left(r, \frac{1}{F-b}\right) - m\left(r, \frac{1}{F-c}\right) + S(r, f)$$

$$i.e., \liminf_{r \rightarrow \infty} \frac{m\left(r, \frac{1}{f-d}\right)}{T(r, f)} \leq 2 \liminf_{r \rightarrow \infty} \frac{T(r, F)}{T(r, f)} - \liminf_{r \rightarrow \infty} \frac{m\left(r, \frac{1}{F-b}\right)}{T(r, f)} - \liminf_{r \rightarrow \infty} \frac{m\left(r, \frac{1}{F-c}\right)}{T(r, f)}$$

$$i.e., \delta(d; f) \leq 2.1 - \delta_R^F(b; f) - \delta_R^F(c; f)$$

$$i.e., \delta(d; f) + \delta_R^F(b; f) + \delta_R^F(c; f) \leq 2.$$

This proves the theorem.

**Theorem 3:** Let  $f$  be a transcendental meromorphic function of finite order  $\rho_f$  and satisfying the condition  $m(r, f) = S(r, f)$ . If  $a$  and  $c$  are any two distinct complex numbers and let  $F = f^n Q[f]$  is a differential polynomial in  $f$  with  $n \geq 1$  then

$$\delta(0; f) + \delta(c; f) + \Delta_R^F(\infty; f) \leq \Delta(\infty; f) + 2\Delta_R^F(0; f).$$

**Proof:** Consider the following identity

$$\frac{c}{f} = \left[ \left\{ 1 - \frac{f-c}{F} \cdot \frac{F}{f} \right\} \left\{ \frac{F}{f-a} \cdot \frac{1}{F} \right\} \right] \cdot (f-a)$$

Since  $m\left(r, \frac{1}{f}\right) \leq m\left(r, \frac{c}{f}\right) + O(1)$  and  $m(r, f-a) \leq m(r, f) + O(1)$ , we get from the above identity in view of Lemma 1 that

$$m\left(r, \frac{c}{f}\right) \leq m\left(r, \frac{f-c}{F}\right) + m\left(r, \frac{1}{F}\right) + m(r, f) + S(r, f) + O(1)$$

$$i.e., m\left(r, \frac{1}{f}\right) \leq T\left(r, \frac{f-c}{F}\right) - N\left(r, \frac{f-c}{F}\right) + T\left(r, \frac{1}{F}\right) - N\left(r, \frac{1}{F}\right) + m(r, f) + S(r, f) + O(1). \quad (6)$$

Now by Nevanlinna's first fundamental theorem and in view of Lemma 1 it follows from (6) that

$$m\left(r, \frac{1}{f}\right) \leq T\left(r, \frac{F}{f-c}\right) - N\left(r, \frac{f-c}{F}\right) + T(r, F) - N\left(r, \frac{1}{F}\right) + m(r, f) + S(r, f) + O(1).$$

$$i.e., m\left(r, \frac{1}{f}\right) \leq N\left(r, \frac{F}{f-c}\right) - N\left(r, \frac{f-c}{F}\right) - N\left(r, \frac{1}{F}\right) + T(r, F) + m(r, f) + S(r, f) + O(1). \quad (7)$$

In view of {p.34, [2]} it follows from (7) that

$$m\left(r, \frac{1}{f}\right) \leq N(r, F) + N\left(r, \frac{1}{f-c}\right) - N(r, f-c) - N\left(r, \frac{1}{F}\right) - N\left(r, \frac{1}{F}\right) + T(r, F) + m(r, f) + S(r, f) + O(1)$$

$$i.e., \liminf_{r \rightarrow \infty} \frac{m(r, \frac{1}{f})}{T(r, f)} \leq \liminf_{r \rightarrow \infty} \frac{N(r, F)}{T(r, f)} - \liminf_{r \rightarrow \infty} \frac{N(r, f)}{T(r, f)} - 2 \liminf_{r \rightarrow \infty} \frac{N(r, \frac{1}{F})}{T(r, f)} + \limsup_{r \rightarrow \infty} \frac{N(r, \frac{1}{f-c})}{T(r, f)} + \limsup_{r \rightarrow \infty} \frac{T(r, F)}{T(r, f)}$$

$$i.e., \delta(0; f) \leq \{1 - \Delta_R^F(\infty; f)\} - \{1 - \Delta(\infty; f)\} - 2\{1 - \Delta_R^F(0; f)\} + \{1 - \delta(c; f)\} + 1$$

$$i.e., \delta(0; f) + \delta(c; f) + \Delta_R^F(\infty; f) \leq \Delta(\infty; f) + 2\Delta_R^F(0; f).$$

Thus the theorem is established.

## REFERENCES

- [1] S.K. Datta and S. Mondal: Relative defects of a special type of differential polynomial, J. Mech. Cont. & Math. Soc., Vol. 5, No. 2, January (2011), pp. 691-706.
- [2] W.K. Hayman: Meromorphic Functions, The Clarendon Press, Oxford (1964).
- [3] Q.L. Xiong : A fundamental inequality in the theory of meromorphic functions and its applications, Chinese Mathematics, Vol. 9, No. 1 (1967), pp. 146-167.

**Source of support: Nil, Conflict of interest: None Declared**