

SOME COMMON FIXED POINT THEOREMS IN FUZZY METRIC SPACE

T. Veerapandi

Associate Professor of Mathematics, P. M. T. College, Melaneelithanallur-627953, India

V. Raji Muniyammal*

Department of Mathematics, Mano College, Nagampatti, India

J. Paulraj Joseph

Associate Professor of Mathematics, Manonmaniam Sundaranar University, Tirunelveli, India

(Received on: 07-08-12; Revised & Accepted on: 08-09-12)

ABSTRACT

This paper presents some common fixed point theorems for occasionally weakly compatible mappings in fuzzy metric space under various conditions.

Mathematics Subject Classification: 47H10, 54H25.

Keywords: Occasionally weakly compatible mappings, fuzzy metric space.

INTRODUCTION

Fuzzy set was defined by Zadeh [24]. Kramosil and Michalek [12] introduced fuzzy metric space, George and Veeramani [4] modified the notion of fuzzy metric spaces with the help of continuous t-norms. Many researchers have obtained common fixed point theorems for mappings satisfying different types of commutativity conditions.

Vasuki [23] proved fixed point theorems for R-weakly commuting mappings. Pant [16, 17, 18] introduced the new concept reciprocally continuous mappings and established some common fixed point theorems.

Balasubramaniam et al. [2] have shown that Rhoades [20] open problem on the existence of contractive definition which generates a fixed point but does not force the mappings to be continuous at the fixed point, possesses an affirmative answer.

Pant and Jha [18] obtained some analogous results proved by Balasubramaniam et al.

Recent literature on fixed point in fuzzy metric space can be viewed in [7, 14, 22]. This paper presents some common fixed point theorems for more general commutative condition i.e., occasionally weakly compatible mappings in fuzzy metric space. Before giving the results, some preliminary definitions are given below.

1. PRELIMINARIES

Definition 1.1 [21]: A fuzzy set A in X is a function with domain X and values in $[0, 1]$

Definition 1.2 [18]: A binary operation $*$: $[0,1] \times [0,1] \rightarrow [0,1]$ is a continuous t-norm if $*$ is satisfying conditions:

- (1) $*$ is an commutative and associative
- (2) $*$ is continuous
- (3) $a * 1 = a$ for all $a \in [0,1]$
- (4) $a * b \leq c * d$ whenever $a \leq c$ and $b \leq d$ and $a, b, c, d \in [0,1]$

Example 1.3 [4]: Example for continuous t-norms are

- (1) $a * b = ab$ for all $a, b \in [0,1]$
- (2) $a * b = \min\{a, b\}$ for all $a, b \in [0,1]$

Corresponding author: V. Raji Muniyammal*
Department of Mathematics, Mano College, Nagampatti, India

Definition 1.4[4]: A 3-tuple $(X, M, *)$ is said to be a fuzzy metric space if X is an arbitrary set, $*$ is a continuous t-norm and M is a fuzzy set on $X^2 \times [0, \infty]$ satisfying the following conditions for all $x, y, z \in X$ such that t is in $[0, \infty]$

- (FM-1) $M(x, y, t) > 0$;
- (FM-2) $M(x, y, t) = 1$ if and only if $x = y$
- (FM-3) $M(x, y, t) = M(y, x, t)$;
- (FM-4) $M(x, y, t) * M(y, z, s) \leq M(x, z, t + s)$;
- (FM-5) $M(x, y, *) : (0, \infty) \rightarrow (0, 1]$ is continuous

Then M is called a fuzzy metric on X .

Then $M(x, y, t)$ denotes the degree of nearness between x and y with respect to t

Example 1.5 [4]: Let (X, d) be a metric space. Denote $a * b = ab$ for all $a, b \in [0, 1]$ and let M_d be fuzzy sets on $X^2 \times (0, \infty)$ defined as follows:

$$M_d(x, y, t) = \frac{t}{t + d(x, y)}$$

Then $(X, M_d, *)$ is a fuzzy metric space.

We call this fuzzy metric induced by a metric d the standard intuitionistic fuzzy metric

Definition 1.6 [4]: Let $(X, M, *)$ be a fuzzy metric space. Then

- (1) a sequence $\{x_n\}$ in X is said to converge to x in X if for each $\varepsilon > 0$ and each $t > 0$, there exists $n_0 \in \mathbb{N}$ such that $M(x_n, x, t) > 1 - \varepsilon$ for all $n \geq n_0$
- (2) a sequence $\{x_n\}$ in X is said to be Cauchy if for each $\varepsilon > 0$ and each $t > 0$ there exists $n_0 \in \mathbb{N}$ such that $M(x_n, x_m, t) > 1 - \varepsilon$ for all $n, m \geq n_0$
- (3) A fuzzy metric space in which every Cauchy sequence is convergent is said to be complete.

Definition 1.7[20]: A pair of self mappings (f, g) of a fuzzy metric space $(X, M, *)$ is said to be

- (1) weakly commuting if $M(fgx, gfx, t) \geq M(fx, gx, t)$ for all $x \in X$ and $t > 0$
- (2) R-weakly commuting if there exists some $R > 0$ such that $M(fgx, gfx, t) \geq M(fx, gx, t/R)$ for all $x \in X$ and $t > 0$

Definition 1.8[11]: Two self-mappings f and g of a fuzzy metric space $(X, M, *)$ are called compatible if $\lim_{n \rightarrow \infty} M(fgx_n, gfx_n, t) = 1$ whenever $\{x_n\}$ is a sequence in X such that $\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = x$ for some $x \in X$

Definition 1.9 [5]: Two self-maps f and g of a fuzzy metric $(X, M, *)$ are called reciprocally continuous on X if $\lim_{n \rightarrow \infty} fgx_n = fx$ and $\lim_{n \rightarrow \infty} gfx_n = gx$ whenever $\{x_n\}$ is a sequence in X such that $\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = x$ for some $x \in X$

Definition 1.10: Let X be a set, f, g self-maps of X . A point $x \in X$ is called a coincidence point f and g if and only if $fx = gx$.

We shall call $w = fx = gx$ a point of coincidence of f and g .

Definition 1.11[11]: A pair maps S and T is called weakly compatible pair if they commute at coincidence points

Definition 1.12: Two self-maps f and g of a set X are occasionally weakly compatible if and only if there is point $x \in X$ which is a coincidence point of f and g at which f and g are commute

Remark 1.13: A. Al-Thagafi and Naseer Shahzad [1] have shown that occasionally weakly is weakly compatible but converse is not true

Example 1.14 [1]: Let R be usual metric space.

Define $S, T: R \rightarrow R$ by $Sx = 3x$ and $Tx = x^2$ for all $x \in R$.

Then $Sx = Tx$ for $x = 0, 3$ but $ST0 = TS0$ and $3 \neq TS$. S and T are occasionally weakly compatible self-maps but not weakly compatible.

Lemma 1.15 [10]: Let X be a set, f, g be occasionally weakly compatible self-maps of X .

If f and g have unique point of coincidence, $w = fx = gx$, then w is the unique common fixed point of f and g .

2. MAIN RESULT

Theorem 2.1: Let $(X, M, *)$ be complete fuzzy metric space and let A, B and $\{S_i\}_{i=1}^n$ and $\{T_j\}_{j=1}^m$ be self-mappings of X .

Let pairs $\{A, S_i\}_{i=1}^n$ and $\{B, T_j\}_{j=1}^m$ be occasionally weakly compatible at x and y respectively.

If there exists $q \in (0, 1)$ such that

$$M(Ax, By, qt) \geq \min \left\{ \begin{array}{l} \{M(S_1x, T_jy, t)\}_{j=1}^m, \{M(S_2x, T_jy, t)\}_{j=1}^m, \dots, \{M(S_nx, T_jy, t)\}_{j=1}^m, \{M(S_ix, Ax, t)\}_{i=1}^n, \\ \{M(By, T_jy, t)\}_{j=1}^m, \frac{\sum_{j=1}^m \alpha_j M(Ax, T_jy, t) + \sum_{i=1}^n \beta_i M(By, S_ix, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \end{array} \right\} \quad (1)$$

for all $x, y \in X$, for all $t > 0$ and α_j 's ≥ 0 and β_i 's ≥ 0 with atleast one of α_j 's or one of β_i 's is non-zero, then there exists a unique point $w \in X$ such that $Aw = S_iw = w$ for all i and a unique point $z \in X$ such that $Bz = T_jz = z$ for all j .

More over $z = w$ so that there is a unique common fixed point of $A, B, \{S_i\}_{i=1}^n$ and $\{T_j\}_{j=1}^m$

Proof 2.1: Let the pairs $\{A, S_i\}_{i=1}^n$ and $\{B, T_j\}_{j=1}^m$ be occasionally weakly compatible at the points $x, y \in X$ respectively.

Then $Ax = S_ix$ and $By = T_jy$ for all i and j

Claim: $Ax = By$

If not,

$$\begin{aligned} M(Ax, By, t) &\geq \min \left\{ \begin{array}{l} \{M(S_1x, T_jy, t)\}_{j=1}^m, \{M(S_2x, T_jy, t)\}_{j=1}^m, \dots, \{M(S_nx, T_jy, t)\}_{j=1}^m, \{M(S_ix, Ax, t)\}_{i=1}^n, \\ \{M(By, T_jy, t)\}_{j=1}^m, \frac{\sum_{j=1}^m \alpha_j M(Ax, T_jy, t) + \sum_{i=1}^n \beta_i M(By, S_ix, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \end{array} \right\} \\ &= \min \left\{ M(Ax, By, t), M(Ax, Ax, t), M(By, By, t), \frac{\sum_{j=1}^m \alpha_j M(Ax, T_jy, t) + \sum_{i=1}^n \beta_i M(By, S_ix, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \right\} \\ &= \min \left\{ M(Ax, By, t), 1, 1, M(Ax, By, t) \left(\frac{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \right) \right\} \\ &= \min \{M(Ax, By, t), 1, 1, M(Ax, By, t)\} \\ &= M(Ax, By, t) \end{aligned}$$

a contradiction.

Therefore $Ax = By$

i.e., $Ax = S_ix = By = T_jy$

Suppose that there is a another point z such that $Az = S_iz$ then by (1) we have $Az = S_iz = By = T_jy$ so $Az = Ax$ and $w = Ax = S_ix$ is the unique point of coincidence of A and S_i

By lemma 1.15, w is the only common fixed point of A and S_i

Similarly there is a unique point $z \in X$ such that $z = Bz = T_jz$

Uniqueness: Assume that $w \neq z$

We have

$$M(w, z, qt) = M(Aw, Bz, t)$$

$$\begin{aligned} &\geq \min \left\{ \begin{array}{l} \{M(S_1 w, T_j z, t)\}_{j=1}^m, \{M(S_2 w, T_j z, t)\}_{j=1}^m, \dots, \{M(S_n w, T_j z, t)\}_{j=1}^m \\ \{M(S_i w, Aw, t)\}_{i=1}^n, \{M(Bz, T_j z, t)\}_{j=1}^m \\ \frac{\sum_{j=1}^m \alpha_j M(Aw, T_j z, t) + \sum_{i=1}^n \beta_i M(Bz, S_i w, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \end{array} \right\} \\ &= \min \left\{ M(w, z, t), M(w, w, t), M(z, z, t), \frac{\sum_{j=1}^m \alpha_j M(w, z, t) + \sum_{i=1}^n \beta_i M(z, w, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \right\} \\ &= \min \left\{ M(w, z, t), 1, 1, M(w, z, t) \left(\frac{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \right) \right\} \\ &= \min \{M(w, z, t), 1, 1, M(w, z, t)\} \\ &= M(w, z, t) \end{aligned}$$

a contradiction.

Therefore we have $z = w$ by lemma 1.15, z is a common fixed point of $A, B, \{S_i\}_{i=1}^n$ and $\{T_j\}_{j=1}^m$. The uniqueness of the fixed point holds from (1).

Corollary 2.2 [25]: Let $(X, M, *)$ be a complete fuzzy metric space and let A, B, S and T be self-mappings of X . Let the pairs $\{A, S\}$ and $\{B, T\}$ be occasionally weakly compatible.

If there exists $q \in (0, 1)$ such that

$$M(Ax, By, t) \geq \min \left\{ M(Sx, Ty, t), M(Sx, Ax, t), M(By, Ty, t), \left(\frac{M(Ax, Ty, t) + M(By, Sx, t)}{2} \right) \right\} \quad (2)$$

for all $x, y \in X$ and for all $t > 0$ then there exists a unique point $w \in X$ such that $Aw = Sw = w$ and a unique point $z \in X$ such that $Bz = Tz = z$.

Moreover $z = w$, so that there is a unique common fixed point of A, B, S and T .

Proof 2.2: Take $n=m=1$ and $\alpha_1 = \beta_1 = 1$ in the above theorem 2.1, the corollary follows.

Theorem 2.3: Let $(X, M, *)$ be complete fuzzy metric space and let A, B and $\{S_i\}_{i=1}^n$ and $\{T_j\}_{j=1}^m$ be self-mappings of X .

Let pairs $\{A, S_i\}_{i=1}^n$ and $\{B, T_j\}_{j=1}^m$ be occasionally weakly compatible at x and y respectively.

If there exists $q \in (0, 1)$ such that

$$M(Ax, By, qt) \geq \Psi \left(\min \left\{ \begin{array}{l} \{M(S_1 x, T_j y, t)\}_{j=1}^m, \{M(S_2 x, T_j y, t)\}_{j=1}^m, \dots, \{M(S_n x, T_j y, t)\}_{j=1}^m, \{M(S_i x, Ax, t)\}_{i=1}^n, \\ \{M(B_y, T_j y, t)\}_{j=1}^m, \frac{\sum_{j=1}^m \alpha_j M(Ax, T_j y, t) + \sum_{i=1}^n \beta_i M(By, S_i x, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \end{array} \right\} \right) \quad (3)$$

for all $x, y \in X$, for all $t > 0$ and α_j 's ≥ 0 and β_i 's ≥ 0 with at least one of α_j 's or one of β_i 's is non-zero, and $\Psi: [0, 1] \rightarrow [0, 1]$ such that $\Psi(t) > t$ for all $0 < t < 1$ then there exists a unique common fixed point of $A, B, \{S_i\}_{i=1}^n$ and $\{T_j\}_{j=1}^m$

Proof 2.3: Let the pairs $\{A, S_i\}_{i=1}^n$ and $\{B, T_j\}_{j=1}^m$ be occasionally weakly compatible at the points $x, y \in X$ respectively.

Then $Ax = S_i x$ and $By = T_j y$ for all i and j

Claim: $Ax = By$

If not,

$$\begin{aligned}
 M(Ax, By, t) &\geq \Psi \left(\min \left\{ \begin{aligned} &\{M(S_1x, T_jy, t)\}_{j=1}^m, \{M(S_2x, T_jy, t)\}_{j=1}^m, \dots, \{M(S_nx, T_jy, t)\}_{j=1}^m, \{M(S_ix, Ax, t)\}_{i=1}^n, \\ &\{M(By, T_jy, t)\}_{j=1}^m, \frac{\sum_{j=1}^m \alpha_j M(Ax, T_jy, t) + \sum_{i=1}^n \beta_i M(By, S_ix, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \end{aligned} \right\} \right) \\
 &= \Psi \left(\min \left\{ M(Ax, By, t), M(Ax, Ax, t), M(By, By, t), \frac{\sum_{j=1}^m \alpha_j M(Ax, By, t) + \sum_{i=1}^n \beta_i M(By, Ax, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \right\} \right) \\
 &= \Psi \left(\min \left\{ M(Ax, By, t), 1, 1, M(Ax, By, t) \left(\frac{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \right) \right\} \right) \\
 &= \Psi(\min\{M(Ax, By, t), 1, 1, M(Ax, By, t)\}) \\
 &= \Psi(M(Ax, By, t)) > M(Ax, By, t)
 \end{aligned}$$

a contradiction.

Therefore $Ax = By$

i.e., $Ax = S_ix = By = T_jy$

Suppose that there is a another point z such that $Az = S_iz$ then by (3) we have

$Az = S_iz = By = T_jy$ so $Az = Ax$ and $w = Ax = S_ix$ is the unique point of coincidence of A and S_i

By lemma 1.15, w is the only common fixed point of A and S_i

Similarly there is a unique point $z \in X$ such that $z = Bz = T_jz$

Uniqueness: Assume that $w \neq z$

We have

$$M(w, z, qt) = M(Aw, Bz, t)$$

$$\begin{aligned}
 &\geq \Psi \left(\min \left\{ \begin{aligned} &\{M(S_1w, T_jz, t)\}_{j=1}^m, \{M(S_2w, T_jz, t)\}_{j=1}^m, \dots, \{M(S_nw, T_jz, t)\}_{j=1}^m \\ &\{M(S_iz, Aw, t)\}_{i=1}^n, \{M(Bz, T_jz, t)\}_{j=1}^m, \\ &\frac{\sum_{j=1}^m \alpha_j M(Aw, T_jz, t) + \sum_{i=1}^n \beta_i M(Bz, S_iz, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \end{aligned} \right\} \right) \\
 &= \Psi \left(\min \left\{ M(w, z, t), M(w, w, t), M(z, z, t), \frac{\sum_{j=1}^m \alpha_j M(w, z, t) + \sum_{i=1}^n \beta_i M(z, w, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \right\} \right) \\
 &= \Psi \left(\min \left\{ M(w, z, t), 1, 1, M(w, z, t) \left(\frac{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i} \right) \right\} \right) \\
 &= \Psi(\min\{M(w, z, t), 1, 1, M(w, z, t)\}) \\
 &= \Psi(M(w, z, t)) \\
 &> M(w, z, t)
 \end{aligned}$$

a contradiction.

Therefore we have $z = w$ by lemma 1.15, z is a common fixed point of $A, B, \{S_i\}_{i=1}^n$ and $\{T_j\}_{j=1}^m$. The uniqueness of the fixed point holds from (3).

Corollary 2.4 [25]: Let $(X, M, *)$ be a complete fuzzy metric space and let A, B, S and T be self-mappings of X . Let the pairs $\{A, S\}$ and $\{B, T\}$ be occasionally weakly compatible.

If there exists $q \in (0,1)$ such that

$$M(Ax, By, t) \geq \Psi \left(\min \left\{ M(Sx, Ty, t), M(Sx, Ax, t), M(By, Ty, t), \left(\frac{M(Ax, Ty, t) + M(By, Sx, t)}{2} \right) \right\} \right) \quad (4)$$

for all $x, y \in X$ and $\Psi: [0,1] \rightarrow [0,1]$ such that $0 < t < 1$, then there exists a unique common fixed point of A, B, S and T .

Proof 2.4: Take $n=m=1$ and $\alpha_1 = \beta_1 = 1$ in the above theorem 2.1, the corollary follows.

Theorem 2.5: Let $(X, M, *)$ be complete fuzzy metric space and let A, B and $\{S_i\}_{i=1}^n$ and $\{T_j\}_{j=1}^m$ be self-mappings of X .

Let pairs $\{A, S_i\}_{i=1}^n$ and $\{B, T_j\}_{j=1}^m$ be occasionally weakly compatible at x and y respectively.

If there exists $q \in (0,1)$ such that

$$M(Ax, By, qt) \geq \Psi \left\{ \begin{array}{l} \left\{ M(S_1x, T_jy, t) \right\}_{j=1}^m, \left\{ M(S_2x, T_jy, t) \right\}_{j=1}^m, \dots, \left\{ M(S_nx, T_jy, t) \right\}_{j=1}^m, \left\{ M(S_ix, Ax, t) \right\}_{i=1}^n, \\ \left\{ M(By, T_jy, t) \right\}_{j=1}^m, \frac{\sum_{j=1}^m \alpha_j M(Ax, T_jy, t) + \sum_{i=1}^n \beta_i M(By, S_ix, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i}, \left\{ M(By, S_ix, t) \right\}_{i=1}^n \end{array} \right\} \quad (5)$$

for all $x, y \in X$, for all $t > 0$ and α_j 's ≥ 0 and β_i 's ≥ 0 with atleast one of α_j 's or one of β_i 's is non-zero, and $\Psi: [0,1]^{nm+2n+m+1} \rightarrow [0,1]$ such that $\Psi(t, t, \dots, t_{(nm \text{ times})}, 1, 1, \dots, 1_{(n \text{ times})}, 1, 1, \dots, 1_{(m \text{ times})}, t, t, \dots, t_{(n \text{ times})}, t) > t$ for all $0 < t < 1$, then there exists a unique common fixed point of $A, B, \{S_i\}_{i=1}^n$ and $\{T_j\}_{j=1}^m$.

Proof 2.5: Let the pairs $\{A, S_i\}_{i=1}^n$ and $\{B, T_j\}_{j=1}^m$ be occasionally weakly compatible at the points $x, y \in X$ respectively.

Then $Ax = S_ix$ and $By = T_jy$ for all i and j

Claim: $Ax = By$

If not,

$$\begin{aligned} M(Ax, By, qt) &\geq \Psi \left\{ \begin{array}{l} \left\{ M(S_1x, T_jy, t) \right\}_{j=1}^m, \left\{ M(S_2x, T_jy, t) \right\}_{j=1}^m, \dots, \left\{ M(S_nx, T_jy, t) \right\}_{j=1}^m, \left\{ M(S_ix, Ax, t) \right\}_{i=1}^n, \\ \left\{ M(By, T_jy, t) \right\}_{j=1}^m, \frac{\sum_{j=1}^m \alpha_j M(Ax, T_jy, t) + \sum_{i=1}^n \beta_i M(By, S_ix, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i}, \left\{ M(By, S_ix, t) \right\}_{i=1}^n \end{array} \right\} \\ &= \Psi \left\{ \begin{array}{l} M(Ax, By, t), \dots, M(Ax, By, t)_{(nm \text{ times})}, M(Ax, Ax, t), \dots, M(Ax, Ax, t)_{(n \text{ times})}, \\ M(By, By, t), \dots, M(By, By, t)_{(m \text{ times})}, \frac{\sum_{j=1}^m \alpha_j M(Ax, By, t) + \sum_{i=1}^n \beta_i M(By, Ax, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i}, \\ M(By, Ax, t), \dots, M(By, Ax, t)_{(n \text{ times})} \end{array} \right\} \\ &= \Psi \left\{ \begin{array}{l} M(Ax, By, t), \dots, M(Ax, By, t)_{(nm \text{ times})}, 1, 1, \dots, 1_{(n+m \text{ times})}, M(Ax, By, t), \dots, \\ M(Ax, By, t)_{(n+1 \text{ times})} \end{array} \right\} \\ &> M(Ax, By, t) \end{aligned}$$

a contradiction.

Therefore $Ax = By$

i.e., $Ax = S_ix = By = T_jy$

Suppose that there is a another point z such that $Az = S_iz$ then by (5) we have $Az = S_iz = By = T_jy$ so $Az = Ax$ and $w = Ax = S_ix$ is the unique point of coincidence of A and S_i

By lemma 1.15, w is the only common fixed point of A and S_i

Similarly there is a unique point $z \in X$ such that $z = Bz = T_j z$

Uniqueness: Assume that $w \neq z$

We have

$$\begin{aligned}
 M(w, z, qt) &= M(Aw, Bz, t) \\
 &\geq \Psi \left\{ \begin{array}{l} \{M(S_1 w, T_j z, t)\}_{j=1}^m, \{M(S_2 w, T_j z, t)\}_{j=1}^m, \dots, \{M(S_n w, T_j z, t)\}_{j=1}^m \\ \{M(S_i w, Aw, t)\}_{i=1}^n, \{M(Bz, T_j z, t)\}_{j=1}^m, \\ \frac{\sum_{j=1}^m \alpha_j M(Aw, T_j z, t) + \sum_{i=1}^n \beta_i M(Bz, S_i w, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i}, \{M(Bz, S_i w, t)\}_{i=1}^n \end{array} \right\} \\
 &= \Psi \left\{ \begin{array}{l} M(w, z, t), \dots, M(w, z, t)_{(nm \text{ times})}, M(w, w, t), \dots, M(w, w, t)_{(n \text{ times})}, \\ M(z, z, t), \dots, M(z, z, t)_{(m \text{ times})}, \frac{\sum_{j=1}^m \alpha_j M(w, z, t) + \sum_{i=1}^n \beta_i M(z, w, t)}{\sum_{j=1}^m \alpha_j + \sum_{i=1}^n \beta_i}, \\ M(w, z, t), \dots, M(w, z, t)_{(n \text{ times})} \end{array} \right\} \\
 &= \Psi \left\{ \begin{array}{l} M(w, z, t), \dots, M(w, z, t)_{(nm \text{ times})}, 1, 1, \dots, 1_{(n+m \text{ times})}, M(w, z, t), \dots, \\ M(w, z, t)_{(n+1 \text{ times})} \end{array} \right\} \\
 &> M(w, z, t)
 \end{aligned}$$

a contradiction.

Therefore we have $z = w$ by lemma 1.15, z is a common fixed point of $A, B, \{S_i\}_{i=1}^n$ and $\{T_j\}_{j=1}^m$. The uniqueness of the fixed point holds from (5).

Corollary 2.6 [25]: Let $(X, M, *)$ be a complete fuzzy metric space and let A, B, S and T be self-mappings of X . Let the pairs $\{A, S\}$ and $\{B, T\}$ be occasionally weakly compatible.

If there exists $q \in (0, 1)$ such that

$$M(Ax, By, t) \geq \Psi \left(M(Sx, Ty, t), M(Sx, Ax, t), M(By, Ty, t), \frac{M(Ax, Ty, t) + M(By, Sx, t)}{2}, M(By, Sx, t) \right) \quad (6)$$

for all $x, y \in X$ and $\Psi: [0, 1]^5 \rightarrow [0, 1]$ such that $\Psi(t, 1, 1, t, t) > t$ for all $0 < t < 1$, then there exists a unique common fixed point of A, B, S and T .

Proof 2.6: Take $n=m=1$ and $\alpha_1 = \beta_1 = 1$ in the above theorem 2.5, the corollary follows.

REFERENCES

- [1] A. Al Thagafi and Naseer Shahzad, Generalized I-Non expansive Selfmaps And Invariant Approximations, Acta Mathematica Sinica, English Series May, 2008, Vol.24, No.5, pp.867-876.
- [2] P. Balasubramaniam, S. Muralisankar, R.P. Pant, "Common fixed points of four mappings in a fuzzy metric space", J. Fuzzy Math . 10(2) (2002), 379-384.
- [3] Y.J. Cho, H.K. Pathak, S.M. Kang, J.S. Jung "common on fixed points of compatible maps of type (A) on fuzzy metric spaces", Fuzzy sets and systems 93 (1998), 99-111.
- [4] A. George, P. Veeramani "on some results in fuzzy metric spaces" Fuzzy sets and systems 64(1994), 395-399.
- [5] M. Grabiec , "Fixed points in fuzzy metric spaces" Fuzzy sets and systems 27 (1988), 385-389.
- [6] O. Hadzic, "Common fixed point for theorems for families of mapping in complete metric Space", Math. Japon. 29 (1984), 127-134.

- [7] Mohd.Imdad and Javid Ali, "Some common fixed point theorems in fuzzy metric space", *Mathematical Communications* 11 (2006), 153-163.
- [8] G. Jungck, "Compatible mappings and common fixed points(2)", *Internet J. Math. Sci* (1988), 285-288.
- [9] G. Jungck and B.E. Rhoades, "Fixed point set valued functions without continuity" *Indian J. Pure .Appl. Math.*, 29(3),(1998),pp.771-779.
- [10] G, Jungck and B.E. Rhoades "Fixed point theorems for occasionally weakly compatible mappings", *Fixed point theory*, Vol.7, No.2, 2006, 287-296.
- [11] G, Jungck and B.E. Rhoades "Fixed point theorems for occasionally weakly compatible mappings", *Erratum*, *Fixed point theory*, Vol.9, No.1, 2008, 383-384.
- [12] O. Kramosil and J. Michalek, "Fuzzy metric and statistical metric spaces", *Kybernetika*, 11(1975), 326-334.
- [13] S.Kutukcu, "A fixed point theorem for contraction type mappings in Menger spaces", *Am. J. Appl. Sci.* 4(6) (2007), 371-373.
- [14] Servet Kutuchu, Sushil Sharma and Hanifi Tokgoz, "A fixed point theorem in fuzzy metric spaces", *Int. Journal of Math. Analysis*, Vol.1, 2007, no.18, 861-872.
- [15] S.N. Mishra, "Common fixed points of compatible mappings in PM spaces", *Math. Japon.* 36(1991), 283-289.
- [16] R.P. Pant, "Common fixed points of four mappings", *Bull.Cal.Math.Soc.*90 (1998), 281-286.
- [17] R.P. Pant, "Common fixed point theorems for contractive maps", *J.Math.Appl.*226 (1998), 251-258.
- [18] R.P. Pant, K. Jha, "A remark on common fixed points of four mappings in fuzzy metric space", *Fuzzy Math.*12(2) (2004), 433-437.
- [19] H.K. Pathak and Prachi Singh, "Common fixed point theorem for weakly compatible mapping", *International Mathematical forum*, 2,2007, no.57, 2831-2839.
- [20] B.E. Rhoades, "Contractive definitions and continuity", *Contemporary Math.*72 (1988), 233-245.
- [21] B. Schweizer and A. Sklar, "Statistical metric spaces", *Pacific J.Math.*10 (1960), 313-334.
- [22] Seong Hoon Cho, "On common fixed point in fuzzy metric space", *Int. Math Forum*, 1, 2006, 10,471-479.
- [23] R. Vauki, "Common fixed point for R-weakly commuting maps in fuzzy metric spaces", *Indian J .Pure Appl. Math.* 30 (1999), 419-423.
- [24] L.A. Zadeh, *Fuzzy sets*, *Inform and Control* 8(1965), 338-353.
- [25] Rajesh Kumar Mishra, Sanjay Choudhary "On fixed point theorems in fuzzy metric spaces", *IMACST*, Vol.1, 2010, no.1, 45-47.

Source of support: Nil, Conflict of interest: None Declared