

SOME RESULTS RELATED TO CAUCHY'S PROPER BOUND FOR THE ZEROS  
OF ENTIRE FUNCTIONS OF ORDER ZERO

Sanjib Kumar Datta<sup>1\*</sup> & Manab Biswas<sup>2</sup>

<sup>1</sup>Department of Mathematics, University of Kalyani, Kalyani,  
Dist-Nadia, Pin-741235, West Bengal, India.

<sup>2</sup>Barabilla High School, P.O. Haptiagach, Dist-Uttar Dinajpur, Pin-733202, West Bengal, India.

(Received on: 24-06-13; Revised & Accepted on: 23-10-13)

ABSTRACT

A single valued function of one complex variable which is analytic in the finite complex plane is called an entire function. In this paper we would like to establish the bounds for the moduli of zeros of entire functions of order zero. Some examples are provided to clear the notions.

AMS Subject Classification 2010: Primary 30C15, 30C10, Secondary 26C10.

Keywords and phrases: Zeros of entire functions of order zero, Cauchy's bound, Proper ring shaped region.

1. INTRODUCTION, DEFINITIONS AND NOTATIONS

Let  $P(z) = a_0 + a_1z + a_2z^2 + a_3z^3 + \dots + a_{n-1}z^{n-1} + a_nz^n$ ;  $|a_n| \neq 0$  be a polynomial of degree  $n$ . Datt and Govil [2]; Govil and Rahaman [5]; Marden [9]; Mohammad [10]; Chattopadhyay, Das, Jain and Konwar [1]; Joyal, Labelle and Rahaman [6]; Jain { [7], [8] }; Sun and Hsieh [11]; Zilovic, Roytman, Combettes and Swamy [13]; Das and Datta [4] etc. worked in the theory of the distribution of the zeros of polynomials and obtained some newly developed results.

In this paper we intend to establish some sharper results concerning the theory of distribution of zeros of entire functions of order zero.

The following definitions are well known:

**Definition: 1** The order  $\rho$  and lower order  $\lambda$  of a meromorphic function  $f$  are defined as

$$\rho = \limsup_{r \rightarrow \infty} \frac{\log T(r,f)}{\log r} \quad \text{and} \quad \lambda = \liminf_{r \rightarrow \infty} \frac{\log T(r,f)}{\log r}.$$

If  $f$  is entire, one can easily verify that

$$\rho = \limsup_{r \rightarrow \infty} \frac{\log [2]M(r,f)}{\log r} \quad \text{and} \quad \lambda = \liminf_{r \rightarrow \infty} \frac{\log [2]M(r,f)}{\log r},$$

where  $\log^{[k]} x = \log(\log^{[k-1]} x)$  for  $k = 1, 2, 3, \dots$  and  $\log^{[0]} x = x$ .

If  $\rho < \infty$  then  $f$  is of finite order. Also  $\rho = 0$  means that  $f$  is of order zero. In this connection Datta and Biswas [3] gave the following definition:

**Definition: 2** Let  $f$  be a meromorphic function of order zero. Then the quantities  $\rho^*$  and  $\lambda^*$  of  $f$  are defined by:

$$\rho^* = \limsup_{r \rightarrow \infty} \frac{T(r,f)}{\log r} \quad \text{and} \quad \lambda^* = \liminf_{r \rightarrow \infty} \frac{T(r,f)}{\log r}.$$

Corresponding author: Sanjib Kumar Datta<sup>1\*</sup>

<sup>1</sup>Department of Mathematics, University of Kalyani, Kalyani,  
Dist-Nadia, Pin-741235, West Bengal, India.

If  $f$  is an entire function then clearly

$$\rho^* = \limsup_{r \rightarrow \infty} \frac{\log M(r,f)}{\log r} \quad \text{and} \quad \lambda^* = \liminf_{r \rightarrow \infty} \frac{\log M(r,f)}{\log r}.$$

## 2. LEMMAS

In this section we present a lemma which will be needed in the sequel.

**Lemma: 1** If  $f(z)$  is an entire function of order  $\rho = 0$ , then for every  $\varepsilon > 0$  the inequality  $N(r) \leq (\log r)^{\rho^* + \varepsilon}$  holds for all sufficiently large  $r$  where  $N(r)$  is the number of zeros of  $f(z)$  in  $|z| \leq \log r$ .

**Proof:** Let us suppose that  $f(z) = 1$ . This supposition can be made without loss of generality because if  $f(z)$  has a zero of order ' $m$ ' at the origin then we may consider  $g(z) = c \frac{f(z)}{z^m}$  where  $c$  is so chosen that  $g(0) = 1$ . Since the function  $g(z)$  and  $f(z)$  have the same order therefore it will be unimportant for our investigations that the number of zeros of  $g(z)$  and  $f(z)$  differ by  $m$ .

We further assume that  $f(z)$  has no zeros on  $|z| = \log 2r$  and the zeros  $z_i$ 's of  $f(z)$  in  $|z| < \log r$  are in non decreasing order of their moduli so that  $|z_i| \leq |z_{i+1}|$ . Also let  $\rho^*$  suppose to be finite where  $\rho = 0$  is the zero of order of  $f(z)$ .

Now we shall make use of Jensen's formula as state below

$$\log|f(0)| = - \sum_{i=1}^n \log \frac{R}{|z_i|} + \frac{1}{2\pi} \int_0^{2\pi} \log|f(R e^{i\phi})| d\phi. \quad (1)$$

Let us replace  $R$  by  $2r$  and  $n$  by  $N(2r)$  in (1).

$$\therefore \log|f(0)| = - \sum_{i=1}^{N(2r)} \log \frac{2r}{|z_i|} + \frac{1}{2\pi} \int_0^{2\pi} \log|f(2r e^{i\phi})| d\phi.$$

Since  $f(0) = 1$ ,  $\therefore \log|f(0)| = \log 1 = 0$ .

$$\therefore \sum_{i=1}^{N(2r)} \log \frac{2r}{|z_i|} = \frac{1}{2\pi} \int_0^{2\pi} \log|f(2r e^{i\phi})| d\phi \quad (2)$$

$$\text{L.H.S.} = \sum_{i=1}^{N(2r)} \log \frac{2r}{|z_i|} \geq \sum_{i=1}^{N(r)} \log \frac{2r}{|z_i|} \geq N(r) \log 2 \quad (3)$$

because for large values of  $r$ ,

$$\log \frac{2r}{|z_i|} \geq \log 2.$$

$$\begin{aligned} \text{R.H.S.} &= \frac{1}{2\pi} \int_0^{2\pi} \log|f(2r e^{i\phi})| d\phi \\ &\leq \frac{1}{2\pi} \int_0^{2\pi} \log M(2r) d\phi = \log M(2r). \end{aligned} \quad (4)$$

Again by definition of order  $\rho^*$  of  $f(z)$  we have for every  $\varepsilon > 0$ ,

$$\log M(2r) \leq \{\log(2r)\}^{\rho^* + \varepsilon/2}. \quad (5)$$

Hence from (2) by the help of (3), (4) and (5) we have

$$N(r) \log 2 \leq (\log 2r)^{\rho^* + \varepsilon/2}$$

$$\text{i.e., } N(r) \leq \frac{(\log 2)^{\rho^* + \varepsilon/2}}{\log 2} \cdot \frac{(\log r)^{\rho^* + \varepsilon}}{(\log r)^{\varepsilon/2}} \leq (\log r)^{\rho^* + \varepsilon}.$$

This proves the lemma.

### 3. THEOREMS

In this section we present the main results of the paper.

**Theorem: 1** Let  $P(z)$  be an entire function defined by

$$P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n + \dots$$

whose order  $\rho = 0$ . Also for all sufficiently large  $r$  in the disc  $|z| \leq \log r$ ,  $|a_{N(r)}| \neq 0$ ,  $|a_0| \neq 0$  and also  $a_n \rightarrow 0$  as  $n > N(r)$ . Then all the zeros of  $P(z)$  lie in the ring shaped region

$$\frac{1}{t_0} \leq |z| \leq t_0$$

where  $t_0$  is the greatest positive root of

$$g(t) \equiv |a_{N(r)}|t^{N(r)+1} - (|a_{N(r)}| + M)t^{N(r)} + M = 0$$

and  $t'_0$  is the greatest positive root of

$$f(t) \equiv |a_0|t^{N(r)+1} - (|a_0| + M')t^{N(r)} + M' = 0$$

where  $M = \max\{|a_0|, |a_1|, \dots, |a_{N(r)-1}|\}$  and  $M' = \max\{|a_1|, |a_2|, \dots, |a_{N(r)}|\}$ .

**Proof:** Now,  $P(z) \approx a_0 + a_1z + a_2z^2 + \dots + a_{N(r)}z^{N(r)}$  because  $N(r)$  exists for  $|z| \leq \log r$ ;  $r$  is sufficiently large and  $a_n \rightarrow 0$  as  $n > N(r)$ . Then all the zeros of  $P(z)$  lie in the ring shaped region given in Theorem 1 which we are to prove.

Now

$$\begin{aligned} |P(z)| &\approx |a_0 + a_1z + a_2z^2 + \dots + a_{N(r)}z^{N(r)}| \\ &\geq |a_{N(r)}||z|^{N(r)} - |a_0 + a_1z + a_2z^2 + \dots + a_{N(r)-1}z^{N(r)-1}|. \end{aligned}$$

Also

$$\begin{aligned} |a_0 + a_1z + a_2z^2 + \dots + a_{N(r)-1}z^{N(r)-1}| &\leq |a_0| + \dots + |a_{N(r)-1}||z|^{N(r)-1} \\ &\leq M(1 + |z| + \dots + |z|^{N(r)-1}) \\ &= M \frac{|z|^{N(r)} - 1}{|z| - 1} \text{ if } |z| \neq 1. \end{aligned} \tag{6}$$

Therefore using (6) we obtain that

$$\begin{aligned} |P(z)| &\geq |a_{N(r)}||z|^{N(r)} - |a_0 + a_1z + a_2z^2 + \dots + a_{N(r)-1}z^{N(r)-1}| \\ &\geq |a_{N(r)}||z|^{N(r)} - M \frac{|z|^{N(r)} - 1}{|z| - 1}. \end{aligned}$$

Hence

$$|P(z)| \geq 0 \text{ if } |a_{N(r)}||z|^{N(r)} - M \frac{|z|^{N(r)} - 1}{|z| - 1} > 0$$

$$\text{i.e., if } |a_{N(r)}||z|^{N(r)} > M \frac{|z|^{N(r)} - 1}{|z| - 1}$$

$$\text{i.e., if } |a_{N(r)}||z|^{N(r)+1} - |a_{N(r)}||z|^{N(r)} > M(|z|^{N(r)} - 1)$$

$$\text{i.e., if } |a_{N(r)}||z|^{N(r)+1} - |a_{N(r)}||z|^{N(r)} - M|z|^{N(r)} + M > 0$$

$$\text{i.e., if } |a_{N(r)}||z|^{N(r)+1} - (|a_{N(r)}| + M)|z|^{N(r)} + M > 0.$$

Therefore on  $|z| \neq 1$ ,

$$|P(z)| \geq 0 \text{ if } |a_{N(r)}||z|^{N(r)+1} - (|a_{N(r)}| + M)|z|^{N(r)} + M > 0.$$

Now let us consider

$$g(t) \equiv |a_{N(r)}|t^{N(r)+1} - (|a_{N(r)}| + M)t^{N(r)} + M = 0. \quad (7)$$

Clearly the maximum number of changes in sign in (7) is two. So the maximum number of positive roots of  $g(t) = 0$  is two and by Descartes' rule of sign if it is less, less by two. Clearly  $t = 1$  is one positive root of (7). So  $g(t) = 0$  must have another positive root  $t_1$  (say). Let us take  $t_0 = \max\{1, t_1\}$ . Clearly for  $t > t_0$ ,  $g(t) > 0$ . If not, for some  $t = t_2 > t_0$ ,  $g(t_2) < 0$ .

Now  $g(t_2) < 0$  and  $g(\infty) > 0$  imply that  $g(t) = 0$  has another positive root in  $(t_2, \infty)$  which gives a contradiction.

Therefore for  $t > t_0$ ,  $g(t) > 0$  and so  $t_0 > 1$ .

Hence  $|P(z)| \geq 0$  for  $|z| > t_0$ .

Therefore all the zeros of  $P(z)$  lie in the disc  $|z| \leq t_0$ . (8)

Again let us consider

$$\begin{aligned} Q(z) &= z^{N(r)} P\left(\frac{1}{z}\right) \\ &\approx z^{N(r)} \left\{ a_0 + \frac{a_1}{z} + \dots + \frac{a_{N(r)}}{z^{N(r)}} \right\} \\ &= a_0 z^{N(r)} + a_1 z^{N(r)-1} + \dots + a_{N(r)} \end{aligned}$$

i.e.,  $|Q(z)| \geq |a_0||z|^{N(r)} - |a_1 z^{N(r)-1} + \dots + a_{N(r)}|$  for  $|z| \neq 1$ .

Now

$$\begin{aligned} |a_1 z^{N(r)-1} + \dots + a_{N(r)}| &\leq |a_1||z|^{N(r)-1} + \dots + |a_{N(r)}| \\ &\leq M'(|z|^{N(r)-1} + \dots + 1) \\ &= M' \left( \frac{|z|^{N(r)} - 1}{|z| - 1} \right) \text{ for } |z| \neq 1. \end{aligned} \quad (9)$$

Using (9) we get that

$$\begin{aligned} |Q(z)| &\geq |a_0||z|^{N(r)} - |a_1 z^{N(r)-1} + \dots + a_{N(r)}| \\ &\geq |a_0||z|^{N(r)} - M' \left( \frac{|z|^{N(r)} - 1}{|z| - 1} \right) \text{ for } |z| \neq 1. \end{aligned}$$

Therefore for  $|z| \neq 1$ ,

$$|Q(z)| \geq 0 \text{ if } |a_0||z|^{N(r)} - M' \left( \frac{|z|^{N(r)} - 1}{|z| - 1} \right) > 0$$

$$\text{i.e., if } |a_0||z|^{N(r)} > M' \left( \frac{|z|^{N(r)} - 1}{|z| - 1} \right)$$

$$\text{i.e., if } |a_0||z|^{N(r)+1} - |a_0||z|^{N(r)} - M'|z|^{N(r)} + M' > 0$$

$$\text{i.e., if } |a_0||z|^{N(r)+1} - (|a_0| + M')|z|^{N(r)} + M' > 0.$$

So for  $|z| \neq 1$ ,

$$|Q(z)| \geq 0 \text{ if } |a_0||z|^{N(r)+1} - (|a_0| + M')|z|^{N(r)} + M' > 0.$$

Let us consider

$$f(t) \equiv |a_0|t^{N(r)+1} - (|a_0| + M')t^{N(r)} + M' = 0.$$

Since the maximum number of changes in sign in  $f(t)$  is two, the maximum number of positive roots of  $f(t) = 0$  is two and by Descartes' rule of sign if it is less, less by two. Clearly  $t = 1$  is one positive root of  $f(t) = 0$ . So  $f(t) = 0$  must have another positive root  $t_2$  (say). Let us take  $t'_0 = \max \{1, t_2\}$ . Clearly for  $t > t'_0$ ,  $f(t) > 0$ . If not, for some  $t_3 > t'_0$ ,  $f(t_3) < 0$ .

Now  $f(t_3) < 0$  and  $f(\infty) > 0$  implies that  $f(t) = 0$  has another positive root in  $(t_3, \infty)$  which is a contradiction.

Therefore for  $t > t'_0$ ,  $f(t) > 0$ .

Also  $t'_0 \geq 1$ . So  $|Q(z)| \geq 0$  for  $|z| > t'_0$ .

Therefore  $Q(z)$  does not vanish in  $|z| > t'_0$ .

Hence all the zeros of  $Q(z)$  lie in  $|z| \leq t'_0$ .

Let  $z = z_0$  be a zero of  $P(z)$ . Therefore  $P(z_0) = 0$ . Clearly  $z_0 \neq 0$  as  $a_0 \neq 0$ . Putting  $z = \frac{1}{z_0}$  in  $Q(z)$  we get that  $Q\left(\frac{1}{z_0}\right) = \left(\frac{1}{z_0}\right)^n \cdot P(z_0) = \left(\frac{1}{z_0}\right)^n \cdot 0 = 0$ .

Therefore  $Q\left(\frac{1}{z_0}\right) = 0$ . So  $z = \frac{1}{z_0}$  is a root of  $Q(z) = 0$ . Hence  $\left|\frac{1}{z_0}\right| \leq t'_0$  implies that  $|z_0| \geq \frac{1}{t'_0}$ .

As  $z_0$  is an arbitrary root of  $P(z) = 0$ .

Therefore all the zeros of  $P(z)$  lie in  $|z| \geq \frac{1}{t'_0}$ . (10)

From (8) and (10) we get that all the zeros of  $P(z)$  lie in the proper ring shaped region

$$\frac{1}{t'_0} \leq |z| \leq t_0$$

where  $t_0$  and  $t'_0$  are the greatest positive roots of the equations

$$g(t) \equiv |a_{N(r)}|t^{N(r)+1} - (|a_{N(r)}| + M)t^{N(r)} + M = 0$$

and

$$f(t) \equiv |a_0|t^{N(r)+1} - (|a_0| + M')t^{N(r)} + M' = 0$$

where  $M$  and  $M'$  are given in the statement of Theorem 1.

This proves the theorem.

**Remark: 1** The limit in Theorem 1 is attained by  $P(z) = z^2 - z - 1$ . Here  $\rho = 0$ ,  $\rho^* = 2$  and  $N(r) = 2 \leq (\log r)^{2+\varepsilon}$ . For  $\varepsilon > 0$  and sufficiently large  $r$ , all  $a_n = 0$ ,  $n \geq 2$ . Also  $a_0 = -1$ ,  $a_1 = -1$ ,  $a_2 = 1$ .

Therefore

$$M = \max \{|a_0|, |a_1|\} = 1 \text{ and } M' = \max \{|a_1|, |a_2|\} = 1$$

and

$$g(t) \equiv |a_2|t^3 - (|a_2| + M)t^2 + M = 0$$

$$\text{i.e., } g(t) \equiv t^3 - (1 + 1)t^2 + 1 = 0$$

$$\text{i.e., } g(t) \equiv t^3 - 2t^2 + 1 = 0.$$

Again

$$f(t) \equiv |a_0|t^3 - (|a_0| + M')t^2 + M' = 0$$

$$\text{i.e., } f(t) \equiv 1 \cdot t^3 - (1 + 1)t^2 + 1 = 0$$

$$\text{i.e., } f(t) \equiv t^3 - 2t^2 + 1 = 0.$$

So  $f(t) = 0$  and  $g(t) = 0$  represent the same equation. Maximum number of positive roots of  $f(t) = 0$  and  $g(t) = 0$  are same. Now

$$g(t) = 0$$

$$\text{implies that } t^3 - 2t^2 + 1 = 0.$$

$$\text{i.e., } (t - 1)(t^2 - t - 1) = 0.$$

Therefore

$$t = 1 \text{ and } t = \frac{1 \pm \sqrt{(-1)^2 - 4 \cdot 1 \cdot (-1)}}{2 \cdot 1} = \frac{1 \pm \sqrt{5}}{2}.$$

Hence the positive roots of  $g(t) = 0$  are 1 and  $\frac{1+\sqrt{5}}{2}$ . So

$$t_0 = \max\left\{1, \frac{1+\sqrt{5}}{2}\right\} = \frac{1+\sqrt{5}}{2}.$$

Also the maximum positive root of  $f(t) = 0$  is

$$t'_0 = \max\left\{1, \frac{1 + \sqrt{5}}{2}\right\} = \frac{1 + \sqrt{5}}{2}.$$

So in view of Theorem 1 all the zeros of  $P(z)$  lie in

$$\frac{1}{t_0} \leq |z| \leq t_0$$

$$\text{i.e., } \frac{1}{\frac{1+\sqrt{5}}{2}} \leq |z| \leq \frac{1+\sqrt{5}}{2}$$

$$\text{i.e., } \frac{\sqrt{5}-1}{2} \leq |z| \leq \frac{1+\sqrt{5}}{2}.$$

Now the zeros of  $P(z)$  are given by solving  $z^2 - z - 1 = 0$ . Therefore  $z = \frac{1+\sqrt{5}}{2}$ . Let us denote the zeros of  $P(z)$  by  $z_1 = \frac{1+\sqrt{5}}{2}$  and  $z_2 = \frac{1-\sqrt{5}}{2} = -\frac{\sqrt{5}-1}{2}$ . Clearly  $z_1$  lies on the upper boundary and  $z_2$  lies on the lower boundary. So the best result is given by  $P(z) = z^2 - z - 1$ .

**Theorem: 2** Let  $P(z)$  be an entire function defined by  $P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n + \dots$  with order  $\rho = 0$ ,  $a_{N(r)} \neq 0$ ,  $a_0 \neq 0$  and also  $a_n \rightarrow 0$  for  $n > N(r)$  for the disc  $|z| \leq \log r$ , when  $r$  is sufficiently large. Further for some  $\rho^* > 0$ ,  $|a_0|(\rho^*)^{N(r)} \geq |a_1|(\rho^*)^{N(r)-1} \geq \dots \geq |a_{N(r)-1}|\rho^* \geq |a_{N(r)}|$ .

Then all the zeros of  $P(z)$  lie in the ring shaped region

$$\frac{1}{\rho^*(1 + \frac{|a_1|}{|a_0|\rho^*})} < |z| < \frac{1}{\rho^*} \left(1 + \frac{|a_0|}{|a_{N(r)}|} (\rho^*)^{N(r)}\right).$$

**Proof:** For the given entire function  $P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n + \dots$  with  $a_n \rightarrow 0$  as  $n > N(r)$ , where  $r$  is sufficiently large,  $N(r)$  exists and  $N(r) \leq (\log r)^{\rho^* + \epsilon}$ .

Therefore

$$P(z) \approx a_0 + a_1z + a_2z^2 + \dots + a_{N(r)}z^{N(r)}$$

as  $a_0 \neq 0$ ,  $a_{N(r)} \neq 0$  and  $a_n \rightarrow 0$  for  $n > N(r)$ .

Let us consider

$$\begin{aligned} R(z) &= (\rho^*)^{N(r)} P\left(\frac{z}{\rho^*}\right) \\ &\approx (\rho^*)^{N(r)} \left(a_0 + a_1 \frac{z}{\rho^*} + a_2 \frac{z^2}{(\rho^*)^2} + \dots + a_{N(r)} \frac{z^{N(r)}}{(\rho^*)^{N(r)}}\right) \\ &= a_0 (\rho^*)^{N(r)} + a_1 (\rho^*)^{N(r)-1} z + \dots + a_{N(r)} z^{N(r)}. \end{aligned}$$

Therefore

$$|R(z)| \geq |a_{N(r)}| |z|^{N(r)} - |a_0 (\rho^*)^{N(r)} + a_1 (\rho^*)^{N(r)-1} z + \dots + a_{N(r)-1} \rho^* z^{N(r)-1}|. \quad (11)$$

Now by the given condition  $|a_0|(\rho^*)^{N(r)} \geq |a_1|(\rho^*)^{N(r)-1} \geq \dots \geq |a_{N(r)-1}|\rho^* \geq |a_{N(r)}|$  provided  $|z| \neq 0$ , we obtain that

$$|a_0(\rho^*)^{N(r)} + a_1(\rho^*)^{N(r)-1}z + \dots + a_{N(r)-1}\rho^*z^{N(r)-1}| \leq |a_0|(\rho^*)^{N(r)} + \dots + |a_{N(r)-1}|\rho^*|z|^{N(r)-1} \\ \leq |a_0|(\rho^*)^{N(r)}|z|^{N(r)} \left( \frac{1}{|z|} + \dots + \frac{1}{|z|^{N(r)}} \right).$$

Therefore on  $|z| \neq 0$ ,

$$-|a_0(\rho^*)^{N(r)} + a_1(\rho^*)^{N(r)-1}z + \dots + a_{N(r)-1}\rho^*z^{N(r)-1}| \geq -|a_0|(\rho^*)^{N(r)}|z|^{N(r)} \left( \frac{1}{|z|} + \dots + \frac{1}{|z|^{N(r)}} \right). \quad (12)$$

Therefore using (12) we get from (11) that

$$|R(z)| \geq |a_{N(r)}||z|^{N(r)} - |a_0|(\rho^*)^{N(r)}|z|^{N(r)} \left( \frac{1}{|z|} + \dots + \frac{1}{|z|^{N(r)}} \right) \\ \geq |a_{N(r)}||z|^{N(r)} - |a_0|(\rho^*)^{N(r)}|z|^{N(r)} \left( \frac{1}{|z|} + \dots + \frac{1}{|z|^{N(r)}} + \dots \right) \\ = |z|^{N(r)} \left[ |a_{N(r)}| - |a_0|(\rho^*)^{N(r)} \sum_{k=1}^{\infty} \frac{1}{|z|^k} \right].$$

Clearly  $\sum_{k=1}^{\infty} \frac{1}{|z|^k}$  is a geometric series which is convergent for  $\frac{1}{|z|} < 1$  i.e., for  $|z| > 1$  and converges to

$$\frac{1}{|z|} \frac{1}{1 - \frac{1}{|z|}} = \frac{1}{|z| - 1}.$$

Therefore

$$\sum_{k=1}^{\infty} \frac{1}{|z|^k} = \frac{1}{|z| - 1} \text{ if } |z| > 1.$$

Hence we get from above that for  $|z| > 1$

$$|R(z)| > |z|^{N(r)} \left( |a_{N(r)}| - (\rho^*)^{N(r)}|a_0| \frac{1}{|z| - 1} \right).$$

Now for  $|z| > 1$ ,

$$|R(z)| > 0 \text{ if } |z|^{N(r)} \left( |a_{N(r)}| - (\rho^*)^{N(r)}|a_0| \frac{1}{|z| - 1} \right) \geq 0$$

$$\text{i.e., if } |a_{N(r)}| - (\rho^*)^{N(r)}|a_0| \frac{1}{|z| - 1} \geq 0$$

$$\text{i.e., if } |a_{N(r)}| \geq (\rho^*)^{N(r)} \frac{|a_0|}{|z| - 1}$$

$$\text{i.e., if } |z| - 1 \geq (\rho^*)^{N(r)} \frac{|a_0|}{|a_{N(r)}|}$$

$$\text{i.e., if } |z| \geq 1 + (\rho^*)^{N(r)} \frac{|a_0|}{|a_{N(r)}|} > 1.$$

Therefore

$$|R(z)| > 0 \text{ if } |z| \geq 1 + (\rho^*)^{N(r)} \frac{|a_0|}{|a_{N(r)}|}.$$

So all the zeros of  $R(z)$  lie in

$$|z| < 1 + \frac{|a_0|}{|a_{N(r)}|} (\rho^*)^{N(r)}.$$

Let  $z_0$  be an arbitrary zero of  $P(z)$ . Therefore  $P(z_0) = 0$ . Clearly  $z_0 \neq 0$  as  $a_0 \neq 0$ . Putting  $z = \rho^*z_0$  in  $R(z)$  we have

$$R(\rho^*z_0) = (\rho^*)^{N(r)} \cdot P(z_0) = (\rho^*)^{N(r)} \cdot 0 = 0.$$

Hence  $z = \rho^* z_0$  is a zero of  $R(z)$ . Therefore

$$|\rho^* z_0| < 1 + \frac{|a_0|}{|a_{N(r)}|} (\rho^*)^{N(r)}$$

$$\text{i.e., } |z_0| < \frac{1}{\rho^*} \left( 1 + \frac{|a_0|}{|a_{N(r)}|} (\rho^*)^{N(r)} \right).$$

Since  $z_0$  is any zero of  $P(z)$  therefore all the zeros of  $P(z)$  lie in

$$|z| < \frac{1}{\rho^*} \left( 1 + \frac{|a_0|}{|a_{N(r)}|} (\rho^*)^{N(r)} \right). \quad (13)$$

Again let us consider

$$F(z) = (\rho^*)^{N(r)} z^{N(r)} P\left(\frac{1}{\rho^* z}\right).$$

Now

$$\begin{aligned} F(z) &= (\rho^*)^{N(r)} z^{N(r)} P\left(\frac{1}{\rho^* z}\right) \\ &\approx (\rho^*)^{N(r)} z^{N(r)} \left\{ a_0 + \frac{a_1}{\rho^* z} + \dots + \frac{a_{N(r)}}{(\rho^* z)^{N(r)}} \right\} \\ &= a_0 (\rho^*)^{N(r)} z^{N(r)} + a_1 (\rho^*)^{N(r)-1} z^{N(r)-1} + \dots + a_{N(r)}. \end{aligned}$$

Therefore

$$|F(z)| \geq |a_0| (\rho^*)^{N(r)} |z|^{N(r)} - |a_1 (\rho^*)^{N(r)-1} z^{N(r)-1} + \dots + a_{N(r)}|.$$

Again

$$\begin{aligned} |a_1 (\rho^*)^{N(r)-1} z^{N(r)-1} + \dots + a_{N(r)}| &\leq |a_1| (\rho^*)^{N(r)-1} |z|^{N(r)-1} + \dots + |a_{N(r)}| \\ &\leq |a_1| (\rho^*)^{N(r)-1} (|z|^{N(r)-1} + \dots + |z| + 1) \end{aligned}$$

provided  $|z| \neq 0$ . So

$$|a_1 (\rho^*)^{N(r)-1} z^{N(r)-1} + \dots + a_{N(r)}| \leq |a_1| (\rho^*)^{N(r)-1} |z|^{N(r)} \left( \frac{1}{|z|} + \dots + \frac{1}{|z|^{N(r)}} \right).$$

So for  $|z| \neq 0$ ,

$$\begin{aligned} |F(z)| &\geq |a_0| (\rho^*)^{N(r)} |z|^{N(r)} - |a_1| (\rho^*)^{N(r)-1} |z|^{N(r)} \left( \frac{1}{|z|} + \dots + \frac{1}{|z|^{N(r)}} \right) \\ &= (\rho^*)^{N(r)-1} |z|^{N(r)} \left[ |a_0| \rho^* - |a_1| \left( \frac{1}{|z|} + \dots + \frac{1}{|z|^{N(r)}} \right) \right]. \end{aligned}$$

Therefore for  $|z| \neq 0$ ,

$$|F(z)| > (\rho^*)^{N(r)-1} |z|^{N(r)} \left[ |a_0| \rho^* - |a_1| \sum_{k=1}^{\infty} \frac{1}{|z|^k} \right]. \quad (14)$$

The geometric series  $\sum_{k=1}^{\infty} \frac{1}{|z|^k}$  is convergent for

$$\frac{1}{|z|} < 1$$

i.e., for  $|z| > 1$

and converges to

$$\frac{1}{|z|} \frac{1}{1 - \frac{1}{|z|}} = \frac{1}{|z| - 1}.$$

Therefore

$$\sum_{k=1}^{\infty} \frac{1}{|z|^k} = \frac{1}{|z|-1} \text{ if } |z| > 1. \quad (15)$$



Using (14) and (15) we have for  $|z| > 1$ ,

$$|F(z)| > (\rho^*)^{N(r)-1} |z|^{N(r)} \left[ |a_0| \rho^* - \frac{|a_1|}{|z|-1} \right].$$

Hence for  $|z| > 1$ ,

$$|F(z)| > 0 \text{ if } (\rho^*)^{N(r)-1} |z|^{N(r)} \left[ |a_0| \rho^* - \frac{|a_1|}{|z|-1} \right] \geq 0$$

$$\text{i.e., if } |a_0| \rho^* - \frac{|a_1|}{|z|-1} \geq 0$$

$$\text{i.e., if } |a_0| \rho^* \geq \frac{|a_1|}{|z|-1}$$

$$\text{i.e., if } |z| \geq 1 + \frac{|a_1|}{|a_0| \rho^*} > 1.$$

Therefore

$$|F(z)| > 0 \text{ for } |z| \geq 1 + \frac{|a_1|}{|a_0| \rho^*}.$$

So  $F(z)$  does not vanish in

$$|z| \geq 1 + \frac{|a_1|}{|a_0| \rho^*}.$$

Equivalently all the zeros of  $F(z)$  lie in

$$|z| < 1 + \frac{|a_1|}{|a_0| \rho^*}.$$

Let  $z = z_0$  be any zero of  $P(z)$ . Therefore  $P(z_0) = 0$ . Clearly  $a_0 \neq 0$  and  $z_0 \neq 0$ .

Now let us put  $z = \frac{1}{\rho^* z_0}$  in  $F(z)$ . So we have

$$\begin{aligned} F\left(\frac{1}{\rho^* z_0}\right) &= (\rho^*)^{N(r)} \left(\frac{1}{\rho^* z_0}\right)^{N(r)} \cdot P(z_0) \\ &= \left(\frac{1}{z_0}\right)^{N(r)} \cdot 0 = 0 \end{aligned}$$

Therefore  $z = \frac{1}{\rho^* z_0}$  is a root of  $F(z)$ .

Hence

$$\left| \frac{1}{\rho^* z_0} \right| < 1 + \frac{|a_1|}{|a_0| \rho^*}$$

$$\text{i.e., } \left| \frac{1}{z_0} \right| < \rho^* \left( 1 + \frac{|a_1|}{|a_0| \rho^*} \right),$$

$$\text{i.e., } |z_0| > \frac{1}{\rho^* \left( 1 + \frac{|a_1|}{|a_0| \rho^*} \right)},$$

As  $z_0$  is an arbitrary zero of  $P(z)$ , all the zeros of  $P(z)$  lie on

$$|z| > \frac{1}{\rho^* \left( 1 + \frac{|a_1|}{|a_0| \rho^*} \right)}. \tag{16}$$

From (13) and (16) we get that all the zeros of  $P(z)$  lie in the proper ring shaped region

$$\frac{1}{\rho^* \left( 1 + \frac{|a_1|}{|a_0| \rho^*} \right)} < |z| < \frac{1}{\rho^*} \left( 1 + \frac{|a_0|}{|a_{N(r)}|} (\rho^*)^{N(r)} \right)$$

where

$$|a_0|(\rho^*)^{N(r)} \geq |a_1|(\rho^*)^{N(r)-1} \geq \dots \geq |a_{N(r)}|$$

for some  $\rho^* > 0$ .

This proves the theorem.

**Corollary: 1** From Theorem 2 we can easily conclude that all the zeros of  $P(z) = a_0 + a_1z + \dots + a_nz^n$  of degree  $n$ ,  $|a_n| \neq 0$  with the property  $|a_0| \geq |a_1| \geq \dots \geq |a_n|$  lie in the proper ring shaped region

$$\frac{1}{\left(1 + \frac{|a_1|}{|a_0|}\right)} < |z| < \left(1 + \frac{|a_0|}{|a_n|}\right)$$

just on putting  $\rho^* = 1$ .

**Theorem: 3** Let  $P(z)$  be an entire function with order  $\rho = 0$ . For sufficiently large values of  $r$  in the disc  $|z| \leq \log r$ , the Taylor's series expansion of  $P(z)$

$$P(z) = a_0 + a_{p_1}z^{p_1} + a_{p_2}z^{p_2} + \dots + a_{p_m}z^{p_m} + a_{N(r)}z^{N(r)}, \quad a_0 \neq 0 \text{ be such that } 1 \leq p_1 < p_2 < \dots < p_m \leq N(r) - 1, p_i \text{'s are integers and for some } \rho^* > 0,$$

$$|a_0|(\rho^*)^{N(r)} \geq |a_{p_1}|(\rho^*)^{N(r)-p_1} \geq \dots \geq |a_{p_m}|(\rho^*)^{N(r)-p_m} \geq |a_{N(r)}|.$$

Then all the zeros of  $P(z)$  lie in the proper ring shaped region

$$\frac{1}{\rho^* t_0} < |z| < \frac{1}{\rho^*} t_0$$

where  $t_0$  and  $t_0'$  are the unique positive roots of the equations

$$g(t) \equiv |a_{N(r)}|t^{N(r)-p_m} - |a_{N(r)}|t^{N(r)-p_m-1} - |a_0|(\rho^*)^{N(r)} = 0$$

and

$$f(t) \equiv |a_0|(\rho^*)^{p_1}t^{p_1} - |a_0|(\rho^*)^{p_1}t^{p_1-1} - |a_{p_1}| = 0$$

respectively.

**Proof:** Let

$$P(z) = a_0 + a_{p_1}z^{p_1} + a_{p_2}z^{p_2} + \dots + a_{p_m}z^{p_m} + a_{N(r)}z^{N(r)}, \quad |a_{N(r)}| \neq 0. \tag{17}$$

Also for some  $\rho^* > 0$ ,

$$|a_0|(\rho^*)^{N(r)} \geq |a_{p_1}|(\rho^*)^{N(r)-p_1} \geq \dots \geq |a_{p_m}|(\rho^*)^{N(r)-p_m} \geq |a_{N(r)}|.$$

Let us consider

$$\begin{aligned} R(z) &= (\rho^*)^{N(r)} P\left(\frac{z}{\rho^*}\right) \\ &= (\rho^*)^{N(r)} \left\{ a_0 + a_{p_1} \frac{z^{p_1}}{(\rho^*)^{p_1}} + \dots + a_{p_m} \frac{z^{p_m}}{(\rho^*)^{p_m}} + a_{N(r)} \frac{z^{N(r)}}{(\rho^*)^{N(r)}} \right\} \\ &= a_0(\rho^*)^{N(r)} + a_{p_1}(\rho^*)^{N(r)-p_1}z^{p_1} + \dots + a_{p_m}(\rho^*)^{N(r)-p_m}z^{p_m} + a_{N(r)}z^{N(r)}. \end{aligned}$$

Therefore

$$|R(z)| \geq |a_{N(r)}z^{N(r)}| - |a_0(\rho^*)^{N(r)} + a_{p_1}(\rho^*)^{N(r)-p_1}z^{p_1} + \dots + a_{p_m}(\rho^*)^{N(r)-p_m}z^{p_m}|. \tag{18}$$

Now for  $|z| \neq 0$ ,

$$\begin{aligned} &|a_0(\rho^*)^{N(r)} + a_{p_1}(\rho^*)^{N(r)-p_1}z^{p_1} + \dots + a_{p_m}(\rho^*)^{N(r)-p_m}z^{p_m}| \\ &\leq |a_0|(\rho^*)^{N(r)} + |a_{p_1}|(\rho^*)^{N(r)-p_1}|z|^{p_1} + \dots + |a_{p_m}|(\rho^*)^{N(r)-p_m}|z|^{p_m} \\ &\leq |a_0|(\rho^*)^{N(r)}(1 + |z|^{p_1} + \dots + |z|^{p_m}) \\ &= |a_0|(\rho^*)^{N(r)}|z|^{p_m+1} \left( \frac{1}{|z|} + \dots + \frac{1}{|z|^{p_m+1-p_2}} + \frac{1}{|z|^{p_m+1-p_1}} + \frac{1}{|z|^{p_m+1}} \right). \end{aligned} \tag{19}$$

Using (18) and (19), we have for  $|z| \neq 0$

$$\begin{aligned}
 |R(z)| &\geq |a_{N(r)}| |z|^{N(r)} - |a_0| (\rho^*)^{N(r)} |z|^{p_m+1} \left( \frac{1}{|z|} + \dots + \frac{1}{|z|^{p_m+1-p_2}} + \frac{1}{|z|^{p_m+1-p_1}} + \frac{1}{|z|^{p_m+1}} \right) \\
 &> |a_{N(r)}| |z|^{N(r)} - |a_0| (\rho^*)^{N(r)} |z|^{p_m+1} \left( \frac{1}{|z|} + \dots + \frac{1}{|z|^{p_m+1-p_2}} + \frac{1}{|z|^{p_m+1-p_1}} + \frac{1}{|z|^{p_m+1}} + \dots \right) \\
 &= |a_{N(r)}| |z|^{N(r)} - |a_0| (\rho^*)^{N(r)} |z|^{p_m+1} \sum_{k=1}^{\infty} \frac{1}{|z|^k}.
 \end{aligned} \tag{20}$$

The geometric series  $\sum_{k=1}^{\infty} \frac{1}{|z|^k}$  is convergent for

$$\frac{1}{|z|} < 1$$

i.e., for  $|z| > 1$

and converges to

$$\frac{1}{|z|} \frac{1}{1 - \frac{1}{|z|}} = \frac{1}{|z| - 1}.$$

Therefore

$$\sum_{k=1}^{\infty} \frac{1}{|z|^k} = \frac{1}{|z| - 1} \text{ for } |z| > 1.$$

So on  $|z| > 1$ ,

$$|R(z)| > 0 \text{ if } |a_{N(r)}| |z|^{N(r)} - \frac{|a_0| (\rho^*)^{N(r)} |z|^{p_m+1}}{|z| - 1} \geq 0$$

$$\text{i.e., if } |a_{N(r)}| |z|^{N(r)} \geq \frac{|a_0| (\rho^*)^{N(r)} |z|^{p_m+1}}{|z| - 1}$$

$$\text{i.e., if } |a_{N(r)}| |z|^{N(r)+1} - |a_{N(r)}| |z|^{N(r)} \geq |a_0| (\rho^*)^{N(r)} |z|^{p_m+1}$$

$$\text{i.e., if } |z|^{p_m+1} (|a_{N(r)}| |z|^{N(r)-p_m} - |a_{N(r)}| |z|^{N(r)-p_m-1} - |a_0| (\rho^*)^{N(r)}) \geq 0.$$

Let us consider

$$g(t) \equiv |a_{N(r)}| t^{N(r)-p_m} - |a_{N(r)}| t^{N(r)-p_m-1} - |a_0| (\rho^*)^{N(r)} = 0.$$

Clearly,  $g(t) = 0$  has one positive root because the maximum number of changes in sign in  $g(t)$  is one and  $g(0) = -|a_0| \rho^{N(r)}$  is  $-ve$ ,  $g(\infty)$  is  $+ve$ . Let  $t_0$  be the positive root of  $g(t) = 0$  and  $t_0 > 1$ . Clearly for  $t > t_0$ ,  $g(t) \geq 0$ . If not, for some  $t_1 > t_0$ ,  $g(t_1) < 0$ . Then  $g(t_1) < 0$  and  $g(\infty) > 0$ . Therefore  $g(t) = 0$  must have another positive root in  $(t_1, \infty)$  which gives a contradiction.

Hence for  $t \geq t_0$ ,  $g(t) \geq 0$  and  $t_0 > 1$ . So  $|R(z)| > 0$  for  $|z| \geq t_0$ .

Thus  $R(z)$  does not vanish in  $|z| \geq t_0$ .

Hence all the zeros of  $R(z)$  lie in  $|z| < t_0$ .

Let  $z = z_0$  be any zero of  $P(z)$ . So  $P(z_0) = 0$ . Clearly  $z_0 \neq 0$  as  $a_0 \neq 0$ . Putting  $z = \rho^* z_0$  in  $R(z)$  we have

$$R(\rho^* z_0) = (\rho^*)^{N(r)} \cdot P(z_0) = (\rho^*)^{N(r)} \cdot 0 = 0.$$

Therefore  $R(\rho^* z_0) = 0$  and so  $z = \rho^* z_0$  is a zero of  $R(z)$  and consequently  $|\rho^* z_0| < t_0$  which implies  $|z_0| < \frac{t_0}{\rho^*}$ .

As  $z_0$  is an arbitrary zero of  $P(z)$ , all the zeros of  $P(z)$  lie in  $|z| < \frac{t_0}{\rho^*}$ . (21)

Again let us consider

$$F(z) = (\rho^*)^{N(r)} z^{N(r)} P\left(\frac{1}{\rho^* z}\right).$$

Now

$$\begin{aligned} F(z) &= (\rho^*)^{N(r)} z^{N(r)} \left\{ a_0 + a_{p_1} \frac{1}{(\rho^*)^{p_1} z^{p_1}} + \dots + a_{p_m} \frac{1}{(\rho^*)^{p_m} z^{p_m}} + a_{N(r)} \frac{1}{(\rho^*)^{N(r)} z^{N(r)}} \right\} \\ &= a_0 (\rho^*)^{N(r)} z^{N(r)} + a_{p_1} (\rho^*)^{N(r)-p_1} z^{N(r)-p_1} + \dots + a_{p_m} (\rho^*)^{N(r)-p_m} z^{N(r)-p_m} + a_{N(r)}. \end{aligned}$$

Also

$$\begin{aligned} &|a_{p_1} (\rho^*)^{N(r)-p_1} z^{N(r)-p_1} + \dots + a_{p_m} (\rho^*)^{N(r)-p_m} z^{N(r)-p_m} + a_{N(r)}| \\ &\leq |a_{p_1}| (\rho^*)^{N(r)-p_1} |z|^{N(r)-p_1} + \dots + |a_{p_m}| (\rho^*)^{N(r)-p_m} |z|^{N(r)-p_m} + |a_{N(r)}| \\ &\leq |a_{p_1}| (\rho^*)^{N(r)-p_1} (|z|^{N(r)-p_1} + |z|^{N(r)-p_2} + \dots + |z|^{N(r)-p_m} + 1). \end{aligned}$$

So for  $|z| \neq 0$ ,

$$\begin{aligned} |F(z)| &\geq |a_0| (\rho^*)^{N(r)} |z|^{N(r)} - |a_{p_1} (\rho^*)^{N(r)-p_1} z^{N(r)-p_1} + \dots + a_{p_m} (\rho^*)^{N(r)-p_m} z^{N(r)-p_m} + a_{N(r)}| \\ &\geq |a_0| (\rho^*)^{N(r)} |z|^{N(r)} - |a_{p_1}| (\rho^*)^{N(r)-p_1} (|z|^{N(r)-p_1} + |z|^{N(r)-p_2} + \dots + |z|^{N(r)-p_m} + 1) \\ &= |a_0| (\rho^*)^{N(r)} |z|^{N(r)} - |a_{p_1}| (\rho^*)^{N(r)-p_1} |z|^{N(r)-p_1+1} \left( \frac{1}{|z|} + \frac{1}{|z|^{p_2-p_1+1}} + \dots + \frac{1}{|z|^{N(r)-p_1+1}} \right) \end{aligned}$$

i.e., on  $|z| \neq 0$ ,

$$|F(z)| > |a_0| (\rho^*)^{N(r)} |z|^{N(r)} - |a_{p_1}| (\rho^*)^{N(r)-p_1} |z|^{N(r)-p_1+1} \left( \sum_{k=1}^{\infty} \frac{1}{|z|^k} \right).$$

The geometric series  $\sum_{k=1}^{\infty} \frac{1}{|z|^k}$  is convergent for

$$\frac{1}{|z|} < 1$$

i.e., for  $|z| > 1$

and converges to

$$\frac{1}{|z|} \frac{1}{1 - \frac{1}{|z|}} = \frac{1}{|z| - 1}.$$

Therefore

$$\sum_{k=1}^{\infty} \frac{1}{|z|^k} = \frac{1}{|z| - 1} \text{ for } |z| > 1.$$

Therefore for  $|z| > 1$

$$\begin{aligned} |F(z)| &> |a_0| (\rho^*)^{N(r)} |z|^{N(r)} - |a_{p_1}| (\rho^*)^{N(r)-p_1} |z|^{N(r)-p_1+1} \left( \frac{1}{|z|-1} \right) \\ &= (\rho^*)^{N(r)-p_1} \left( (\rho^*)^{p_1} |a_0| |z|^{N(r)} - |a_{p_1}| \frac{|z|^{N(r)-p_1+1}}{|z|-1} \right) \\ &= (\rho^*)^{N(r)-p_1} |z|^{N(r)-p_1+1} \left( |a_0| (\rho^*)^{p_1} |z|^{p_1-1} - \frac{|a_{p_1}|}{|z|-1} \right). \end{aligned}$$

For  $|z| > 1$ ,

$$|F(z)| > 0 \text{ if } |a_0|(\rho^*)^{p_1}|z|^{p_1-1} - \frac{|a_{p_1}|}{|z|^{-1}} \geq 0$$

$$\text{i.e., if } |a_0|(\rho^*)^{p_1}|z|^{p_1-1} \geq \frac{|a_{p_1}|}{|z|^{-1}}$$

$$\text{i.e., if } |a_0|(\rho^*)^{p_1}|z|^{p_1} - |a_0|(\rho^*)^{p_1}|z|^{p_1-1} - |a_{p_1}| \geq 0. \quad (22)$$

Therefore on  $|z| > 1$ ,  $|F(z)| > 0$  if (22) holds.

Let us consider

$$f(t) \equiv |a_0|(\rho^*)^{p_1}t^{p_1} - |a_0|(\rho^*)^{p_1}t^{p_1-1} - |a_{p_1}| = 0.$$

Clearly  $f(t) = 0$  has exactly one positive root and is greater than one. Let  $t'_0$  be the positive root of  $f(t) = 0$ .

Therefore  $t'_0 > 1$ . Obviously if  $t \geq t'_0$  then  $f(t) \geq 0$ . So for  $|F(z)| > 0$ ,  $|z| \geq t'_0$ .

Therefore  $F(z)$  does not vanish in  $|z| \geq t'_0$ .

Hence all the zeros of  $F(z)$  lie in  $|z| < t'_0$ .

Let  $z = z_0$  be any zero of  $P(z)$ . Therefore  $P(z_0) = 0$ . Clearly  $z_0 \neq 0$  as  $a_0 \neq 0$ .

Now putting  $z = \frac{1}{\rho^*z_0}$  in  $F(z)$  we obtain that

$$\begin{aligned} F\left(\frac{1}{\rho^*z_0}\right) &= (\rho^*)^{N(r)} \left(\frac{1}{\rho^*z_0}\right)^{N(r)} \cdot P(z_0) \\ &= \left(\frac{1}{z_0}\right)^{N(r)} \cdot P(z_0) \\ &= \left(\frac{1}{z_0}\right)^{N(r)} \cdot 0 \\ &= 0. \end{aligned}$$

Therefore  $z = \frac{1}{\rho^*z_0}$  is a zero of  $F(z)$ . Now

$$\left|\frac{1}{\rho^*z_0}\right| < t'_0$$

$$\text{i.e., } \left|\frac{1}{z_0}\right| < \rho^*t'_0$$

$$\text{i.e., } |z_0| > \frac{1}{\rho^*t'_0}.$$

$$\text{As } z_0 \text{ is an arbitrary zero of } P(z) \text{ therefore we obtain that all the zeros of } P(z) \text{ lie in } |z| > \frac{1}{\rho^*t'_0}. \quad (23)$$

Using (21) and (23) we get that all the zeros of  $P(z)$  lie in the ring shaped region

$$\frac{1}{\rho^*t'_0} < |z| < \frac{t_0}{\rho^*}$$

where  $t_0, t'_0$  are the unique positive roots of the equations  $g(t) = 0$  and  $f(t) = 0$  respectively whose form is given in the statement of Theorem 3.

This proves the theorem.

**Corollary: 2** In view of Theorem 3 we may state that all the zeros of the polynomial

$P(z) = a_0 + a_{p_1}z^{p_1} + \dots + a_{p_m}z^{p_m} + a_nz^n$  of degree  $n$  with  $1 \leq p_1 < p_2 < \dots < p_m \leq n - 1$ ,  $p_i$ 's are integers such that  $|a_0| \geq |a_{p_1}| \geq \dots \geq |a_n|$  lie in the ring shaped region  $\frac{1}{t'_0} < |z| < t_0$

where  $t_0, t'_0$  are the unique positive roots of the equations

$$g(t) \equiv |a_n|t^{n-p_m} - |a_n|t^{n-p_m-1} - |a_0| = 0$$

and

$$f(t) \equiv |a_0|t^{p_1} - |a_0|t^{p_1-1} - |a_{p_1}| = 0$$

respectively just substituting  $\rho^* = 1$ .

## REFERENCES

1. Chattopadhyay, A.; Das, S.; Jain, V. K. and Konwer, H.: Certain generalization of Eneström-Keakeya theorem, J. Indian Math. Soc., Vol. 72, No. 1 - 4(2005), pp.147 - 156.
2. Datta, B. and Govil, N. K.: On the location of the zeros of polynomial, J. Approximation Theory, Vol. 24(1978), pp. 78 - 82.
3. Datta, S.K. and Biswas, T.: On the definition of a meromorphic function of order zero, Int. Mat. Forum, Vol. 4, No. 37(2009), pp.1851 - 1861.
4. Das, S. and Datta, S. K.: On Cauchy's proper bound for zeros of a polynomial, International J. of Math. Sci. and Engg. Appls. (IJMSEA), Vol. 2, No. IV (2008), pp. 241 - 252.
5. Govil, N. K. and Rahaman, Q. I.: On the Eneström-Keakeya theorem, Tohoku Math. J. Vol. 20(1968), pp.126 - 136.
6. Joyal, A., Labelle, G. and Rahaman, Q. I.: On the location of zeros of polynomials, Canad. Math. Bull., Vol. 10(1967), pp.53 - 63.
7. Jain, V. K. : On the location of zeros of polynomials, Ann. Univ. Mariae Curie-Sklodowska, Lublin-Polonia Sect. A, Vol. 30(1976), pp. 43 - 48.
8. Jain, V. K.: On Cauchy's bound for zeros of a polynomial, Truck. J. Math. Vol. 30(2006), pp.95 - 100.
9. Marden, M. : Geometry of polynomials, Amer. Math-Soc. Providence, R.I., 1966.
10. Mohammad, Q. G.: Location of zeros of polynomials, Amer. Math. Monthly, Vol. 74(1967), pp. 290 - 292.
11. Sun, Y. J. and Hsieh, J. G.: A note on circular bound of polynomial zeros, IEEE Trans. Circuit Syst. Vol. 143(1996), pp. 476 - 478.
12. Valiron, G.: Lectures on the general theory of integral functions, Chelsea Publishing Company (1949).
13. Zilovic, M. S.; Roytman, L. M.; Combettes, P. L. and Swami, M. N. S.: A bound for the zeros of polynomials, *ibid* Vol. 39(1992), pp. 476 - 478.

**Source of support: Nil, Conflict of interest: None Declared**