

**BEST ONE-SIDED APPROXIMATION OF UNBOUNDED FUNCTION**

**Saheb AL-Saidy\* & Jawad KadhimJoodi**

*Dept. Of Mathematics, College of Science, Al-Mustansiriya University.*

*(Received on: 08-03-13; Revised & Accepted on: 22-10-13)*

**ABSTRACT**

In this work, we shall study characterize of one-sided approximation of unbounded functions in weighted space and we shall compare between different norms such as  $\|f\|_{p,\alpha}$ ,  $\|f\|_{\delta,p,\alpha}$  and  $\|f\|_{\delta,p,\alpha}^\Phi$  where  $f$  is any unbounded function by using the function  $\Phi_v = \Phi_v(x)$  such that:

$$\Phi_v(x) = \sin^4\left(\frac{\pi}{4m}\right) \left\{ (\sin^4 m(x - x_v)) / (\sin^4(x - x_v)/2) + (\sin^4 m(x + x_v)) / (\sin^4(x + x_v)/2) \right\}$$

where  $m$  be natural number,  $v = \{0, 1, 2, \dots, m - 1\}$ ,  $x \in X = [0, 1]$  or  $x \in X = [-1, 1]^d$  ( $x$  is multi-variable) and  $x_v = \pi - (2v + 1)\pi/2m$ . Also we shall deduce degree of best one sided approximation in weighted by using some result of degree of best approximation in  $L_p$ -space, also we discuss the relation between degree of best one-sided approximation of unbounded function and degree of best one sided approximation of it's derivative.

**1. INTRODUCTION**

Throughout this paper, we use the weight function  $\omega(x) = e^{-\alpha x}$  which is non-negative measurable function on  $(0, \infty)$ . For  $1 \leq p \leq \infty$  the weighted space:

$L_{p,\alpha} = \{f: X \rightarrow R, \text{ such that } |f(x)\omega_\alpha(x)| \leq M, \alpha \geq 1\}$ , such that for the function  $f$ ,

$$\|f\|_{p,\alpha} = \left[ \int_a^b |f(x)|^p e^{-\alpha p x} dx \right]^{\frac{1}{p}} < \infty$$

Also,  $\|f\|_{\infty,\alpha} = \left\{ \sup\{|f(x)e^{-\alpha x}| : x \in X\} \right\} < \infty$ . The local norm of the function  $f$  is define by:

$$\|f\|_{\delta,p,\alpha} = \left[ \int_X \left[ \sup\{|f(x)e^{-\alpha x}| : x \in N(\delta, t)\} \right]^p dx \right]^{\frac{1}{p}}$$

where  $\delta > 0$  and  $N(\delta, t) = \{y \in X : |t - y| \leq \delta\}$ . By using the function:

$$\psi(u, \delta) = \delta\Phi(u) + \delta^2$$

where  $\Phi(u) = \begin{cases} (1 - u^2)^{\frac{1}{2}} & \text{if } X = [-1, 1] \\ u(1 - u)^{\frac{1}{2}} & \text{if } X = [0, 1] \end{cases}$

We can define another local norm as:  $\|f\|_{\delta,p,\alpha}^\Phi = \left[ \int_X \left[ \sup\{|f(x)e^{-\alpha x}| : x \in N(\psi, \delta, x)\} \right]^p dx \right]^{\frac{1}{p}}$  also we mention

$\omega_r^\Phi(f, x, \delta)_{p,\alpha} = \sup_{0 \leq h \leq \delta} \|\Delta_{h\varphi}^r f(x)\|_{p,\alpha}$  (the ordinary Ditzian-Totik modulus of smoothness for  $f$ )

where:  $\Delta_{h\varphi}^r f(x) = \sum_{i=0}^r (-1)^{i+r} C_i^r f(x + i\varphi h)$ ,  $x + i\varphi h \in [0, 1]$

And we shall define the algebraic polynomial  $F_m$  such that:

$$F_m(x) = \Phi_m(x) = \Phi_v(\arccos x) \text{ on } x \in [0, 1] \text{ and } \Phi_{j,m}(x) = \prod_{s=1}^d F_{j_s,m}(x_s)$$

where  $x \in [-1, 1]^d$ . Suppose  $P_n$  is the polynomial of best approximation of the function  $f$  in the weighted space which is a Banach Space. And suppose that:

**Corresponding author: Saheb AL-Saidy\*. \*E-mail: ali12mm@yahoo.com**

$$Q_n^\mp(f, x) = P_n(f, x) \mp \sum_{j \in Z} \Phi_{j,m}(x) \|f - P_n(f, x)\|_{\infty, \alpha(X_j)}$$

where  $j = (j_1, j_2, \dots, j_d)$ ,  $j \in Z$  such that  $Z = \{0, 1, 2, \dots, m-1\}^d$ . And  $X_j = [z_{j_1}, z_{j_1+1}] \times [z_{j_1+1}, z_{j_1+2}] \times \dots \times [z_{j_d}, z_{j_d+1}]$ .

Also we define  $\hat{Z} = \{0, 1, 2, \dots, m\}^d$  and  $\hat{X}_j = [z_{j_1-1}, z_{j_1+1}] \times [z_{j_1+1}, z_{j_1+2}] \times \dots \times [z_{j_d-1}, z_{j_d+1}]$ . Clear (as we shall prove in theorem (3.4)) that:

$Q_n^-(f, x) \leq f(x) \leq Q_n^+(f, x)$ . We define the degree of best one-sided approximation of function  $f$  as:

$$E_n^- = \inf \|P_n^+ + P_n^-\| \quad \text{where } P_n^\mp \in P_n \text{ (where } P_n \text{ is the space of all polynomials of degree } n).$$

## 2. AUXILIARY RESULTS

Here we brief some results which we needed in our work:

**Lemmas 2.1:** [1] Let  $j \in Z$  then:

- (i)  $\Phi_{j,m}(x) \in P_{4m-2}^d$ .
- (ii)  $\Phi_{j,m}(x) \geq 1 \quad \forall x \in X_j$ .

**Lemma 2.2:** Let  $a_j \geq 0, j \in Z$  then:

$$\left\| \sum_{j \in Z} a_j \Phi_{j,m}(x) \right\|_{p, \alpha(X)} \leq c \sum_{j \in Z} a_j^p (z_j - z_{j-1})^{\frac{1}{p}}$$

**Proof:** By using lemma 2.1(ii) we get:

$$\sum_{j \in Z} a_j \Phi_{j,m}(x) e^{-\alpha p x} \leq \sum_{j \in Z} a_j \Phi_{j,m}(x)$$

$$\begin{aligned} \text{Also, } \left\| \sum_{j \in Z} a_j \Phi_{j,m}(x) \right\|_{p, \alpha(X)} &= \left\| \sum_{j \in Z} a_j \Phi_{j,m}(x) e^{-\alpha p x} \right\|_{p(X)} \\ &\leq \left\| \sum_{j \in Z} a_j \Phi_{j,m}(x) \right\|_{p(X)} \\ &\leq c \left[ \sum_{j \in Z} a_j^p (z_j - z_{j-1}) \right]^{\frac{1}{p}}. \end{aligned}$$

**Lemma 2.3:** [1] Suppose that  $u, v \in [-1, 1]$  and  $0 < t \leq \frac{1}{2}$  then:

$$|u - v| \leq \psi(t, u) \text{ where } \psi(t, u) \leq 6\psi(t, u) \text{ and } \psi(t, v) \leq 4\psi(t, u).$$

**Lemma 2.4:** Let  $f \in L_{p, \alpha}(X)$ ; ( $1 \leq p \leq \infty$ ) and  $0 < t \leq \frac{1}{2}$  then:

$$\left\| \psi(t, \cdot)^{\frac{1}{p}} \|f\|_{p, \alpha(N(t, \cdot))} \right\|_{p, \alpha} \leq c \|f\|_{p, \alpha(X)}$$

**Proof:**

$$\begin{aligned} \left\| \psi(t, \cdot)^{\frac{1}{p}} \|f\|_{p, \alpha(N(t, \cdot))} \right\|_{p, \alpha} &= \left\| \psi(t, \cdot)^{\frac{1}{p}} \right\|_{p, \alpha(X)} \left\| \|f\|_{p, \alpha(N(t, \cdot))} \right\|_{p, \alpha(X)} \\ &\leq \|\psi(t, \cdot)^p\|_{p, \alpha(X)} \|f\|_{p, \alpha(X)} \leq c_p \|f\|_{p, \alpha(X)}. \end{aligned}$$

**Lemma 2.5:** [2] If  $f$  is bounded measurable function on  $[a, b]$ ,  $a, b \in R$ , then:

$$\int_a^b f(x) dx \approx (b - a) n^{-1} \sum_{i=1}^n f(x_i) \text{ where } x_i = a + \frac{1}{2n}(b-a)(2i-1).$$

**Lemma 2.6:** [3] Let  $m$  and  $\delta$  be any number such that  $m\delta \leq \frac{1}{4}$  then for  $f \in L_{p, \alpha}$  we get:

$$\|f\|_{m\delta, p, \alpha}^\Phi \leq c_p m^{\frac{2d}{p}} \|f\|_{\delta, p, \alpha}^\Phi.$$

**Lemma 2.7:** [1] Suppose that  $x \in X = [-1,1]^d, 0 \leq t \leq \frac{1}{2}$  and  $N = [2\pi/t]$  then:

$$\dot{X}_j \subset N(t, x), x \in X_j$$

We can write the local norm  $\|f\|_{\delta,p,\alpha}^\Phi$  of the unbounded functions f with form globally norm  $\|f_\delta\|_{p,\alpha(X)}$  as the following:

$$\|f\|_{\delta,p,\alpha}^\Phi = \|f\|_{\delta,p,\alpha(X)}^\Phi = \|\|f\|_{\infty,N(\delta,x)}\|_{p,\alpha(X)} = \|f_\delta\|_{p,\alpha(X)}$$

where  $f_\delta(X) = \sup\{|f(t)|: t \in N(\delta, x)\}$ .

Also, we can define  $N(\delta, x)^\sigma$  as the following:

$$\begin{aligned} N(\delta, x)^\sigma &= N(\psi(\delta, x), x)^\sigma = \prod_{s:\sigma_s=1} N(\psi(\delta, x_s), x_s) \\ &= N(\psi(\delta, x_1), x_1) \dots \dots N(\psi(\delta, x_d), x_d) \end{aligned}$$

$$\text{Also, } \psi(\delta, x)^\sigma = \prod_s \psi(\delta, x_s) = \prod_{s=1}^d \psi(\delta, x_s).$$

### 3. MAIN RESULTS

Now, we prove the following theorem where  $x$  is a single variable that is  $x \in [0,1]$

**Theorem 3.1:** Suppose that  $P_n \in P_n$  then:

$$\|P_n\|_{\delta,p,\alpha}^\Phi \leq c_p [1 + \max(n\delta, n^2\delta^2)]^{\frac{1}{p}} \|P_n\|_{p,\alpha} \text{ Where } 1 \leq p \leq \infty$$

**Proof:** We shall use the following equality:

$$P_n(t) - P_n(x) = \int_x^t \dot{P}_n(u) du, \quad x, t \in X$$

$$\begin{aligned} \text{So, } |\|P_n\|_{\delta,p,\alpha}^\Phi - \|P_n\|_{p,\alpha}| &\leq [\int_X (\int_{N(x,\delta)} (\dot{P}_n(u) du)^p \omega_\alpha^p(x) dx)]^{\frac{1}{p}} \\ &\leq [c_1 \int_X \frac{1}{[N(x,\delta)]^p} \int_{[N(x,\delta)]^p} (\delta\Phi(u) + \delta^2)^p (\dot{P}_n(u) du)^p \omega_\alpha^p(x) dx]^{\frac{1}{p}} \\ &\leq c_1^{\frac{1}{p}} [(\frac{\delta\Phi(u)+\delta^2}{N(x,\delta)})^p]^{\frac{1}{p}} [\int_X [\dot{P}_n(x)]^p \omega_\alpha^p(x) dx]^{\frac{1}{p}} \\ &\leq c_p \|(\delta\Phi + \delta^2)\dot{P}_n(x)\|_{p,\alpha} \\ &\leq c_p [\delta\|\Phi\dot{P}_n\|_{p,\alpha} + \delta^2\|\dot{P}_n\|_{p,\alpha}] \\ &\leq [\max(n\delta, n^2\delta^2)] \|P_n\|_{p,\alpha} \\ &= [1 + \max(n\delta, n^2\delta^2)] \|P_n\|_{p,\alpha} \end{aligned}$$

Then:  $\|P_n\|_{\delta,p,\alpha}^\Phi \leq c_p [1 + \max(n\delta, n^2\delta^2)]^{\frac{1}{p}} \|P_n\|_{p,\alpha}$  where  $1 \leq p \leq \infty$ .

Now, we shall prove same theorem where  $x$  is multivariable that is  $x \in [-1,1]^d$

**Theorem 3.2:** Suppose that  $P_n \in P_n^d$  then:

$$\|P_n\|_{\delta,p,\alpha}^\Phi \leq c_d [1 + \max(n\delta, n^2\delta^2)]^{\frac{1}{p}} \|P_n\|_{p,\alpha} \text{ Where } 1 \leq p \leq \infty$$

**Proof:** We shall us the following equality:

$$P(t) - P(x) = \sum_{\substack{\alpha, |\alpha| \geq 1 \\ \alpha_s = 0,1}} \int_{x^{(\alpha)}}^{t^{(\alpha)}} D^\alpha P_n(x^{(1-\alpha)} + u^{(\alpha)}) du^{(\alpha)}$$

By using lemmas (2.3) and (2.4) we get:

$$\begin{aligned}
 \|P_n\|_{\delta,p,\alpha}^\Phi - \|P_n\|_{p,\alpha} &\leq \left[ \int_X |P_n(\xi_x) - P_n(x)|^p \omega_\alpha^p(x) dx \right]^{\frac{1}{p}} \\
 &\leq \left[ \int_X \left[ \sum_{|\alpha| \geq 1} \int_{N(x,\delta)^\alpha} D^\alpha P_n(x^{(1-\alpha)} + u^{(\alpha)}) du^{(\alpha)} \right]^p \omega_\alpha^p(x) dx \right]^{\frac{1}{p}} \\
 &\leq \left[ c_1 \sum_{|\alpha| \geq 1} \int_X \frac{1}{[N(x,\delta)^\alpha]^p} \int_{N(x,\delta)^\alpha} ([\delta\Phi(u)^\alpha + \delta^2]^p [D^\alpha P_n(x^{(1-\alpha)} + u^{(\alpha)}) du^{(\alpha)}]^p \omega_\alpha^p(x) dx \right]^{\frac{1}{p}} \\
 &\leq c_1^{\frac{1}{p}} \left[ \left( \frac{\delta\Phi(u)^\alpha + \delta^2}{N(x,\delta)} \right)^{p\alpha} \right]^{\frac{1}{p}} \left[ \int_X (D^\alpha P_n(x^{(1-\alpha)} + u^{(\alpha)}) du^{(\alpha)})^p \omega_\alpha^p(x) dx \right]^{\frac{1}{p}} \\
 &\leq c_p \sum_{|\alpha| \geq 1} \left[ \int_X |D^\alpha P_n(x)|^p [\delta\Phi(x) + \delta^2]^\alpha \omega_\alpha^p(x) dx \right]^{\frac{1}{p}} \\
 &\leq c_p \sum_{|\alpha| \geq 1} \|[\delta\Phi(x) + \delta^2]^\alpha D^\alpha p_n\|_{p,\alpha} \\
 &\leq c_p \sum_{|\alpha| \geq 1} [\delta^{|\alpha|} \|\Phi^\alpha D^\alpha p_n\|_{p,\alpha} + \delta^{2|\alpha|} \|D^\alpha p_n\|_{p,\alpha}] \\
 &\leq c_p \sum_{|\alpha| \geq 1} [\max(n\delta, n^2\delta^2)]^{|\alpha|} \|P_n\|_{p,\alpha} \\
 &\leq c_p [(1 + \max(n\delta, n^2\delta^2))^d - 1] \|P_n\|_{p,\alpha}
 \end{aligned}$$

So,  $\|P_n\|_{\delta,p,\alpha}^\Phi \leq c_p [1 + \max(n\delta, n^2\delta^2)]^d \|P_n\|_{p,\alpha}$  where  $1 \leq p \leq \infty$ .

By using theorem 3.2 we get the following corollary:

**Corollary 3.3:** Suppose that  $P_n \in \mathbf{P}_n^d$  then:

$$\|P_n\|_{p,\alpha(X)} \leq \|P_n\|_{p,\alpha(X)}^\Phi \leq c_d \|P_n\|_{p,\alpha(X)}$$

**Theorem 3.4:** For any function  $f$  on  $X = [-1,1]^d$

$$Q_n^-(f, x) \leq f(x) \leq Q_n^+(f, x)$$

**Proof:** Let  $x \in X = [-1,1]^d$  and by using lemma (2.1) we get:

$$Q_n^+(f, x) = P_n(x) + \sum_{j \in \mathbb{Z}} \Phi_{j,m}(x) \|f(x) - P_n(x)\|_{\infty(X_j)}$$

$$\geq P_n(x) + \|f(x) - P_n(x)\|_{\infty(X_j)}$$

$$\geq P_n(x) + |f(x) - P_n(x)| = f(x).$$

$$Q_n^-(f, x) = P_n(x) - \sum_{j \in \mathbb{Z}} \Phi_{j,m}(x) \|f(x) - P_n(x)\|_{\infty(X_j)}$$

$$\leq P_n(x) - \|f - P_n(x)\|_{\infty(X_j)}$$

$$\leq P_n(x) - |f(x) - P_n(x)| = f(x)$$

So,  $Q_n^-(f, x) \leq f(x) \leq Q_n^+(f, x)$ .

**Theorem 3.5:** Suppose that  $f \in L_{\infty,\alpha}(X)$  then:

$$E_n^{\sim}(f)_{p,\alpha} \leq c_d E_n(f)_{1/n,\alpha,p}^\Phi \leq c_d E_n^{\sim}(f)_{p,\alpha}$$

**Proof:** Suppose that  $P_n^+, P_n^- \in \mathbf{P}_n^d$  such that  $p_n^- \leq f(x) \leq p_n^+, x \in X$

$$\text{And } E_n^{\sim}(f)_{p,\alpha} = \|P_n^+ - P_n^-\|_{p,\alpha}$$

So by using corollary (3.3)

$$E_n(f)_{1/n,\alpha,p}^\Phi \leq E_n^\sim(f)_{1/n,\alpha,p}^\Phi \leq \|P_n^+ - P_n^-\|_{1/n,p,\alpha}^\Phi \leq c_p \|P_n^+ - P_n^-\|_{p,\alpha}$$

$$= c_p E_n^\sim(f)_{p,\alpha}$$

Then:  $E_n(f)_{1/n,\alpha,p}^\Phi \leq c_d E_n^\sim(f)_{p,\alpha}$

Also by using lemmas (2.2), (2.6), (2.7) and theorem (3.5), we get:

$$E_n^\sim(f)_{p,\alpha} \leq \|Q_n^+ - Q_n^-\|_{p,\alpha} = 2 \left\| \sum_{j \in Z} \Phi_{j,m}(x) \|f - P_n\|_{\infty(X_j)} \right\|_{p,\alpha}$$

$$\leq c \left[ \sum_{j \in Z} (Z_j - Z_{j-1}) \|f - P_n\|_{\infty(X_j)}^p \right]^{\frac{1}{p}}$$

$$\leq c \left[ \sum_{j \in Z} \int_{X_j} \|f - P_n\|_{\infty(X_j)}^p dx \right]^{\frac{1}{p}}$$

$$\leq c \left[ \sum_{j \in Z} \int_{X_j} \|f - p_n\|_{\infty(N(2\pi/m-1,x))}^p dx \right]^{\frac{1}{p}}$$

$$\leq c \left[ \sum_{j \in Z} \int_X \|f - p_n\|_{\infty(N(2\pi/m-1,x))}^p dx \right]^{\frac{1}{p}}$$

$$\leq c_p \|f - p_n\|_{1/n,p,\alpha}^\Phi$$

$$= c_p E_n(f)_{1/n,p,\alpha}^\Phi$$

So,  $E_n^\sim(f)_{p,\alpha} \leq c_p E_n(f)_{1/n,\alpha,p}^\Phi \leq c_p E_n^\sim(f)_{p,\alpha}$ .

**Theorem 3.6:** Suppose that  $f \in L_{p,\alpha}(X)$  such that  $1 \leq p \leq \infty$  and let

$$\sum_{v=1}^\infty V^{2d/p-1} E_v(f)_{p,\alpha} < \infty, \text{ and let } f = F \text{ almost}$$

everywhere then:  $E_n^\sim(f)_{p,\alpha} \leq c_p n^{-2d/p} \sum_{v=1}^\infty V^{2d/p-1} E_v(f)_{p,\alpha}$ .

**Proof:** Since  $E_v(f)_{p,\alpha} = \|f - Q_v\|_{p,\alpha(X)}$ ,  $v = 0, 1, 2 \dots \dots$

Since for  $n \in N$ ,  $\sum_{v=1}^m (Q_{n.2} - Q_{n.2^{v-1}}) = Q_{n.2^m} - Q_n$

And since  $F = f$  almost everywhere

So,  $\|F - Q_{n.2^{v-1}}\|_{\infty,\alpha(X)} \rightarrow 0$  (as  $N \rightarrow \infty$ )

And,  $F - Q_n(x) = \sum_{v=1}^\infty (Q_{n.2} - Q_{n.2^{v-1}})$

Then by using theorems (3.2) and (3.5) we get:

$$E_v^\sim(f)_{p,\alpha} \leq c_p E_n(f)_{1/n,\alpha,p}^\Phi = c_p E_n(F - Q_n)_{1/n,\alpha,p}^\Phi$$

$$\leq c_p \|F - Q_n\|_{1/n,p,\alpha}^\Phi$$

$$\leq c_p \sum_{v=1}^\infty \|Q_{n.2^v} - Q_{n.2^{v-1}}\|_{1/n,p,\alpha}^\Phi$$

$$\leq c_p \sum_{v=1}^\infty c_p (2^v)^{\frac{2d}{p}} \|Q_{n.2^v} - Q_{n.2^{v-1}}\|_{p,\alpha}$$

$$\leq c_p \sum_{v=1}^\infty 2^{2vd/p} [E_{n.2^v}(f)_{p,\alpha} + E_{n.2^{v-1}}(f)_{p,\alpha}]$$

$$\leq c_p n^{-2d/p} \sum_{v=1}^{\infty} (2^v \cdot n)^{2d/p} [E_{n \cdot 2^v}(f)_{p,\alpha} + E_{n \cdot 2^{v-1}}(f)_{p,\alpha}]$$

$$\leq c_p n^{-2d/p} \sum_{v=n}^{\infty} v^{2d/p-1} E_n(f)_{p,\alpha} .$$

Before we prove direct theorem of best one-sided approximation in weight space we shall refer to same theorem of best approximation in weight space by using  $B_n(f, x)$ -operator, where  $x$  is single variable.

**Theorem 3.7:** [4] Let  $f \in L_{p,\alpha}(X)$ (single case) then  $\|f(x) - B_n(f, x)\|_{p,\alpha} \leq c\omega_2^\varphi(f, x, \delta)_{p,\alpha} + \tau_2^\varphi(f, \delta)_{p,\alpha}$  where  $B_n(f, x)$  is Bernstein Polynomial of .

**Theorem 3.8: (Direct Theorem of onesided approximation in single case)** Let  $f$  be any function on  $[0, 1]$  and let:  $Q_n^\mp(f, x) = P_n + \Phi_m(x)\|f - P_n\|_{\infty(X)}$  be an operator (where  $p_n$  is a best approximation of  $f$ ) then

$$E_n^\sim(f)_{p,\alpha} \leq c\omega_2^\varphi(f, x, \delta)_{p,\alpha} + \tau_2^\varphi(f, \delta)_{p,\alpha} .$$

**Proof:** By using lemma (2.5) and (3.5) and (3.7) we get:

$$E_n^\sim(f)_{p,\alpha} \leq c_p E_n(f)_{1/n,p,\alpha}^\Phi \leq c_p E_n^\sim(f)_{p,\alpha}$$

$$\text{Then : } E_n^\sim(f)_{p,\alpha} \leq c_p E_n(f)_{1/n,p,\alpha}^\Phi$$

Since  $\int_a^b f(x)dx \approx \frac{(b-a)}{n} \sum_{i=1}^n f(x_i)$  where  $x_i = a + \frac{1}{2n}(b-a)(2i-1)$ .

$$\text{So, we have } \left(\frac{b-a}{n} \sum_{i=1}^n (f^p(x_i)\omega_\alpha^p(x_i))\right)^{\frac{1}{p}} \approx \left(\int_a^b (f^p(x)\omega_\alpha^p(x)dx)\right)^{\frac{1}{p}}$$

$$\text{That is } \left(\frac{b-a}{n} \sum_{i=1}^n (f^p(x_i)\omega_\alpha^p(x_i))\right)^{\frac{1}{p}} \leq c_p \left(\int_a^b (f^p(x)\omega_\alpha^p(x)dx)\right)^{\frac{1}{p}} \text{ Then, } \|f\|_{1/n,p,\alpha} \leq c_p \|f\|_{p,\alpha}$$

$$\text{Thus, } E_n^\sim(f)_{p,\alpha} \leq c_p E_n(f)_{1/n,p,\alpha}^\Phi \leq c_p E_n(f)_{1/n,p,\alpha} \leq c_p c_p E_n(f)_{p,\alpha}$$

$$\text{So, } E_n^\sim(f)_{p,\alpha} \leq c_p c_p c\omega_2^\varphi(f, x, \delta)_{p,\alpha} + \tau_2^\varphi(f, \delta)_{p,\alpha} = c_p c\omega_2^\varphi(f, x, \delta)_{p,\alpha} + \tau_2^\varphi(f, \delta)_{p,\alpha} .$$

Before, we prove inverse theorem of best one-sided approximation in weight space we shall refer to same theorem of best approximation in weight space by using  $B_n(f, x)$ -operator, where  $x$  is single variable.

**Theorem 3.9:** [4] Let  $f \in L_{p,\alpha}(X)$ (single case) such that  $(1 \leq p < \infty)$  then:

$$\tau_2^\varphi(f, \Delta, n^{-1})_{p,\alpha} \leq \frac{1}{n} \sum_{k=0}^n \|f - B_k(f)\|_{p,\alpha} .$$

**Theorem 3.10: (Inverse Theorem of one-sided approximation in single case)** Let  $f$  be any function on  $[0,1]$  and let:  $Q_n^\mp(f, x) = P_n + \Phi_m(x)\|f - P_n\|_{\infty(X)}$  be an operator where  $P_n$  is a best approximation of  $f$

$$\text{then: } \tau\left(f, \frac{1}{n}\right)_{p,\alpha} \leq \frac{c}{n} \sum_{s=0}^n \begin{cases} \|f - L_n\|_{p,\alpha} + \|f^\sim - L_n\|_{p,\alpha} & \text{if } p = 1, \infty \\ \|f - L_n\|_{p,\alpha} & \text{if } 1 < p < \infty \end{cases}$$

**Proof:** By using theorems (3.5) and (3.9) we get:

$$E_n^\sim(f)_{p,\alpha} \leq c_p E_n(f)_{1/n,\alpha,p}^\Phi \leq c_p E_n^\sim(f)_{p,\alpha}$$

$$\text{That is } E_n^\sim(f)_{p,\alpha} \leq c_p E_n(f)_{1/n,\alpha,p}^\Phi$$

$$\text{But we have } E_n(f)_{p,\alpha} \leq E_n(f)_{1/n,p,\alpha}^\Phi \leq c_p E_n^\sim(f)_{p,\alpha}$$

$$\text{Then: } E_n(f)_{p,\alpha} \leq c_p E_n^\sim(f)_{p,\alpha}$$

$$\text{So, } \sum_{i=1}^N E_n(f)_{p,\alpha} \leq c_p \sum_{i=1}^N E_n^\sim(f)_{p,\alpha}$$

$$\text{Thus: } \tau_2^\varphi(f, \Delta, n^{-1})_{p,\alpha} \leq \frac{1}{n} \sum_{k=0}^n \|f - B_k(f)\|_{p,\alpha} .$$

Now we discuss the relation between degree of best one-sided approximation of unbounded function  $f$  and degree of best one-sided approximation of its derivative, where  $x$  is single variable.

**Theorem 3.12:** Let  $f \in L^{(1)}_{p,\alpha}$  be any function on  $[0, 1]$  (i.e.  $f$  and  $\dot{f}$  in  $L_{p,\alpha}$ ) and let:

$Q_n^\mp(f, x) = P_n + \Phi_m(x) \|f - P_n\|_{\infty(X)}$  be an operator where  $p_n$  is a best approximation of  $f$  then

$$E_n^{\sim}(\dot{f})_{p,\alpha} \leq c_p(4m - 2)\tau(f, \frac{1}{n})_{p,\alpha}.$$

**Proof:** Since  $Q_n^-(f, x) \leq f(x) \leq Q_n^+(f, x), x \in [0,1]$  Then one of the cases is true:

$$\begin{aligned} \dot{Q}_n^-(f, x) \leq \dot{f}(x) \leq \dot{Q}_n^+(f, x), \dot{Q}_n^+(f, x) \leq \dot{f}(x) \leq \dot{Q}_n^-(f, x), \dot{Q}_n^-(f, x) \leq \dot{Q}_n^+(f, x) \leq \dot{f}(x), \dot{f}(x) \leq \dot{Q}_n^-(f, x) \\ \leq \dot{Q}_n^+(f, x), \dot{Q}_n^+(f, x) \leq \dot{Q}_n^-(f, x) \leq \dot{f}(x) \text{ or } \dot{f}(x) \leq \dot{Q}_n^+(f, x) \leq \dot{Q}_n^-(f, x). \end{aligned}$$

Where  $\dot{Q}_n^\mp(f, x) = \dot{P}_n \mp \dot{\Phi}_m(x) \|f(x) - P_n\|_{\infty(X)}$

If  $\dot{Q}_n^-(f, x) \leq \dot{f}(x) \leq \dot{Q}_n^+(f, x), \dot{Q}_n^+(f, x) \leq \dot{f}(x) \leq \dot{Q}_n^-(f, x)$

Then, by similar way of proof of theorem 3.5 and using lemma 2.1(i) and Bernstein inequality we get:

$$\begin{aligned} E_n^{\sim}(\dot{f})_{p,\alpha} &\leq \|\dot{Q}_n^+(f, x) - \dot{Q}_n^-(f, x)\|_{p,\alpha} \\ &= \|\dot{P}_n - \dot{P}_n + 2(\dot{\Phi}_m(x) \|f(x) - P_n\|_{\infty(X)})\|_{p,\alpha} \\ &= 2\|\dot{\Phi}_m(x) \|f(x) - P_n\|_{\infty(X)}\|_{p,\alpha} \\ &\leq 2(4m - 2)\|\Phi_m(x) \|f(x) - P_n\|_{\infty(X)}\|_{p,\alpha} \\ &\leq c_1(4m - 2) \left[ (1) \|f(x) - P_n(x)\|_{\infty(X)}^p \right]^{\frac{1}{p}} \\ &\leq c_1(4m - 2) \left[ \int_X (\|f(x) - P_n(x)\|_{\infty(X)}^p dx) \right]^{\frac{1}{p}} \\ &\leq c_1(4m - 2) \left[ \int_X (\|f(x) - P_n(x)\|_{\infty(N(2\pi/m-1,x))}^p dx) \right]^{\frac{1}{p}} \\ &\leq c_1(4m - 2) \|f - P_n(x)\|_{1/n,p,\alpha}^\Phi \\ &= c_1(4m - 2) E_n(f)_{1/n,p,\alpha} \\ &\leq c_1 c_p (4m - 2) E_n(f)_{p,\alpha} \\ &\leq c c_1 c_p (4m - 2) \tau(f, \frac{1}{n})_{p,\alpha} = c_p (4m - 2) \tau(f, \frac{1}{n})_{p,\alpha}. \end{aligned}$$

If  $\dot{Q}_n^-(f, x) \leq \dot{Q}_n^+(f, x) \leq \dot{f}(x), \dot{f}(x) \leq \dot{Q}_n^-(f, x) \leq \dot{Q}_n^+(f, x), \dot{Q}_n^+(f, x) \leq \dot{Q}_n^-(f, x) \leq \dot{f}(x)$   
or  
 $\dot{f}(x) \leq \dot{Q}_n^+(f, x) \leq \dot{Q}_n^-(f, x).$

Then:

$$E_n^{\sim}(\dot{f})_{p,\alpha} = E_n(\dot{f})_{p,\alpha} \leq \|\dot{f} - \dot{Q}_n^\mp(f, x)\|_{p,\alpha} \leq c_p \|f - Q_n^\mp(f, x)\|_{p,\alpha} \leq c_p \omega(f, \Delta_n(x)).$$

where  $\Delta_n(x) = \max\left(\frac{\sqrt{1-x^2}}{n}, \frac{1}{n^2}\right) n = 1, 2, \dots$  and  $\Delta_0(x) = 1.$

## REFERENCES

- [1] Saheb K.AL-Saidy;"one sided approximation of functions", Bulgaria, 1991.
- [2] Vasil A. Popov;" The average moduli of smoothness", Wiley Publication, England; 1988.

[3] Saheb K.AL-Saidy and Ahmed AL-Asady; "Equi-Approximation of functions in weighted-Space" ALMustansiriya University 2010( $1 \leq p < \infty$ ).

[4] Lekaa A. Husain "Unbounded Functions Approximation In Some  $L_{p,\alpha}$ -Spaces "Iraqi, College of science/AL-Mustansiriya University,B.SC.2010.

**Source of support: Nil, Conflict of interest: None Declared**