



ON SYLOW NUMBERS OF THE ALTERNATING SIMPLE GROUPS¹

Rulin Shen*

Department of Mathematics, Hubei University for Nationalities, Enshi, Hubei, P.R. China, 445000

E-mail: rulinshen@gmail.com

Ronghua Tan

Department of Mathematics, Hubei University for Nationalities, Enshi, Hubei, P.R. China, 445000

(Received on: 29-04-11; Accepted on: 11-05-11)

ABSTRACT

In this paper we give the formula of numbers of Sylow subgroups of alternating simple groups. This partially solves a problem posed by Jiping Zhang.

Keywords: symmetric groups, alternating groups, Sylow numbers.

MR (2000): 20D60, 20D06.

1. INTRODUCTION AND LEMMAS:

The structure of normalizer of Sylow 2-subgroup of symmetric groups S_n is studied by P. Hall (see Lemma 4 of [1], or [2]), he proved that Sylow 2-subgroup is self-normalized. The study in the numbers of Sylow subgroups (also called the Sylow numbers) is due to Professor Jiping Zhang [5]. He proved a conjecture of Huppert's. Then he put forward a problem in the end of his paper [5]. determine the number of Sylow p -subgroup of finite simple groups. In this paper we give the formula of Sylow numbers of the alternating simple group A_n , which partially solves above Zhang's

problem. Let $\Omega = \{1, 2, 3, \dots, n\}$, $S(\Omega)$ or S_n be symmetric group on Ω and P_n be a Sylow p -subgroup of

symmetric group S_n . Then $|P_n| = (n!)_p = p^{s(n)}$, where $s(n) = \left[\frac{n}{p} \right] + \left[\frac{n}{p^2} \right] + \left[\frac{n}{p^3} \right] + \dots$.

Suppose that $n = a_r p^r + a_{r-1} p^{r-1} + \dots + a_1 p + a_0$ is the p -adic expansion of the number n , in which

$0 \leq a_i \leq p-1, i = 0, 1, \dots, r$ and $a_r \neq 0$. In the following paper, the number p is always a prime. We will prove:

Theorem: The number of Sylow p -subgroup of the alternating group $A_n (n \geq 6)$ is

$$\frac{n!}{a_0! a_1! \dots a_r! p^{s(n)} (p-1)^{a_1 + 2a_2 + \dots + ra_r}}$$

Corresponding author: Rulin Shen, *E-mail: rulinshen@gmail.com

Lemma: 1 Let $X = \{1, 2, \dots, p\}$, $Y = \{1, 2, \dots, t, i_1, i_2, \dots, i_{p-t}\}$ and $X \cap Y = \{1, 2, \dots, t\}$ with $1 \leq t \leq p-1$.

Suppose that a, b are p -cycles on X and Y , respectively. Then the group $\langle a, b \rangle$ is not a p -group.

Proof: Suppose that $\langle a, b \rangle$ is a p -group. Clearly, it is a transitive permutation on the set $X \cap Y$, and its degree n is $|X \cap Y| = 2p - t$. Since $1 \leq t \leq p-1$, we have $1 < n < 2p$. On the other hand, the degree of $\langle a, b \rangle$ is a multiple of p since it is a p -group. In fact, we only need to choose a subgroup P of $\langle a, b \rangle$ whose kernel is 1 and order is maximal, then $n = |\langle a, b \rangle : P|$. So that $p \mid n$, which contradicts the facts that $p < n < 2p$.

Lemma: 2 The normalizer of Sylow p -subgroup of the symmetric group S_p ($p \geq 5$) is the Frobenius group $Z_p : Z_{p-1}$.

Proof: Let $Z_p = \langle (123\dots p) \rangle$ is a Sylow p -subgroup of S_p . Choose an element $b = (12435687\dots)$. It is easy to check that $(123\dots p)^b \in Z_p$, and hence $Z_p : \langle a, b \rangle \leq N_{S_p}(Z_p)$. On the other hand, S_p has $(p-1)!$ elements with order p exactly, so the number of subgroups of order p in S_p is $(p-2)!$. Obviously, the number $|S_p : N_{S_p}(Z_p)|$ of Sylow p -subgroups is also $(p-2)!$, thus $|N_{S_p}(Z_p)| = p(p-1)$. Therefore, $N_{S_p}(Z_p)$ is a Frobenius group with order $p(p-1)$.

2. Proof of Theorem: Denote by $A \wr B$ the wreath product of A and B , in which B regarded as a permutation group. Denote by $G^m := \overbrace{G \times G \times \dots \times G}^{m \text{ copies}}$, $G^{\wr m} := \overbrace{G \wr G \wr \dots \wr G}^{m \text{ copies}}$.

It is known that the Sylow p -subgroup P_{p^r} of S_{p^r} is $Z_p^{\wr r}$ (see 1.6.19 in [4]). Of course, it is a transitive group with degree $P^{s(p^r)}$. If G acts on the set Δ and $X \subseteq \Delta$, then N_X stands for the global stabilizer of subset X in G . We will use the following steps to complete the proof of the Theorem.

Step: 1 If $n = p^r$, then $N_{S_{p^r}}(P_{p^r}) \cong Z_p^{\wr r} : Z_{p-1}^r$.

Let $X = \{1, 2, \dots, p^r\}$, $X_i = \{p(i-1)+1, p(i-1)+2, \dots, p(i-1)+p\}$, where $1 \leq i \leq p^{r-1}$. Set $D = \{X_1, X_2, \dots, X_{p^{r-1}}\}$. Then

$$S(X)_D = (S(X_1) \times S(X_2) \times \dots \times S(X_{p^{r-1}})) : S_{p^{r-1}} \cong S_p \wr S_{p^{r-1}}.$$

Clearly, $S(X)_D$ must include a Sylow p -subgroup P_{p^r} of $S(X)$.

Let $K_i := S(X_i)$, $1 \leq i \leq p^{r-1}$, and $K := K_1$. The class of subgroups conjugate to K in $S(X)$ referred to as

the set of the fundamental subgroups in $S(X)$. Set

$$\Delta = Fun_{S(X)}(P_{p^r}) = \{K^x \mid x \in S(X), K^x \cap P_{p^r} \in Syl_p(K^x)\}.$$

By the definition of Δ , we have $\{K_i \mid 1 \leq i \leq p^{r-1}\} \subseteq \Delta$. Suppose that there exists an element $g \in S(X)$ such that $K^g \in \Delta$, and $K^g \neq K_i$ for each i . Now we let $K^g := S(g(X_1))$. of course $g(X_1) \neq K_i$ for each i . By the definition of Δ , we know that $K^g \cap P_{p^r} \in Syl_p(S(X))$. Then there exists a Sylow p -subgroup P of K^g such that $P \leq P_{p^r}$. Without loss of generality, we can assume that $a = (123\dots p) \in K \cap P_n$, then $\langle a, P \rangle \leq P_{p^r}$. Suppose that $X_i \cap g(X_1) \neq \emptyset$ for some i , and set $|X_i \cap g(X_1)| = t$, where $1 \leq t \leq p-1$. Then we can assume that $g(X_1) = \{1, 2, \dots, t, i_1, i_2, \dots, i_{p-t}\}$. By the Lemma 1, for any p -cycle b on $g(X_1)$, we have $\langle a, b \rangle$ is not a p -group. Then such P does not exist, contradicts. So $X_i \cap g(X_1) = \emptyset$ for any i , that is $g(X_1) \cap X = g(X_1) \cap (\cup_{i=1}^{p^{r-1}} X_i) = \emptyset$, which contradicts the fact that $K^g \leq S(X)$. Thus $\{K_i \mid 1 \leq i \leq p^{r-1}\} = \Delta$. Moreover, by the definition of Δ , we know that $K^x \cap P_{p^r} \in Syl_p(K^x)$ for any $K^x \in \Delta$. If $g \in N_{S(X)}(P_{p^r})$, i.e., $P_{p^r}^g = P_{p^r}$, then $K^{xg} \cap P_{p^r} \in Syl_p(K^{xg})$, and hence $K^{xg} \in \Delta$. Thus $g \in N_{S(X)}(\Delta)$, then

$$N_{S(X)}(P_{p^r}) \leq N_{S(X)}(\Delta) = S(X)_D \cong S_p \wr S_{p^{r-1}}.$$

Without loss of generality, we suppose that P_{p^r} is a Sylow p -subgroup of $S_p \wr S_{p^{r-1}}$. of course, Now we

$$N_{S(X)}(P_{p^r}) = N_{S_p \wr S_{p^{r-1}}}(P_{p^r}). \quad \text{denote by} \quad G := S_p \wr S_{p^{r-1}}, \quad A := S_p^{p^{r-1}},$$

$A := S_p^{p^{r-1}}, B := S_{p^{r-1}}$, $P_A = P_{p^r} \cap A, P_B = P_{p^r} \cap B$. Since $G = A : B$, we have $N_B(P_B) \cong N_{G/A}(P_{p^r} A / A) = N_G(P_{p^r}) A / A \cong N_G(P_{p^r}) / A \cap N_G(P_{p^r}) = N_G(P_{p^r}) / N_A(P_{p^r})$, so that

$N_G(P_{p^r}) = N_A(P_{p^r}) : N_B(P_B)$. In the sequel, we determine the structure of $N_A(P_{p^r})$. Obviously, $P_{p^r} = Z_p^{1r} = Z_p \wr Z_p^{1r-1} = Z_p^{p^{r-1}} : Z_p^{1r-1}$. Choose an element $x = (x_1, x_2, \dots, x_{p^{r-1}}; 1) \in N_A(P_{p^r})$, then we have $g^x \in P_{p^r}$

for any element $g = (g_1, g_2, \dots, g_{p^{r-1}}; g_0) \in P_{p^r}$, where $g_0 \in Z_p^{1r-1}$. That is

$$g^x = (x_1^{-1} g_1 x_1^{g_0}, x_2^{-1} g_2 x_2^{g_0}, \dots, x_{p^{r-1}}^{-1} g_{p^{r-1}} x_{(p^{r-1})}^{g_0}; g_0) \in Z_p^{p^{r-1}} : Z_p^{1r-1}.$$

Since g_0 can be chosen randomly, we can get $x_i \in N_{S_p}(Z_p)$ if we choose $g_0 = 1$, and

hence $x_i = x_{i^{g_0}}$. Furthermore, $Z_p^{i_{r-1}}$ is a transitive group with degree $p^{s(p^{r-1})}$, then for any pairs $i, j \in \{1, 2, \dots, p^{r-1}\}$, there exists an element $g_0 \in Z_p^{i_{r-1}}$ such that $i^{g_0} = j$, and

hence $x_1 = x_2 = \dots = x_{p^{r-1}}$. Conversely, if $x = (x_1, x_2, \dots, x_{p^{r-1}}; 1)$ and $x_1 \in N_{S_p}(Z_p)$, then $x \in N_A(P_{p^r})$. Thus

$N_A(P_{p^r}) = Z_p^{p^{r-1}} : Z_{p-1}$. So we have $N_A(P_{p^r}) = N_A(P_{p^r}) : N_B(P_B) \cong (Z_p^{p^{r-1}} : Z_{p-1}) : N_B(P_B)$. Therefore, $N_{S_{p^r}}(P_{p^r}) \cong (Z_p^{p^{r-1}} : Z_{p-1}) : N_{S_{p^{r-1}}}(P_{p^{r-1}})$. By the induction for r , we can obtain $N_{S_{p^r}}(P_{p^r}) \cong Z_p^{i_r} : Z_{p-1}^r$.

Step: 2 If $n = a_r p^r + k$, then $N_{S_n}(P_n) \cong N_{S_k}(P_k) \times (N_{S_{p^r}}(P_{p^r}) \wr S_{a_r})$, here $0 \leq k < p^r$.

Let $\Omega = \{1, 2, 3, \dots, n\}$, $X_i = \{p^{i-1}, p^{i-1} + 1, \dots, p^{i-1} + p^r - 1\}$ in which $1 \leq i \leq a_r$ and $X_0 = \Omega \setminus \cup_{i=1}^{a_r} X_i$. Set

$D = \{X_0, X_1, X_2, \dots, X_{p^{r-1}}\}$. Then

$$S(\Omega)_D = S(X_0) \times ((S(X_1) \times S(X_2) \times \dots \times S(X_{p^{r-1}})) : S_{a_r}) \cong S_k \times (S_{p^r} \wr S_{a_r}).$$

Clearly, $S(\Omega)_D$ includes a Sylow p -subgroup P_n of $S(\Omega)$. Denote by $K_i := S(X_i)$, $1 \leq i \leq a_r$, and $K := K_1$.

Set $\Delta = Fun_{S(\Omega)}(P_n) = \{K^x \mid x \in S(\Omega), K^x \cap P_n \in Syl_p(K^x)\}$.

In this case, Δ is a maximal set of pair-wise commuting fundamental subgroups of K in S_{p^r} . It is clear

that $\{K_i \mid 1 \leq i < p^{r-1}\} \subseteq \Delta$. If there exists an element $g \in S(\Omega)$ such that $K^g \in \Delta$ and $K^g \neq K_i$ for each i .

Denote by $K^g := S(g(X_1))$. Obviously, $g(X_1) \neq K_i$ for each i . Since $K^g \cap P_{p^r} \in Syl_p(S(\Omega))$, then there is a

Sylow p -subgroup P of K^g such that $P \leq P_n$. Now we can assume that $a = (123 \dots p) \in K \cap P_n$,

then $\langle a, P \rangle \leq P_n$. Suppose that $X_i \cap g(X_1) \neq \emptyset$ for some i , and set $|X_i \cap g(X_1)| = t$, where $1 \leq t \leq p-1$.

Without loss of generality, we can assume that $g(X_1) = \{1, 2, \dots, t, i_1, i_2, \dots, i_{p-t}\}$. Similarly, for any p -cycle b

on $g(X_1)$, we have $\langle a, b \rangle$ is not a p -group by Lemma 1. Then such P does not exist, contradicts. So

$X_i \cap g(X_1) = \emptyset$ for any i , then $g(X_1) \subseteq X_0$, which contradicts the fact that $|g(X_1)| > |X_0|$. Therefore,

$\{K_i \mid 1 \leq i \leq p^{r-1}\} = \Delta$. Then

$$N_{S(\Omega)}(P_n) \leq N_{S(\Omega)}(\Delta) = S(\Omega)_D \cong S_k \times (S_{p^r} \wr S_{a_r}).$$

We can assume that P_n is a chosen Sylow p -subgroup of It is $S_k \times (S_{p^r} \wr S_{a_r})$. obvious that

$$N_{S(\Omega)}(P_n) = N_{S_k \times (S_{p^r} \wr S_{a_r})}(P_n).$$

It is easy to see that $P_n = P_k \times P_{p^r}^{a_r}$, so $N_{S(\Omega)}(P_n) = N_{S_k}(P_k) \times (N_{S_{p^r}}(P_{p^r}) \wr S_{a_r})$.

Step: 3 Now we continue to decompose k into the sum $a_{r-1}p^{r-1} + k_1$, in which $0 \leq k_1 < p^{r-1}$. Next we repeat to

decompose k_1 , and so on. At last we can get the p -adic expansion $a_r p^r + a_{r-1} p^{r-1} + \dots + a_1 p + a_0$ by r steps.

Using the induction for r by the conclusions of step 1 and 2, we can obtain the result of the normalizer for general n :

$$N_{S_n}(P_n) \cong ((Z_p^{a_r} : Z_{p-1}^r) \wr S_{a_r}) \times ((Z_p^{a_{r-1}} : Z_{p-1}^{r-1}) \wr S_{a_{r-1}}) \times \dots \times ((Z_p^{a_1} : Z_{p-1}) \wr S_{a_1}) \times S_{a_0}.$$

In the case of $p = 2$, it is not hard to see $|N_{S_n}(P_n)| = |P_n|$ since $a_i = 0$ or 1 ,

then S_n has a self-normalized Sylow 2-subgroup. Since $|S_n : A_n| = 2$, we have the Sylow p -number of S_n is

same as the one of A_n for $p > 2$. If $p = 2$, we use the result of Kondrat'ev's [3]: A_n has a self-normalized

Sylow 2-subgroup for $n \geq 6$.

So the number of Sylow p -subgroups of the alternating group A_n ($n \geq 6$) is

$$\frac{n!}{a_0! a_1! \dots a_r! p^{s(n)} (p-1)^{a_1 + 2a_2 + \dots + ra_r}}.$$

REFERENCE:

[1] Carter R., Fong P., The Sylow 2-subgroups of the finite classical groups, J. Algebra, 1(1964) 139-151.
 [2] Dmitruk, Yu. V., Sushchanskii V. I., Structure of Sylow 2-subgroups of the alternating groups and normalizers of Sylow subgroups in the symmetric and alternating groups, Ukrainian Mathematical Journal, 3(1981)235-241.
 [3] Kondrat'ev A. S., Normalizers of the Sylow 2-Subgroups in Finite Simple Groups, Mathematical Notes 78(34), 338-346.
 [4] Robinson D. J. S., A Course in the Theory of Groups, Springer-Verlag, 1982.
 [5] Zhang J., Sylow numbers of finite groups, J. Algebra., 176(1995), 111-12.

¹Project supported by the NNSF of China (No.11026195) and the foundation of Educational Department of Hubei Province in China (No.Q20111901).
