

EXISTENCE OF QUASILINEAR NEUTRAL IMPULSIVE INTEGRODIFFERENTIAL EQUATIONS

R. Prabakaran\*<sup>1</sup> and V. Vinoba<sup>2</sup>

<sup>1</sup>*Department of Science and Humanities, Sri Krishna College of Technology, Coimbatore-42, India.*

<sup>2</sup>*Department of Mathematics, K. N. Govt. arts College for women, Thanjavur, India.*

(Received on: 17-05-14; Revised & Accepted on: 08-07-14)

ABSTRACT

In this paper, we devoted to study the existence of mild solutions for quasilinear impulsive integrodifferential equation in Banach spaces. The results are established by using Hausdorff's measure of noncompactness and the fixed point theorems. Application is provided to illustrate the theory.

**Keywords:** Mild solution, neutral differential equation, impulsive condition, Hausdorff measure of noncompactness, fixed point theorem.

**2010 AMS Subject Classification:** 34A37, 34K05, 34K40, 47H10.

1. INTRODUCTION

In various fields of engineering and physics, many problems that are related to linear viscoelasticity, nonlinear elasticity have mathematical models and are described by the problems of differential or integral equations or integrodifferential equations. Our work centers on the problems described by the integrodifferential models. It is important to note that when we describe the systems which are functions of space and time by partial differential equations, in some situations, such a formulation may not accurately model the physical system because, while describing the system as a function at a given time, it may fail to take into account the effect of past history. Neutral differential equations arise in many areas of applied mathematics and for this reason these equations have received much attention during the last few decades [1, 2, 3]. A good guide to the literature for neutral functional differential equations is the book by Hale and Verduyn Lunel [4] and the references therein. The existence of solution to evolution equations with nonlocal conditions in Banach space was studied first by Byszewski [5, 6]. Byszewski and Lakshmikanthan [7] proved an existence and uniqueness of solutions of a nonlocal Cauchy problem in Banach spaces. Ntouyas and Tsamatos [8] studied the existence for semilinear evolution equations with nonlocal conditions. The problem of existence of solutions of evolution equations in Banach space has been studied by several authors [9, 10].

However, one may easily visualize that abrupt changes such as shock, harvesting and disasters may occur in nature. These phenomena are short time perturbations whose duration is negligible in comparison with the duration of the whole evolution process. Consequently, it is natural to assume, in modeling these problems, that these perturbations act instantaneously, that is in the form of impulses. The theory of impulsive differential equation [11, 12, 13] is much richer than the corresponding theory of differential equations without impulsive effects. The impulsive condition

$$\Delta u(t_i) = u(t_i^+) - u(t_i^-) = I_i(u(t_i^-)), i=1, 2, \dots, m,$$

is a combination of traditional initial value problems and short-term perturbations whose duration is negligible in comparison with the duration of the process. Liu [14] discussed the iterative methods for the solution of impulsive functional differential systems.

Measures of noncompactness are a very useful tool in many branches of mathematics. They are used in the fixed point theory, linear operators theory, theory of differential and integral equations and others [15]. There are two measures which are the most important ones. The Kuratowski measure of noncompactness  $\sigma(X)$  of a bounded set  $X$  in a metric space is defined as infimum of numbers  $r>0$  such that  $X$  can be covered with a finite number of sets of diameter

*Corresponding author: R. Prabakaran\*<sup>1</sup>*

<sup>1</sup>*Department of Science and Humanities, Sri Krishna College of Technology, Coimbatore-42, India.*

smaller than  $r$ . The Hausdorff measure of noncompactness  $\chi(X)$  defined as infimum of numbers  $r > 0$  such that  $X$  can be covered with a finite number of balls of radii smaller than  $r$ . There exist many formulae on  $\chi(X)$  in various spaces [15, 18].

Let  $E$  be a Banach space and  $F$  be a subspace of  $E$ . Let  $\chi_E(X), \chi_F(X), \sigma_E(X), \sigma_F(X)$  denote Hausdorff and Kuratowski measures in spaces  $E, F$ , respectively. Then, for any bounded  $X \subset F$  we have  $\chi_E(X) \leq \chi_F(X) \leq \sigma_F(X) = \sigma_E(X) \leq 2\chi_E(X)$ . The notion of a measure of weak compactness was introduced by De Blasi [16] and was subsequently used in numerous branches of functional analysis and the theory of differential and integral equations. Several authors have studied the measures of noncompactness in Banach spaces [17, 18, 19]. Motivated by [9, 15, 20, 21], in this paper, we study the existence results for quasilinear equation represented by first-order neutral integrodifferential equations using the semigroup theory and the measure of noncompactness.

**2. PRELIMINARIES**

We consider the quasilinear integrodifferential equations with impulsive and nonlocal condition of the form

$$\frac{d}{dt}[x(t) + e(t, x(t), \int_0^t k(t, s, x(s))ds)] + A(t, x(t))x(t) = f(t, x(t)) + \int_0^t g(t, s, x(s))ds, t \in [0, b], t \neq t_k, \tag{1}$$

$$x(0) + h(x) = x_0, \tag{2}$$

$$\Delta x(t_k) = I_k(x(t_k)), \quad k = 1, 2, 3, \dots, n, \tag{3}$$

where  $A: [0, b] \times X \rightarrow X$  is a continuous function in Banach space  $X, x_0 \in X, f: [0, b] \times X \rightarrow X, g: I \times X \rightarrow X, h: PC([0, b], X) \rightarrow X, e: [0, b] \times X \times X \rightarrow X, k: I \times X \rightarrow X$  and  $\Delta x(t_k) = x(t_k^+) - x(t_k^-)$ , for all  $k=1, 2, \dots, m, 0=t_0 < t_1 < t_2 < \dots < t_m < t_{m+1} = b$ ; constitutes an impulsive condition. Here  $A = \{(t, s): 0 \leq s \leq t \leq b\}$ .

Let  $X$  be a Banach space with norm  $\|\cdot\|$  Let  $PC([0, b], X)$  consist of functions  $u$  from  $[0, b]$  into  $X$ , such that  $x(t)$  is continuous at  $t \neq t_i$  and left continuous at  $t = t_i$  and the right limit  $x(t_i^+)$  exists, for  $i=1, 2, 3, \dots, n$ . Evidently  $PC([0, b], X)$  is a Banach space with the norm

$$\|x\|_{PC} = \sup_{t \in [0, b]} \|x(t)\|,$$

and denoted  $L([0, b], X)$  by the space of  $X$ -valued Bochner integrable functions on  $[0, b]$  with the form

$$\|x\|_L = \int_0^b \|x(t)\| dt.$$

The Hausdorff's measure of noncompactness  $\chi_Y$  is defined by

$$\chi(B) = \inf \{r > 0, B \text{ can be covered by finite number of balls with radii } r\},$$

for bounded set  $B$  in a Banach space  $Y$ .

**Lemma 2.1:** [15]. Let  $Y$  be a real Banach space and  $B, E \subseteq Y$  be bounded, with the following properties:

1.  $B$  is precompact if and only if  $\chi_X(B) = 0$ .
2.  $\chi_Y(B) = \chi_Y(\bar{B}) = \chi_Y(\text{con}B)$ , where  $\bar{B}$  and  $\text{con}B$  mean the closure and convex hull of respectively.  $B$
3.  $\chi_Y(B) \leq \chi_Y(E)$ , where  $B \subseteq E$ .
4.  $\chi_Y(B + E) \leq \chi_Y(B) + \chi_Y(E)$ , where  $B + E = \{x + y: x \in B, y \in E\}$ .
5.  $\chi_Y(B \cup E) \leq \max \{\chi_Y(B), \chi_Y(E)\}$ .
6.  $\chi_Y(\lambda B) \leq |\lambda| \chi_Y(B)$ , for any  $\lambda \in R$ .
7. If the map  $F: D(F) \subseteq Y \rightarrow Z$  is Lipschitz continuous with constant  $r$ , then  $\chi_Z(FB) \leq r\chi_Y(B)$ , for any bounded subset  $B \subseteq D(F)$ , where  $Z$  be a Banach space.

8.  $\chi_Y(B) = \inf \{d_Y(B, E); E \subseteq Y \text{ is precompact}\} = \inf \{d_Y(B, E); E \subseteq Y \text{ is finite valued}\}$ , where  $d_Y(B, E)$  means the non-symmetric (or symmetric) Hausdorff distance between  $B$  and  $E$  in  $Y$ .
9. If  $\{W_n\}_{n=1}^{+\infty}$  is decreasing sequence of bounded closed nonempty subsets of  $Y$  and  $\lim_{n \rightarrow \infty} \chi_Y(W_n) = 0$ , then  $+\infty W_n$  is nonempty and compact in  $Y$ .

The map  $F:W \subseteq Y \rightarrow Y$  is said to be a  $\chi_Y$ -contraction if there exists a positive constant  $r < 1$  such that  $\chi_Y(F(B)) \leq r\chi_Y(B)$  for any bounded closed subset  $B \subseteq W$ , where  $Y$  is a Banach space.

**Lemma 2.2:** (Darbo-Sadovskii [15]). If  $W \subseteq Y$  is bounded closed and convex, the continuous map  $F:W \rightarrow W$  is a  $\chi_Y$ -contraction, the map  $F$  has atleast one fixed point in  $W$ .

We denote by  $\chi$  the Hausdorff's measure of noncompactness of  $X$  and also denote  $\chi_c$  by the Hausdorff's measure of noncompactness of  $PC([0, b], X)$ .

Before we prove the existence results, we need the following Lemmas.

**Lemma 2.3:** [22] If  $W \subseteq PC([0, b], X)$  is bounded, then  $\chi(W(t)) \leq \chi_c(W)$ , for all  $t \in [0, b]$ , where  $W(t) = \{u(t); u \in W\} \subseteq X$ . Furthermore if  $W$  is equicontinuous on  $[0, b]$ , then  $\chi(W(t))$  is continuous on  $[0, b]$  and  $\chi_c(W) = \sup \{\chi(W(t)), t \in [0, b]\}$ .

**Lemma 2.4:** [22, 23]. If  $\{u_n\}_{n=1}^{\infty} \subset L^1([0, b], X)$  is uniformly integrable, then the function  $\chi(\{u_n(t)\}_{n=1}^{\infty})$  is measurable and

$$\chi\left(\left\{\int_0^t u_n(s) ds\right\}_{n=1}^{\infty}\right) \leq 2 \int_0^t \chi(\{u_n(s)\}_{n=1}^{\infty}) ds. \tag{4}$$

**Lemma 2.5:** If  $W \subseteq PC([0, b], X)$  is bounded and equicontinuous, then  $\chi(W(t))$  is continuous and

$$\chi\left(\int_0^t W(s) ds\right) \leq \int_0^t \chi(W(s)) ds, \text{ for all } t \in [0, b], \tag{5}$$

where  $\int_0^t W(s) ds = \{\int_0^t u(s) ds : u \in W\}$ .

The  $C_0$  semigroup  $U_u(t, s)$  is said to be equicontinuous if  $(t, s) \rightarrow \{U_u(t, s)u(s); u \in B\}$  is equicontinuous for  $t > 0$ , for all bounded set  $B$  in  $X$ . The following lemma is obvious.

**Lemma 2.6:** If the evolution family  $\{U_u(t, s)\}_{0 \leq s \leq t \leq b}$  is equicontinuous and  $\eta \in L([0, b], R^+)$ , then the set  $\{\int_0^t U_u(t, s)u(s) ds, \|u(s)\| \leq \eta(s), \text{ for a.e } s \in [0, b]\}$  is equicontinuous for  $t \in [0, b]$ .

We know that, for any fixed  $u \in PC([0, b], X)$  there exist a unique continuous function  $U_u: [0, b] \times [0, b] \rightarrow B(X)$  defined on  $[0, b] \times [0, b]$  such that

$$U_u(t, s) = I + \int_s^t A_u(w)U_u(w, s)dw, \tag{6}$$

where  $B(X)$  denote the Banach space of bounded linear operators from  $X$  to  $X$  with the norm  $\|F\| = \sup \{\|Fu\|; \|u\| = 1\}$ , and  $I$  stands for the identity operator on  $X$ ,  $A_u(t) = A(t, u(t))$ , we have

$$U_u(t, t) = I, U_u(t, s)U_u(s, r) = U_u(t, r), (t, s, r) \in [0, b] \times [0, b] \times [0, b],$$

$$\frac{\partial U_u(t, s)}{\partial t} = A_u(t)U_u(t, s), \text{ for almost all } t, s \in [0, b].$$

### 3 THE EXISTENCE OF MILD SOLUTION

**Definition 3.1:** A function  $x \in PC([0, b], X)$  is said to be a mild solution of (1)–(3) if it satisfies the integral equation

$$\begin{aligned}
 x(t) = & U_x(t, 0)x_0 - U_x(t, 0)h(x) + U_x(t, 0)e(0, x(0), 0) - e\left(t, x(t), \int_0^t k(t, s, x(s))ds\right) \\
 & + \int_0^t A(s, x(s))U_x(t, s)e\left(s, x(s), \int_0^s k(s, \tau, x(\tau))d\tau\right)ds \\
 & + \int_0^t U_x(t, s)[f(s, x(s)) + \int_0^s g(s, \tau, x(\tau))d\tau]ds + \sum_{0 < t_k < t} U_x(t, t_k)I_k(x(t_k)), 0 \leq t \leq b.
 \end{aligned}$$

In this paper, we denote  $M_0 = \sup\{\|U_x(t, s)\| : (t, s) \in [0, b] \times [0, b]\}$ , for all  $x \in X$ . Without loss of generality, we let  $x_0 = 0$ .

Assume the following conditions:

(H1) The evolution family  $\{U_x(t, s)\}_{0 \leq s \leq t \leq b}$  generated by  $A(t, x(t))$  is equicontinuous and  $\|U_x(t, s)\| \leq M_0$ , for almost  $t, s \in [0, b]$ .

(H2) (i) The function  $h: PC([0, b] \times X) \rightarrow X$  is continuous and compact.

(ii) There exists  $N_0 > 0$  such that  $\|h(x)\| \leq N_0$ , for all  $x \in PC([0, b]; X)$ .

(H3) (i) The nonlinear function  $f: [0, b] \times X \rightarrow X$  satisfies the Carathéodory-type conditions; that is,  $f(\cdot, x)$  is measurable for all  $x \in X$  and  $f(t, \cdot)$  is continuous, for a.e  $t \in [a, b]$ .

(ii) There exists a function  $\alpha \in L([0, b], R^+)$  such that for every  $x \in X$ , we have

$$\|f(t, x)\| \leq \alpha(t) \|x\|, \text{ a.e } t \in [0, b].$$

(iii) There exists a function  $m_f \in L([0, b], R^+)$  such that, for every bounded  $K \subset X$ , we have

$$\chi(f(t, K)) \leq m_f(t) \chi(K), \text{ a.e } t \in [0, b].$$

(H4) (i) The nonlinear function  $g: [0, b] \times [0, b] \times X \rightarrow X$  satisfies the Carathéodory-type conditions; i.e.,  $g(\cdot, \cdot, x)$  is measurable, for all  $x \in X$  and  $g(t, s, \cdot)$  is continuous for a.e  $t \in [a, b]$ .

(ii) There exist two functions  $\beta_1 \in L([0, b], R^+)$  and  $\beta_2 \in L([0, b], R^+)$  such that for every  $x \in X$ , we have

$$\|g(t, s, x(s))\| \leq \beta_1(t) \|x(s)\| + \beta_2(t), \text{ a.e } t \in [0, b].$$

(iii) There exist functions  $m_g, n_g \in L([0, b], R^+)$  such that, for every bounded  $K \subset X$ , we have

$$\chi(g(t, s, K)) \leq m_g(t) n_g(s) \chi(K), \text{ a.e } t \in [0, b].$$

Assume that the finite bound of  $\int_0^t m_g(s) ds$  is  $G_0$ .

(H5) (i) The function  $e: [0, b] \times X \times X \rightarrow X$  satisfies the Carathéodory-type conditions; that is,  $e(\cdot, x, x_1)$  is measurable, for all  $x, x_1 \in X$  and  $e(t, \cdot, \cdot)$  is continuous, for a.e  $t \in [0, b]$ .

(ii) There exists a function  $\gamma \in L([0, b], R^+)$  such that for every  $x, x_1 \in X$ , we have

$$\|e(t, x, x_1)\| \leq \gamma(t) (\|x\| + \|x_1\|), \text{ a.e } t \in [0, b].$$

(iii) The nonlinear function  $q: [0, b] \times [0, b] \times X \rightarrow X$  satisfies the Carathéodory-type conditions; i.e.

$k(\cdot, \cdot, x)$  is measurable, for all  $x \in X$  and  $k(t, s, \cdot)$  is continuous, for a.e  $t \in [0, b]$ .

(iv) There exist two functions  $\omega_1 \in L([0, b], R^+)$  and  $\omega_2 \in L([0, b], R^+)$  such that for every  $x \in X$ , we have

$$\|k(t, s, x(s))\| \leq \omega_1(t) \|x(s)\| + \omega_2(t), \text{ a.e } t \in [0, b].$$

(v) There exists a function  $\eta \in L([0, b], R^+)$  such that for every  $x, x_1 \in X$ , we have

$$\|A(t, x(t))e(t, x, x_1)\| \leq e(t, x, x_1), \text{ a.e } t \in [0, b],$$

(vi) There exists a function  $m_e \in L([0, b], R^+)$  such that, for every bounded  $K, K_1 \subset X$ , we have

$$\chi(e(t, K, K_1)) \leq m_e(t)(\chi(K) + \chi(K_1)), \text{ a.e } t \in [0, b].$$

Assume that the finite bound of  $\int_0^t m_e(s)ds$  is  $G_1$ .

(vii) There exist functions  $m_k, n_k \in L([0, b], R^+)$  such that for every bounded  $K \subset X$ , we have

$$\chi(k(t, s, K)) \leq m_k(t)n_k(s)\chi(K), \text{ a.e } t \in [0, b].$$

Assume that the finite bound of  $\int_0^t m_e(s)ds$  is  $G_2$ .

(H6) For every  $t \in [0, b]$  and there exist positive constants  $N_1$  and  $N_2$ , the scalar equation

$$m(t) = M_0 N_0 + \gamma_1(1 + m(s)) + M_0 \gamma_0 + M_0 C_1 \omega(t)(1 + m(s)) + \gamma(t) C_1 \int_0^t \eta(t) \omega_1(s) ds \\ + M_0 \int_0^t [\alpha(s)(1 + m(s)) ds + C_0 \int_0^t \beta_1(s)(1 + m(s)) ds + \sum_{k=1}^n d_k],$$

where  $C_0 = \int_0^s \beta(t) dt$ .

(H7)  $I_k : X \rightarrow X$  is continuous. There exist constants  $d_k > 0, k = 1, 2, 3, \dots, n$  such that

$$\|I_k(x(t_k))\| \leq \sum_{k=1}^n d_k, \text{ where, } k = 1, 2, 3, \dots, n.$$

For any bounded subset  $K \subset X$ , and there is a constant  $l_k > 0$  such that

$$\chi(I_k(K)) \leq \sum_{k=1}^n l_k \chi(K), \text{ } k = 1, 2, \dots, n.$$

**Theorem 3.1:** If assumptions  $(H_1) - (H_7)$  holds, then the quasilinear neutral impulsive problem (1) - (3) has at least one mild solution.

**Proof:** Let  $m(t)$  be a solution of the scalar equation

$$m(t) = M_0 N_0 + \gamma_1(1 + m(s)) + M_0 \gamma_0 + M_0 C_1 \omega(t)(1 + m(s)) + \gamma(t) C_1 \int_0^t \eta(t) \omega_1(s) ds \\ + M_0 \int_0^t [\alpha(s)(1 + m(s)) ds + C_0 \int_0^t \beta_1(s)(1 + m(s)) ds + \sum_{k=1}^n d_k]. \tag{7}$$

Let us assume that the finite bound of  $\int_0^t \beta_2(s) ds$  is  $C_0$ , for  $t \in [0, b]$ . is  $C_0$ , for  $t \in [0, b]$ . Consider the map

$F : PC([0, b], X) \rightarrow PC([0, b], X)$  defined by

$$(Fx)(t) = U_x(t, 0)h(x) + U_x(t, 0)e(0, x(0), 0) - e\left(t, x(t), \int_0^t k(t, s, x(s)) ds\right) \\ + \int_0^t A(s, x(s))U_x(t, s)e\left(s, x(s), \int_0^s k(s, \tau, x(\tau)) d\tau\right) ds \\ + \int_0^t U_x(t, s)[f(s, x(s)) + \int_0^s g(s, \tau, x(\tau)) d\tau] ds \\ + \sum_{0 < t_k < t} U_x(t, t_k)I_k(x(t_k)), \text{ } 0 \leq t \leq b, \text{ for all } x \in PC([0, b], X). \tag{8}$$

Let us take  $W_0 = \{x \in PC([0, b], X), \|x(t)\| \leq m(t), \text{ for all } t \in [0, b]\}$ . Then  $W_0 \subseteq PC([0, b], X)$  is bounded and convex. We define  $W_1 = \overline{con} K(W_0)$ , where  $\overline{con}$  means the closure of the convex hull in  $PC([0, b], X)$ . As  $U_x(t, s)$  is equicontinuous,  $h$  is compact and  $W_0 \subseteq PC([0, b], X)$  is bounded, due to Lemma 2.6 and using the assumptions,  $W_1 \subseteq PC([0, b], X)$  is bounded closed convex nonempty and equicontinuous on  $[0, b]$ .

For any  $x \in F(W_0)$ , we know

$$\begin{aligned} \|x(t)\| &\leq \|U_x(t,0)h(x)\| + \|U_x(t,0)e(0,x(0),0)\| + \left\| e\left(t,x(t),\int_0^t k(t,s,x(s))ds\right) \right\| \\ &+ \int_0^t \left\| A(t,x(t))U_x(t,s)e\left(t,x(t),\int_0^s k(s,\tau,x(\tau))d\tau\right) \right\| ds \\ &+ \int_0^t \left\| U_x(t,s)[f(s,x(s)) + \int_0^s g(s,\tau,x(s))d\tau] \right\| ds + \sum_{0 < t_k < t} \|U_x(t,t_k)I_i(x(t_k))\| \\ &\leq M_0N_0 + M_0\gamma_0 + \gamma_1(1 + \|x\|) + \int_0^t k(t,s,x(s))ds \\ &+ M_0\eta(t) \int_0^t \left\| e\left(t,x(t),\int_0^s k(s,\tau,x(\tau))d\tau\right) \right\| ds \\ &+ M_0 \left[ \int_0^t \|f(s,x(s))\| ds + \int_0^t \int_0^s \|g(s,\tau,x(\tau))\| d\tau ds \right] + M_0 \sum_{k=1}^n \|I_k(x(t_k))\| \\ &\leq M_0N_0 + M_0\gamma_0 + \gamma_1(1 + \|x\|) + \omega_1(t) \int_0^t \omega_2(s)(1 + \|x\|) ds \\ &+ M_0 \left[ \int_0^t [\eta(s)\gamma(s)ds + \int_0^s \omega_1(s)\omega_2(\tau)dsd\tau](1 + \|x\|) \right] \\ &+ M_0 \int_0^t \alpha(s)(1 + \|x(s)\|) ds + M_0 \int_0^t \int_0^s \beta_1(s)\beta_2(\tau)(1 + \|x(\tau)\|) d\tau ds + M_0 \sum_{k=1}^n d_k \\ &\leq M_0N_0 + \gamma_1(1 + m(s)) + M_0\gamma_0 + M_0C_1\omega(t)(1 + m(s)) + \eta(t)\gamma(t)C_1 \int_0^t \omega_1(s) ds \\ &+ M_0 \left[ \int_0^t [\alpha(s)(1 + m(s))ds + C_0 \int_0^t \beta_1(s)(1 + m(s))ds + \sum_{k=1}^n d_k] \right] \\ &= m(t). \end{aligned}$$

It follows that  $W_1 \subset W_0$ . We define  $W_{n+1} = \overline{\text{con } F(W_n)}$ , for  $n=1,2,3,\dots$ . From above we know that  $\{W_n\}_{n=1}^\infty$  is a decreasing sequence of bounded, closed, convex, equicontinuous on  $[0,b]$  and nonempty subsets in  $\text{PC}([0,b], X)$ .

Now for  $n \geq 1$  and  $t \in [0,b]$ ,  $W_n(t)$  and  $F(W_n(t))$  are bounded subsets of  $X$ , hence, for any  $\varepsilon > 0$ , there is a sequence  $\{x_k\}_{k=1}^\infty \subseteq W_n$ , such that (see, e.g. [24], pp.125).

$$\begin{aligned} \chi(W_{n+1}(t)) &= \chi(FW_n(t)) \\ &\leq 2\chi(e(t,\{x_k(s)\}_{k=1}^\infty, \int_0^t k(t,s,\{x_k(s)\}_{k=1}^\infty)ds)) \\ &+ 2M_0\eta(t) \int_0^t \chi(e(s,\{x_k(s)\}_{k=1}^\infty, \int_0^s k(s,\tau,\{x_k(\tau)\}_{k=1}^\infty)d\tau)) ds \\ &+ 2M_0 \int_0^t \chi(f(s,\{x_k(s)\}_{k=1}^\infty)) ds + 4M_0 \int_0^t \int_0^s \chi(g(s,\tau,\{u_k(\tau)\}_{k=1}^\infty)) d\tau ds \\ &+ 2M_0 \sum_{i=1}^n \chi(I_i(\{u_k(t_k)\}_{k=1}^\infty)) + \varepsilon \\ &\leq 2m_\varepsilon(t)\chi\{x_k(t)\}_{k=1}^\infty + 2m_k(t) \int_0^t m_k(s)\chi\{x_k(s)\}_{k=1}^\infty ds \\ &+ 2M_0\eta(t) \left[ \int_0^t m_\varepsilon(s)\chi\{x_k(s)\}_{k=1}^\infty ds + 2 \int_0^t \int_0^s m_k(s)m_k(\tau)\chi\{x_k(\tau)\}_{k=1}^\infty d\tau ds \right] \\ &+ 2M_0 \int_0^t m_f(s)\chi(\{u_k(s)\}_{k=1}^\infty) ds + 4M_0 \int_0^t \int_0^s m_g(s)n_g(\tau)\chi(\{u_k(\tau)\}_{k=1}^\infty) d\tau ds \\ &+ 2M_0 \sum_{i=1}^n l_i \chi(\{u_k(t_k)\}_{k=1}^\infty) + \varepsilon \end{aligned}$$

$$\begin{aligned} &\leq 2[m_e(t) + m_k(t)G_2 + M_0\eta(t)G_1]\chi(W_n(t)) + 2M_0\left[\int_0^t\{2G_2m_k(s) \right. \\ &\left. + m_f(s)\}\chi(W_n(s))ds + 2G_0\int_0^t n_g(s)\chi(W_n(s))ds\right] + 2M_0\sum_{k=1}^n I_k\chi(W_n(t_k)) + \varepsilon. \end{aligned}$$

Since  $\varepsilon > 0$  is arbitrary, it follows that from the above inequality that

$$\begin{aligned} \chi(W_{n+1}(t)) &\leq 2[m_e(t) + m_k(t)G_2 + M_0\eta(t)G_1]\chi(W_n(t)) \\ &\quad + 2M_0\left[\int_0^t[2G_2m_k(s) + m_f(s) + 2G_0n_g(s)]\chi(W_n(s))ds \right. \\ &\quad \left. + 2M_0\sum_{k=1}^n I_k\chi(W_n(t_k))\right], \text{ for all } t \in [0, b]. \end{aligned} \tag{9}$$

Because  $W_n$  is decreasing for  $n$ , we have

$$\lambda(t) = \lim_{n \rightarrow \infty} \chi(W_n(t)),$$

for all  $t \in [0, b]$ . From (9), we have

$$\begin{aligned} \lambda(t) &\leq 2[m_e(t) + m_k(t)G_2 + M_0\eta(t)G_1]\lambda(t) \\ &\quad + 2M_0\left[\int_0^t[2G_2m_k(s) + m_f(s) + 2G_0n_g(s)]\lambda(s)ds + \sum_{k=1}^n I_k\lambda(t_k)\right], \end{aligned}$$

for  $t \in [0, b]$ , which implies that  $\lambda(t) = 0$ , for all  $t \in [0, b]$ . By Lemma 2.3, we know that  $\lim_{n \rightarrow \infty} \chi(W_n(t)) = 0$ .

Using Lemma 2.1 we know that  $W = \bigcap_{n=1}^{\infty} W_n$  is convex compact and nonempty in  $PC([0, b], X)$  and  $F(W) \subset W$ . By the Schauder fixed point theorem, there exist at least one mild solution  $u$  of the initial value problem (1)–(3), where  $x \in W$  is a fixed point of the continuous map  $F$ .

**Remark 3.2:** If the functions  $f, g$  and  $I_i$  are compact or Lipschitz continuous (see e.g [5, 7]), then  $(H_3)$ – $(H_7)$  is automatically satisfied.

In some of the early related results in references and above results, it is supposed that the map  $h$  is uniformly bounded. In fact, if  $h$  is compact, then it must be bounded on bounded set. Here we give an existence result under growth condition of  $f, g$  and  $I_i$ , when  $h$  is not uniformly bounded. Precisely, we replace the assumptions  $(H_3)$ – $(H_6)$  by

(H8) There exists a function  $p \in L([0, b], R^+)$  and a increasing function  $\varphi: R^+ \rightarrow R^+$  such that

$$\|f(t, x)\| \leq p(t)\varphi(\|x\|)$$

for a.e  $t \in [0, b]$ , for all  $x \in PC([0, b], X)$ .

(H9) There exist two functions  $L_g \in L([0, b], R^+)$  and  $\widehat{L}_g \in L([0, b], R^+)$  and a increasing function  $\Psi: R^+ \rightarrow R^+$  such that

$$\|g(t, s, x)\| \leq L_g(t)\widehat{L}_g(s)\Psi(\|x\|),$$

for a.e  $t \in [0, b]$  and for all  $L_g \in PC([0, b], X)$ . Assume that the finite bound of  $\int_0^t L_g(s)ds$  is  $G_3$ .

(H10) There exists a function  $L_e \in L([0, b], R^+)$  and a increasing function  $\Gamma: R^+ \rightarrow R^+$  such that

$$\|e(t, x, x_1)\| \leq L_e(t)\Gamma(\|x\| + \|x_1\|)$$

for a.e  $t \in [0, b]$  and for all  $L_g \in PC([0, b], X)$ . Assume that the finite bound of  $\int_0^t L_e(s)ds$  is  $G_5$ .

(H11) There exist two functions  $L_k \in L([0, b], \mathbb{R}^+)$  and  $\widehat{L}_k \in L([0, b], \mathbb{R}^+)$  and an increasing function  $\Theta: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that

$$\|k(t, s, x)\| \leq L_k(t) L_k(s) \Theta(\|x\|),$$

for a.e.  $t \in [0, b]$  and for all  $L_k \in PC([0, b], X)$ . Assume that the finite bound of  $\int_0^t L_k(s) ds$  is  $G_4$ .

**Theorem 3.2:** Suppose that the assumptions  $(H_1) - (H_2)$  and  $(H_8) - (H_{11})$  are satisfied, then the equation (1) - (3) has at least one mild solution if

$$\begin{aligned} & \limsup_{r \rightarrow \infty} \frac{1}{r} \{M_0[\varphi(r) + L_e(t)] + L_e(t)(\Gamma\|x\|) \\ & + G_3 L_k(t) \Theta(r) + \eta(t) M_0[G_4 \Gamma(r) + G_3 \Theta(r) \int_0^t \widehat{L}_k(s) ds] \\ & + M_0[\phi(r) \int_0^t L_f(s) ds + G_2 \Psi(r) \int_0^t \widehat{L}_g(s) ds + \sum_{k=1}^n d_k]\} < 1, \end{aligned} \tag{10}$$

where  $\varphi(r) = \sup\{\|h(x)\|, \|x\| \leq r\}$ .

**Proof:** The inequality (10) implies that there exist a constant  $r > 0$  such that

$$\begin{aligned} & M_0[\varphi(r) + L_e(t)] + L_e(t)(\Gamma P x P) + G_3 L_k(t) \Theta(r) + \eta(t) M_0[G_4 \Gamma(r) \\ & + G_3 \Theta(r) \int_0^t \widehat{L}_k(s) ds] + M_0[\phi(r) \int_0^t p(s) ds + G_2 \Psi(r) \int_0^t \widehat{L}_g(s) ds + \sum_{k=1}^n d_k] < r, \end{aligned}$$

As in the proof of Theorem 3.1, let  $W_0 = \{x \in PC([0, b], X), \|x(t)\| \leq r\}$  and  $W_1 = \overline{\text{con } FW_0}$ . Then for any  $x \in W_1$ , we have  $W_1 = \overline{\text{con } FW_0}$ . Then for any  $x \in W_1$ , we have

$$\begin{aligned} & \|x(t)\| \leq \|U_x(t, 0)h(x)\| + \|U_x(t, 0)e(0, x(0), 0)\| + \left\| e\left(t, x(t), \int_0^t k(t, s, x(s)) ds\right) \right\| \\ & + \int_0^t \left\| A(t, x(t)) U_x(t, s) e\left(t, x(t), \int_0^s k(s, \tau, x(\tau)) d\tau\right) \right\| ds \\ & + \int_0^t \left\| U_x(t, s) [f(s, x(s)) + \int_0^s g(s, \tau, x(s)) d\tau] \right\| ds + \sum_{0 < t_k < t} \|U_x(t, t_k) I_i(x(t_k))\| \\ & \leq M_0[\varphi(r) + L_e(t)] + L_e(t)(\Gamma\|x\|) + \int_0^t L_k(t) L_k(s) \Theta(\|x\|) ds \\ & + \eta(t) M_0 \int_0^t [p(s) \Gamma(\|x\|) + \int_0^s L_k(s) L_k(\tau) \Theta(\|x\|) d\tau] ds \\ & + M_0 \left[ \int_0^t L_f(s) \phi(\|x(s)\|) ds + \int_0^t \int_0^s L_g(s) L_g(\tau) \Psi(\|x(\tau)\|) d\tau ds + \sum_{k=1}^n d_k \right] \\ & \leq M_0[\varphi(r) + L_e(t)] + L_e(t)(\Gamma\|x\|) + G_3 L_k(t) \Theta(\|x\|) \\ & + \eta(t) M_0 [G_4 \Gamma(\|x\|) + G_3 \int_0^t \widehat{L}_k(s) \Theta(\|x\|) ds] \\ & + M_0 \left[ \int_0^t p(s) \phi(\|x(s)\|) ds + G_2 \int_0^t \widehat{L}_g(s) \Psi(\|x(s)\|) ds + \sum_{k=1}^n d_k \right] \end{aligned}$$



$$\begin{aligned} \|x(t)\| &\leq M_0[\varphi(r) + L_e(t)] + L_e(t)(\Gamma P x P) + G_3 L_k(t)\Theta(r) \\ &\quad + \eta(t)M_0[G_4\Gamma(r) + G_3\Theta(r)] \int_0^t \widehat{L_k}(s) ds \\ &\quad + M_0[\phi(r) \int_0^t p(s) ds + G_2\Psi(r) \int_0^t \widehat{L_g}(s) ds + \sum_{k=1}^n d_k] \\ &< r, \end{aligned}$$

for  $t \in [0, b]$ . It means that  $W_1 \subset W_0$ . So we can complete the proof similarly to Theorem 3.1.

#### 4. WHEN $h$ IS LIPSCHITZ

In this section, we discuss the equation (1)–(3) when  $h$  is Lipschitz and  $f, g$  and  $I_k$  are not Lipschitz. Assume that

(H12) The function  $h$  is a Lipschitz continuous in  $X$ , there exists a constant  $L_0 > 0$  such that

$$\|h(x) - h(y)\| \leq L_0 \|x - y\|, \quad x, y \in PC([0, b], X).$$

**Theorem 4.1:** Suppose that the assumptions  $(H_1) - (H_{12})$  are satisfied, then the equation (1)–(3) has at least one mild solution provided that

$$\begin{aligned} &M_0[L_0 + h_4(t)]\chi_c(B) + 2[m_e(t) + m_k(t)G_2 + M_0\eta(t)G_1] \\ &\quad + 2M_0 \left[ \int_0^t \{2G_2 m_k(s) + m_f(s) + 2G_0 n_g(s)\} ds + \sum_{k=1}^n l_k \right] < 1. \end{aligned} \tag{11}$$

**Proof:** Consider the map  $F: PC([0, b], X) \rightarrow PC([0, b], X)$  is defined by  $F = F_1 + F_2$ , where

$$\begin{aligned} (F_1 x)(t) &= U_x(t, 0)h(x) + U_x(t, 0)e(0, x(0), 0), \\ (F_2 u)(t) &= \int_0^t A(t, x(t))U_x(t, s)e\left(t, x(t), \int_0^s k(s, \tau, x(\tau))d\tau\right) ds - e\left(t, x(t), \int_0^t k(t, s, x(s))ds\right) \\ &\quad + \int_0^t U_x(t, s)[f(s, x(s)) + \int_0^s g(s, \tau, x(\tau))d\tau] ds + \sum_{0 < t_k < t} U_x(t, t_k)I_k(x(t_k)), \end{aligned}$$

for  $x \in PC([0, b], X)$ . As defined in the proof of Theorem 3.1. We define  $W_0 = \{x \in PC([0, b], X) : \|x(t)\| \leq m(t), \text{ for all } t \in [0, b] \text{ and let } W = \overline{\text{con}FW_0}$ . Then from the proof of Theorem 3.1 we know that  $W$  is a bounded closed convex and equicontinuous subset of  $PC([0, b], X)$  and  $FW \subset W$ . We shall prove that  $F$  is  $\chi_c$  contraction on  $W$ . Then Darbo-Sadovskii's fixed point theorem can be used to get a fixed point of  $F$  in  $W$ , which is a mild solution of (1)–(2). First, for every bounded subset  $B \subset W$ , from  $(H_{12})$  and Lemma 2.1 we have

$$\begin{aligned} \chi_c(F_1 B) &= \chi_c(U_B(t, 0)h(B)) + U_B(t, 0)e(0, B(0), 0) \\ &\leq M_0 \chi_c[h(B)] + e(B, 0) \\ &\leq M_0[L_0 + h_4(t)] \chi_c(B) \end{aligned} \tag{12}$$

Next, for every bounded subset  $B \subset W$ , for  $t \in [0, b]$  and every  $\varepsilon > 0$ , there is a sequence  $\{x_k\}_{k=1}^\infty \subset B$  such that

$$\chi(F_2(B(t))) \leq 2\chi(\{F_2 x_i(t)\}_{i=1}^\infty) + \varepsilon.$$

Note that  $B$  and  $F_2B$  are equicontinuous, we can get from Lemma 2.1, Lemma 2.4, Lemma 2.5 and using the assumptions we get

$$\begin{aligned} \chi(F_2B(t)) &\leq 2\chi(e(t, \{x_k(s)\}_{k=1}^\infty, \int_0^t k(t, s, \{x_k(\tau)\}_{k=1}^\infty) ds)) \\ &\quad + 2M_0\eta(t) \int_0^t \chi(e(s, \{x_k(s)\}_{k=1}^\infty, \int_0^s k(s, \tau, \{x_k(\tau)\}_{k=1}^\infty) d\tau)) ds \\ &\quad + 2M_0 \int_0^t \chi(f(s, \{x_k(s)\}_{k=1}^\infty)) ds + 4M_0 \int_0^t \int_0^s \chi(g(s, \tau, \{u_k(\tau)\}_{k=1}^\infty)) d\tau ds \\ &\quad + 2M_0 \sum_{i=1}^n \chi(I_i(\{u_k(t_k)\}_{k=1}^\infty)) + \varepsilon \\ &\leq 2m_e(t) \chi\{x_k(t)\}_{k=1}^\infty + 2m_k(t) \int_0^t m_k(s) \chi\{x_k(s)\}_{k=1}^\infty ds \\ &\quad + 2M_0\eta(t) [\int_0^t m_e(s) \chi\{x_k(s)\}_{k=1}^\infty ds + 2 \int_0^t \int_0^s m_k(s) m_k(\tau) \chi\{x_k(\tau)\}_{k=1}^\infty d\tau ds] \\ &\quad + 2M_0 \int_0^t m_f(s) \chi(\{u_k(s)\}_{k=1}^\infty) ds + 4M_0 \int_0^t \int_0^s m_g(s) n_g(\tau) \chi(\{u_k(\tau)\}_{k=1}^\infty) d\tau ds \\ &\quad + 2M_0 \sum_{i=1}^n l_i \chi(\{u_k(t_k)\}_{k=1}^\infty) + \varepsilon \\ &\leq 2[m_e(t) + m_k(t)G_2 + M_0\eta(t)G_1] \chi(B) \\ &\quad + 2M_0 [\int_0^t \{2G_2 m_k(s) + m_f(s)\} \chi(B) ds + 2G_0 \int_0^t n_g(s) \chi(B) ds] \\ &\quad + 2M_0 \sum_{k=1}^n l_k \chi(B) + \varepsilon. \end{aligned}$$

Since  $\varepsilon > 0$  is arbitrary, it follows that from the above inequality that

$$\begin{aligned} \chi_c(F_2B(t)) &\leq 2[m_e(t) + m_k(t)G_2 + M_0\eta(t)G_1] \chi_c(B) \\ &\quad + 2M_0 [\int_0^t \{2G_2 m_k(s) + m_f(s) + 2G_0 n_g(s)\} ds + \sum_{k=1}^n l_k] \chi_c(B) \end{aligned} \tag{13}$$

for any bounded  $B \subset W$ .

Now, for any subset  $B \subset W$ , due to Lemma 2.1, (12) and (13) we have

$$\begin{aligned} \chi_c(FB) &\leq \chi_c(F_1B) + \chi_c(F_2B) \\ &\leq M_0[L_0 + h_4(t)] \chi_c(B) + 2[m_e(t) + m_k(t)G_2 + M_0\eta(t)G_1] \chi_c(B) \\ &\quad + 2M_0 [\int_0^t \{2G_2 m_k(s) + m_f(s) + 2G_0 n_g(s)\} ds + \sum_{k=1}^n l_k] \chi_c(B) \end{aligned} \tag{14}$$

By (14) we know that  $F$  is a  $\chi_c$ -contraction on  $W$ . By Lemma 2.2, there is a fixed point  $x$  of  $F$  in  $W$ , which is a solution of (1)–(3). This completes the proof.

**Theorem 4.2:** Suppose that the assumptions  $(H_1)–(H_{12})$  are satisfied, then the equation (1)–(3) has at least one mild solution if (15) and the following condition are satisfied.

$$\begin{aligned} &M_0L_0 + \limsup_{r \rightarrow \infty} \frac{1}{r} \{M_0L_e(t) + L_e(t)\Gamma(r) + G_3L_k(t)\Theta(r) \\ &\quad + \eta(t)M_0[G_4\Gamma(r) + G_3\Theta(r)] \int_0^t \widehat{L_k(s)} ds\} \\ &\quad + M_0[\phi(r) \int_0^t L_f(s) ds + G_2\Psi(r) \int_0^t \widehat{L_g(s)} ds + \sum_{k=1}^n d_k] < 1. \end{aligned} \tag{15}$$

**Proof:** From the equation (15) and fact that  $L_0 < 1$ , there exists a constant  $r > 0$  such that

$$\begin{aligned} & M_0(rL_0 + Ph(0)P + L_e(t)) + L_e(t)(\Gamma PxP) + G_3L_k(t)\Theta(r) \\ & + \eta(t)M_0[G_4\Gamma(r) + G_3\Theta(r)] \int_0^t \widehat{L_k(s)} ds \\ & + M_0[\phi(r) \int_0^t L_f(s) ds + G_2\Psi(r) \int_0^t \widehat{L_g(s)} ds + \sum_{k=1}^n d_k] < r. \end{aligned}$$

We define  $W_0 = \{x \in PC([0, b], X), Px(t)P \leq r, \text{ for all } t \in [0, b]\}$ . Then for every  $x \in W_0$ , we have

$$\begin{aligned} \|Fx(t)\| & \leq \|U_x(t, 0)h(x)\| + \|U_x(t, 0)e(0, x(0), 0)\| + \left\| e\left(t, x(t), \int_0^t k(t, s, x(s)) ds\right) \right\| \\ & + \int_0^t \left\| A(t, x(t))U_x(t, s)e\left(t, x(t), \int_0^s k(s, \tau, x(\tau)) d\tau\right) \right\| ds \\ & + \int_0^t \left\| U_x(t, s)[f(s, x(s)) + \int_0^s g(s, \tau, x(s)) d\tau] \right\| ds + \sum_{0 < t_k < t} \|U_x(t, t_k)I_i(x(t_k))\| \\ & \leq M_0(rL_0 + \|h(0)\| + L_e(t)) + L_e(t)(\Gamma\|x\|) + G_3L_k(t)\Theta(r) \\ & + \eta(t)M_0[G_4\Gamma(r) + G_3\Theta(r)] \int_0^t \widehat{L_k(s)} ds \\ & + M_0[\phi(r) \int_0^t L_f(s) ds + G_2\Psi(r) \int_0^t \widehat{L_g(s)} ds + \sum_{k=1}^n d_k] < r. \end{aligned}$$

for  $t \in [0, b]$ . This means that  $FW_0 \subset W_0$ . Define  $W = \overline{con FW_0}$ . The above proof also implies that  $FW \subset W$ . So we can prove the theorem similar with Theorem 4.1 and hence we omit it.

### 5. APPLICATION

The notion of controllability is of great importance in mathematical control theory. Many fundamental problems of control theory such as pole-assignment, stabilizability and optimal control may be solved under the assumption that the system is controllable. It means that it is possible to steer any initial state of the system to any final state in some finite time using an admissible control. During the last few decades, several authors [25, 27] have discussed the existence, uniqueness, and asymptotic behavior of the solution of these systems. Apart from these, the study of controllability and observability properties of a system in control theory is certainly, at present, one of the most active interdisciplinary areas of research. Control theory arises in most modern applications. On the other hand, control theory has remained a discipline where many mathematical ideas and methods have fused to produce a new body of important mathematics. As an application of Theorem 3.1 we shall consider the system (1)–(3) with a control parameter such as

$$\begin{aligned} \frac{d}{dt}[x(t) + e(t, x(t), \int_0^t k(t, s, x(s)) ds)] + A(t, x(t))x(t) \\ = f(t, x(t)) + Cv(t) + \int_0^t g(t, s, x(s)) ds, \quad t \in [0, b], t \neq t_i, \end{aligned} \tag{16}$$

$$x(0) + h(x) = x_0, \tag{17}$$

$$\Delta x(t_k) = I_k(x(t_k)), \quad k = 1, 2, 3, \dots, n, \tag{18}$$

where  $A, f, g, h$  and  $I_k$  are as before and  $C$  is a bounded linear operator from a Banach space  $V$  into  $X$  and the control function  $v \in L^2(J, V)$ . The mild solution of (16)–(18) is given by

$$\begin{aligned} x(t) & = U_x(t, 0)x_0 - U_x(t, 0)h(x) + U_x(t, 0)e(0, x(0), 0) - e\left(t, x(t), \int_0^t k(t, s, x(s)) ds\right) \\ & + \int_0^t A(t, x(t))U_x(t, s)e\left(t, x(t), \int_0^s k(s, \tau, x(\tau)) d\tau\right) ds \\ & + \int_0^t U_x(t, s)[f(s, x(s)) + Cv(s) + \int_0^s g(s, \tau, x(\tau)) d\tau] ds \\ & + \sum_{0 < t_k < t} U_x(t, t_k)I_k(x(t_k)), \quad 0 \leq t \leq b. \end{aligned}$$

**Definition 5.1:** ([26, 27]) System (16)–(18) is said to be controllable on the interval  $J$ , if for every  $x_0, x_1 \in X$ , there exists a control  $v \in L^2(J, V)$  such that the mild solution  $u(\cdot)$  of (16)–(18) satisfies  $x(0) = x_0$  and  $x(b) = x_1$ .

To study the controllability result we need the following additional condition:

1. The linear operator  $W: L^2(J, V) \rightarrow X$ , defined by

$$Wv = \int_0^b U_x(b, s)Cv(s)ds$$

induces an inverse operator  $W^{-1}$  defined on  $L^2(J, V)/\ker W$  and there exists a positive constant  $M_1 > 0$  such that  $\|CW^{-1}\| \leq M_1$ .

**Theorem 5.1:** If the assumptions  $(H_1)–(H_{13})$  are satisfied, then the system (16)–(18) is controllable on  $J$ .

**Proof.** Using the assumption  $(H_{13})$ , for an arbitrary function  $u(\cdot)$ , define the control

$$\begin{aligned} v(t) = & W^{-1}[u_1 - U_x(b, 0)x_0 + U_x(b, 0)h(x) - U_x(b, 0)e(0, x(0), 0) \\ & + e\left(b, x(b), \int_0^b k(b, s, x(s))ds\right) - \int_0^b A(b, x(b))U_x(b, s)e\left(b, x(b), \int_0^s k(s, \tau, x(\tau))d\tau\right)ds \\ & - \int_0^b U_x(b, s)[f(s, x(s)) - \int_0^s g(s, \tau, x(\tau))d\tau]ds - \sum_{0 < t_k < t} U_x(t, t_k)I_k(x(t_k))](t). \end{aligned}$$

We shall show that when using this control, the operator  $H: Z \rightarrow Z$  defined by

$$\begin{aligned} (Hv)(t) = & U_x(t, 0)x_0 - U_x(t, 0)h(x) + U_x(t, 0)e(0, x(0), 0) - e\left(t, x(t), \int_0^t k(t, s, x(s))ds\right) \\ & + \int_0^t A(t, x(t))U_x(t, s)e\left(t, x(t), \int_0^s k(s, \tau, x(\tau))d\tau\right)ds \\ & + \int_0^t U_x(t, s)[f(s, x(s)) + CW^{-1}[u_1 - U_x(b, 0)x_0 + U_x(b, 0)h(x) \\ & - U_x(b, 0)e(0, x(0), 0) + e\left(b, x(b), \int_0^b k(b, s, x(s))ds\right) \\ & + \int_0^b A(b, x(b))U_x(b, s)e\left(b, x(b), \int_0^s k(s, \tau, x(\tau))d\tau\right)ds \\ & - \int_0^b U_x(b, s)[f(s, x(s)) + \int_0^s g(s, \tau, x(\tau))d\tau]ds \\ & - \sum_{0 < t_k < t} U_x(t, t_k)I_k(x(t_k))](s) + \int_0^s g(s, \tau, x(\tau))d\tau]ds + \sum_{0 < t_k < t} U_x(t, t_k)I_k(x(t_k)). \end{aligned}$$

has a fixed point. This fixed point is, then a solution of (16)–(18). Clearly,  $(Hv)(b) = x(b) = x_1$ , which means that the control  $v$  steers the system (16)–(18) from the initial state  $x_0$  to the final state  $x_1$  at time  $b$ , provided we can obtain a fixed point of the nonlinear operator  $H$ . The remaining part of the proof is similar to Theorem 3.1 and hence, it is omitted.

## REFERENCES

- [1] K. Balachandran and E. R. Anandhi, Neutral functional integrodifferential control systems in Banach spaces, *Kybernetika*, 39 (2003), 359-367.
- [2] K. Balachandran and E. R. Anandhi, Controllability of neutral functional integrodifferential infinite delay systems in Banach spaces, *Taiwanese Journal of Mathematics*, 8 (2004), 689-702.

- [3] E. Hernandez and H. R. Henriquez, Impulsive partial neutral differential equations, *Applied Mathematics Letters*, 19 (2006), 215-222.
- [4] J. K. Hale and S. M. Verduyn Lunel, *Introduction to Functional-Differential Equations*, Springer-Verlag, New York, 1993.
- [5] L. Byszewski, *Theorems about the existence and uniqueness of solutions of a semilinear evolution nonlocal Cauchy problem*, J. Math. Anal. Appl. 162 (1992), 494-505.
- [6] L. Byszewski, H. Akca, *Existence of solutions of a semilinear functional-differential evolution nonlocal problem*, Nonlinear Anal. 34 (1998), 65-72.
- [7] L. Byszewski, V. Lakshmikanthan, *Theorems about the existence and uniqueness of solutions of a nonlocal Cauchy problem in Banach spaces*, Applicable Anal. 40 (1990), 11-19.
- [8] S. K. Ntouyas, P. Ch. Tsamatos, *Global existence for semilinear evolution equations with nonlocal conditions*, J. Math. Anal. Appl. 210 (1997), 679-687.
- [9] Q. Dong, G. Li, J. Zhang, *Quasilinear nonlocal integrodifferential equations in Banach spaces*, Electronic J. Diff. Equa. 19 (2008), 1-8.
- [10] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer, New York, 1983.
- [11] D. Guo, X. Liu, *External solutions of nonlinear impulsive integrodifferential equations in Banach spaces*, J. Math. Anal. Appl. 177 (1993), 538-552.
- [12] J. Liang, J. H. Liu, T. J. Xiao, *Nonlocal impulsive problems for nonlinear differential equations in Banach spaces*, Math. Computer Modelling. 49 (2009), 798-804.
- [13] A. M. Samoilenko, N. A. Perestyuk, *Impulsive Differential Equations*, World Scientific, Singapore, 1995.
- [14] Z. Luo, J. J. Nieto, *New results for the periodic boundary value problem for impulsive integrodifferential equations*, Nonlinear Anal. 70 (2009), 2248-2260.
- [15] J. Banas, K. Goebel, *Measure of noncompactness in Banach spaces*, Lecture Notes in Pure and Applied Mathematics, vol. 60, Marcle Dekker, New York, 1980.
- [16] F.S. De Blasi, *On a property of the unit sphere in a Banach space*, Bull. Math. Soc. Sci. Math. R.S. Roumanie, 21 (1977), 259-262.
- [17] J. M. Ayerbe, T. D. Benavides, G. L. Acedo, *Measure of noncompactness in in metric fixed point theorem*, Birkhauser, Basel, 1997.
- [18] J. Banas, W.G. El-Sayed, *Measures of noncompactness and solvability of an integral equation in the class of functions of locally bounded variation*, J. Math. Anal. Appl. 167 (1992), 133-151.
- [19] G. Emmanuele, *Measures of weak noncompactness and fixed point theorems*, Bull. Math. Soc. Sci. Math. R. S. Roumanie, 25 (1981), 353-358.
- [20] Q. Dong, G. Li, *The Existence of solutions for semilinear differential equations with nonlocal conditions in Banach spaces*, Electronic J. Qualitative Theory of Diff. Equa. 47 (2009), 1-13.
- [21] Z. Fan, Q. Dong, G. Li, *Semilinear differential equations with nonlocal conditions in Banach spaces*, International J. Nonlinear Sci. 2 (2006), 131-139.
- [22] H.P. Heinz, *On the behavior of measure of noncompactness with respect to differentiation and integration of vector-valued functions*, Nonlinear Anal. 7 (1983), 1351-1371.
- [23] M. Kamenskii, V. Obukhovskii, P. Zecca, *Condensing multivalued maps and semilinear differential inclusions in Banach spaces*, De Gruyter Series. Nonlinear Anal. Appl. Vol.7, de Gruyter, Berlin, 2001.
- [24] D. Both, *Multivalued perturbation of m-accretive differential inclusions*, Israel J. Math. 108 (1998), 109- 138.

[25] K. Balachandran, J. Y. Park, *Existence of solutions and controllability of nonlinear integrodifferential systems in Banach spaces*, Mathematical Problems in Engineering, 2 (2003), 65-79.

[26] B. Radhakrishnan, K. Balachandran, *Controllability results for semilinear impulsive integrodifferential evolution systems with nonlocal conditions*, J. Control Theory and Appl. 10 (2012), 28-34.

[27] L. Wang and Z. Wang, *Controllability of abstract neutral functional differential systems with infinite delay*, Dynamics of Continuous, Discrete and Impulsive Systems Ser.B: Applications and Algorithms, 9 (2002), 59-70.

**Source of support: Nil, Conflict of interest: None Declared**

***[Copy right © 2014 This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]***