

**A PENALTY FUNCTION APPROACH  
FOR SOLVING INEQUALITY CONSTRAINED OPTIMIZATION PROBLEMS**

**M. H. Farag\*<sup>1, 2</sup>, W. A. Hashem<sup>2</sup> and H. H. Saleh<sup>2</sup>**

<sup>1</sup>*Mathematics Department, Faculty of Science, Taif University, Hawia (888), Taif, Saudi Arabia.*

<sup>2</sup>*Mathematics Department, Faculty of Science, Minia University, Mina, Egypt.*

*(Received On: 08-04-14; Revised & Accepted On: 31-07-14)*

**ABSTRACT**

*In this paper, a computational approach based on an exterior penalty function MPQI method is given for solving a class of continuous inequality constrained optimization problems. The essential steps of the partial quadratic interpolation technique and its modified are given. The numerical algorithm and the flowchart of the penalty function method combined with the Modified Partial Quadratic Interpolation Technique are given. For illustration, three examples are solved using the proposed method. From the solutions obtained, we observe that the values of their object functions are amongst the smallest when compared with those obtained by other existing methods available in the literature. More importantly, our method finds solution which satisfies the continuous inequality constraints.*

**Keywords:** *Constrained optimization, Exterior penalty function, modified partial quadratic interpolation technique.*

**1. INTRODUCTION**

In recent years, there has been a resurgence of interest in penalty methods [1-5] because of their ability to handle degenerate problems and inconsistent constraint linearization. Exact penalty methods have been used successfully to solve mathematical programs with complementarity constraints (MPCCs) [6-7], a class of problems that do not satisfy the Mangasarian-Fromovitz constraint qualification at any feasible point. They are also used in nonlinear programming algorithms to ensure the feasibility of subproblems and to improve the robustness of the iteration [8-9].

Penalty methods have undergone three stages of development since their introduction in the 1950s. They were first seen as vehicles for solving constrained optimization problems by means of unconstrained optimization techniques. This approach has not proved to be effective, except for special classes of applications. In the second stage, the penalty problem is replaced by a sequence of linearly constrained subproblems. These formulations, which are related to the sequential quadratic programming approach, are much more effective than the unconstrained approach but they leave open the question of how to choose the penalty parameter. In the most recent stage of development, penalty methods adjust the penalty parameter at every iteration so as to achieve a prescribed level of linear feasibility. The choice of the penalty parameter then ceases to be a heuristic and becomes an integral part of the step computation.

The general form of a minimization problem with inequality and equality constraints is as follows [10]: Find

$x \in \mathcal{R}^n$  such that minimize  $f(x)$  subject to :

$$g_i(x) \leq 0, \quad i = 1, \dots, m, \tag{1}$$

and

$$h_j(x) = 0, \quad j = 1, \dots, p. \tag{2}$$

Penalty function methods are the most popular constraint handling methods among users. Two main branches of penalty methods have been proposed in the literature: Exterior and Interior which is also called the barrier method.

**Corresponding author: M. H. Farag\*<sup>1, 2</sup>**

<sup>1</sup>*Mathematics Department, Faculty of Science, Taif University, Hawia (888), Taif, Saudi Arabia.*

The widely used form of  $\Psi(x, r_k)$  in the exterior penalty method [11] is:

$$\Psi = \Psi(x, r_k) = f(x) + r_k \left\{ \sum \langle g_i(x) \rangle^2 + \sum (h_j(x))^2 \right\}, \quad r_k \rightarrow \infty \quad (3)$$

where

$$\langle g_i(x) \rangle = \max \{0, g_i(x)\}, \quad i = 1, \dots, m, \quad (4)$$

and  $r_k$  is a parameter, which is modified at the beginning of each round of optimization. Each optimization round is defined here as a complete optimization of  $\Psi(x, r_k)$  for a fixed value of  $r_k$  until the convergence is achieved. The optimum point  $x^*$ , at the end of each round serves as the starting point,  $x_1$  of the next round of optimization with a larger  $r_k$ .

The original form of the interior penalty function  $\Psi_{in}(x, r_k)$  is as follows [12-13]:

$$\Psi_{in}(x, r_k) = f(x) + \left\{ \sum \frac{1}{g_i(x)} + \sum \frac{1}{h_j(x)} \right\}, \quad r_k \rightarrow 0, \quad (5)$$

where  $r_k$  reduces from a high value to zero gradually. Rao [6, 7] has proposed the following function for selecting  $r_k$  at the start of the optimization procedure:

$$r_k = (0.1 \sim 1) \times \frac{f(x_1)}{-\sum \frac{1}{g(x_1)}}, \quad (6)$$

where  $x_1$  is the initial point in the feasible region. The optimization procedure is similar to the exterior penalty function method except that  $r_k$  reduces to 0 gradually. Here, the reduction follows [8]

$$r_{k+1} = \lambda \times r_k \quad (7)$$

where  $\lambda$  is a coefficient less than 1.

In this paper, a computational approach based on an exterior penalty function MPQI method is given for solving a class of continuous inequality constrained optimization problems. The essential steps of the partial quadratic interpolation technique and its modified are given. The numerical algorithm and the flowchart of the penalty function method combined with the Modified Partial Quadratic Interpolation Technique are given. For illustration, three examples are solved using the proposed method. From the solutions obtained, we observe that the values of their object functions are amongst the smallest when compared with those obtained by other existing methods available in the literature. More importantly, our method finds solution which satisfies the continuous inequality constraints.

## 2. PARTIAL QUADRATIC INTERPOLATION TECHNIQUE (PQI)

The essential steps to apply the partial quadratic interpolation technique [14] can be summarized as follows:

- 1) Choose some starting point  $\xi^0 \in \mathfrak{R}_m$  and  $\nu = \mathbf{1}$ .
- 2) Approximate the function  $F(x)$  about  $\xi^r$  in the quadratic form

$$F(x) = a + [B_m(x^r)]^{Tr} [x - x^r] + \frac{1}{2} [x - x^r]^{Tr} [A_m(x^r)] [x - x^r] \quad (8)$$

where  $[A_m]$  and  $[B_m]$  represent the gradient vector and the Hessian matrix of the function  $F(x)$  respectively. To compute particular values for  $a, A_m, B_m$  we choose a set of interpolation points as follows:

- i)  $m$  points  $[x_{i+}^r], i = 1(1)m$

$$[x_{i+}^r] = (x_1^r, x_2^r, \dots, x_{i-1}^r, x_i^r + l_i, x_{i+1}^r, \dots, x_m^r) \quad (9)$$

- ii)  $m$  points  $[x_{i-}^r], i = 1(1)m$

$$[x_{i-}^r] = (x_1^r, x_2^r, \dots, x_{i-1}^r, x_i^r - l_i, x_{i+1}^r, \dots, x_m^r) \quad (10)$$

$$\text{iii) } \frac{m(m-1)}{2} \text{ points } [x_{ij}^r], i = 1(1)m-1, j = i+1(1)m \text{ where}$$

$$[x_{ij}^r] = (x_1^r, x_2^r, \dots, x_{i-1}^r, x_i^r + l_i, x_{i+1}^r, \dots, x_j^r + l_j, \dots, x_m^r) \quad (11)$$

Using these interpolation points it can be shown that

$$a = F(x^r) \quad (12)$$

and the elements  $b_i, a_{ij}$ , of  $B_m, A_m$  respectively are given by

$$b_i = \frac{F(x_{i+}^r) - F(x_{i-}^r)}{2 l_i}, \quad a_{ii} = \frac{F(x_{i+}^r) - 2F(x^r) + F(x_{i-}^r)}{l_i^2} \quad (13)$$

$$a_{ij} = \frac{F(x_{ij}^r) - F(x_{i+}^r) - F(x_{j+}^r) + F(x^r)}{l_i l_j} \quad (14)$$

$$a_{ij} = \frac{F(x_{ij}^r) - F(x_{i+}^r) - F(x_{j+}^r) + F(x^r)}{l_i l_j} \quad (15)$$

The  $l_i$  are a set of constants which determine the accuracy of the interpolation.

3) Extract the symmetric positive definite matrix  $[A_q]$  from the symmetric matrix  $[A_m]$  using Choliski's method,  $q \leq m$ , by cancelling certain rows and columns. Essentially we write  $A_m = [S_m][S_m]^{Tr}$ . From this we have

$$s_{11}^2 = a_{11}$$

If  $a_{11} \leq 0$  then we eliminate the first row and column in each of  $[A_m], [S_m]$  and  $[S_m]^{Tr}$  and perform the calculation on the  $[A_{m-1}], [S_{m-1}]$  and  $[S_{m-1}]^{Tr}$ . If  $a_{11} \geq 0$  then we have

$$s_{11}^2 = \sqrt{a_{11}}, \quad s_{1j} = \frac{a_{1j}}{s_{11}}, \quad j = 2(1)m. \quad (16)$$

Let us now suppose that we have operated on the first  $j-1$  columns of  $[S]$ , i.e. We have either calculated the elements or eliminated them. The operation on the  $j^{th}$  column gives

$$s_{jj}^2 = a_{jj} - \sum_{i \in k1, i \leq j} s_{ij}^2. \quad (17)$$

where  $k1$  is the set of indices of rows and columns not eliminated.

If  $s_{jj} \leq 0$  we eliminate the  $j^{th}$  columns and rows  $[A_q]$  and  $[S_q]$  where  $[A_q]$  and  $[S_q]$  are the current reduced matrices derived to date from  $[A_m]$  and  $[S_m]$ ,  $q \leq m$ .

Otherwise we take

$$s_{jj} = \sqrt{a_{jj} - \sum_{i \in k1, i \leq j} s_{ij}^2}, \quad s_{ij} = \frac{[a_{ij} - \sum_{i \in k1, i \leq j} s_{ij}^2]}{s_{jj}} \quad (18)$$

This process is repeated for each column until we finally obtain the reduced matrix  $[A_q]$  given by

$$[A_q] = [S_q][S_q]^{Tr} \quad (19)$$

4) Solve the system of the linear equations

$$[A_q][\Delta x_i] = [B_q] \quad (20)$$

where  $B_q$  is the reduced form of gradient vector corresponding to  $A_q$ .

5) Compute a new point  $x^{v+1}$  from

$$x_i^{v+1} = \begin{cases} x_i^{v+1} + \beta \Delta x_i & \text{for } x_i \in \mathfrak{R}_q \\ x_i^v & \text{for } x_i \in \mathfrak{R}_q \end{cases}$$

where  $\beta$  is a parameter which takes values  $1, \frac{1}{2}, \frac{1}{4}, \dots$  and we use the first value of  $\beta$  which satisfies

$F(x^{v+1}) < F(x^v)$ . If  $\beta$  becomes too small without satisfying this condition, the calculation can be restarted with a finer approximation of the matrices  $[A_m]$  and  $[B_m]$ , i.e. smaller values  $l_i$ .

### 3. MODIFIED PQI TECHNIQUE (MPQI) [15]

In PQI technique we set  $t = 1, \frac{1}{2}, \frac{1}{4}, \dots$  and we take the first value of  $t$  which satisfies the condition

$$f(x^{r+1}) < f(x^r), \quad x^r \in \mathfrak{R}_n.$$

However, the value of  $t$  taken by this way may not be the optimal value of  $t$ . Since there is a great possibility that the optimal value  $t^*$  lies between these values, i.e. between  $t = 1, \frac{1}{2}, \frac{1}{4}, \dots$  to get the optimal step size  $t^*$  we suggest the following modification:

Let us approximate  $f(t) = f(x^r + t \delta x^r)$  by a polynomial of second degree  $p_2(t)$  over the interval  $[0,1]$  as following:

$$f(t) = p_2(t) = \begin{bmatrix} 1 & \frac{t}{h} & \left(\frac{t}{h}\right)^2 \end{bmatrix} [L_2] \begin{pmatrix} f_0 \\ f_1 \\ f_2 \end{pmatrix} \quad (21)$$

where  $h$  is the interval of the interpolation,  $f_{2i} = f(u^r + i \delta u^r)$ ,  $i = 0, \frac{1}{2}, 1$  and  $L_2$  is the Lagrange matrix where

$$L_2 = \frac{1}{2} \begin{pmatrix} 2 & 0 & 0 \\ -3 & 4 & -1 \\ 1 & -2 & 1 \end{pmatrix} \quad (22)$$

from (21) and (22) we have

$$p(t) = \begin{bmatrix} 1 & \frac{t}{h} & \left(\frac{t}{h}\right)^2 \end{bmatrix} \begin{pmatrix} \frac{1}{2} & 0 & 0 \\ -\frac{3}{2} & 2 & -\frac{1}{2} \\ \frac{1}{2} & -1 & \frac{1}{2} \end{pmatrix} \begin{pmatrix} f_0 \\ f_1 \\ f_2 \end{pmatrix} \quad (23)$$

then,  $p'(t) = \frac{1}{h} \left(-\frac{2}{3} f_0 + 2 f_1 - \frac{1}{2} f_2\right) + \frac{2t^*}{h^2} \left(\frac{1}{2} f_0 - 4 f_1 + f_2\right) = 0$  and taking  $h = \frac{1}{2}$  we obtain

$$t^* = \frac{3 f_0 - 4 f_1 + f_2}{4 (f_0 - 2 f_1 + f_2)} \quad (24)$$

### 4. PENALTY MODIFIED PQI TECHNIQUE (PMPQI)

The following numerical algorithm and the flowchart of the penalty function method combined with the Modified Partial Quadratic Interpolation Technique (Figure 1) are as follows:

1. Given  $g_i(x) \leq 0, i = 1, \dots, m$ , and  $h_j(x) = 0, j = 1, \dots, p, It = 0, \varepsilon' > 0, r_{it} > 0, \varepsilon > 0$ .
2. Initialize  $x^{(it)} \in \mathfrak{R}_n$  arbitrarily.

3. If the optimality conditions are satisfied at  $x^{(It)}$ , then stop.
4. Compute  $\Psi(x^{(It)}, r_{It}) = \min_{x \geq 0} \Psi(x, r_{It})$  and minimize  $x^{(It)}$  using MPQI technique and  $r_{It+1} = 10 \times r_{It}$ .
5. If  $\|x^{(It)} - x^{(It-1)}\| < \varepsilon$  or  $|\Psi(x^{(It)}, r_{It}) - \Psi(x^{(It-1)}, r_{It-1})| < \varepsilon$ ; then stop.  
Else  $It = It + 1$  and go to step 3.




Figure - 1: Penalty Modified Partial Quadratic Interpolation Method

## 5. NUMERICAL EXAMPLES

We present here some numerical examples to minimize the unconstrained and constrained optimization problems which use the modified partial quadratic interpolation technique (MPQI) and combined exterior penalty function-modified partial quadratic interpolation technique (PMPQI) and compare with other methods.

### A) Unconstrained Examples

**P1) Brown and Dennis function [16]:** This example gives the local minimum for the function

$$\text{Minimize } f(X) = \sum_{k=1}^{20} \left\{ (x_1 + t_k x_2 - e^{t_k}) + (x_3 + x_4 \sin(t_k) - \cos(t_k)) \right\}^2$$

where  $t_k = 0.2k$  with  $f(X^*) = 85822.2$

**P2) Extended Powell singular function [16]:** In this case we obtain the local minimum for the function :

$$\text{Minimize } f(X) = (x_1 + 10 x_2)^2 + 5(x_3 - x_4)^2 + (x_2 - 2 x_3)^4 + 10(x_1 - x_4)^4$$

where  $f(X^*) = 0.65 \times 10^{-13}$  at  $X^* = (0.18 \times 10^{-5}, 0.61 \times 10^{-5}, 0.60 \times 10^{-5})$

**P3) Rosenbrock function [16]:** This example gives the local minimum for the function

$$\text{Minimize } f(X) = 100(x_2 - x_1^2)^2 + (1.0 - x_1)^2$$

with  $f(X^*) = 0.0$  at  $X^* = (1, 1)$ .

In Table 1 we compare the numerical results obtained, for various starting points, by applying other methods (Armijo's quadratic method) [16], ARMBIS [17], Fletcher-Reeves [12]) and the modified partial quadratic interpolation technique (MPPQI). The first column in Table 1 contains the problem number and the next two columns of each method contain the total iterations (IT) and the total number of function and gradient evaluations (NFG) of each method.

Table 1									
Numerical results for Armijo, ARMBIS , Fletcher R. and MPQI									
P. NO.	Initial values	Armijo		ARMBIS		Fletcher R.		MPQI	
		IT	NFG	IT	NFG	IT	NFG	IT	NFG
P1	(0,0,0)	54	511	10	98	20	363	7	182
	(1,2,4,8)	72	709	15	146	30	523	8	364
	(-5,-1,1,5)	72	708	13	126	33	565	9	312
P2	(3,-1,0,1)	1197	16515	710	6799	8660	122575	17	417
	(1,-1,1,-1)	1184	16316	598	5712	610	86593	18	443
	(0,1,2,3)	1262	17505	635	6194	9943	140655	18	443
P3	(-1.2,1)	794	8901	957	5303	193	3211	32	482
	(-1.2,-1)	745	8283	485	2800	39	615	30	333
	(0,-1.2)	778	8697	484	2807	31	469	30	397

**B) Constrained Examples [3]**

**P4) Minimize**  $f(X) = 0.4 x_1 + 0.5 x_2$

**Subject to**  $0.3 x_1 + 0.1 x_2 \geq 2.7, 0.5 x_1 + 0.5 x_2 = 6.0, x_i \geq 0, i = \overline{1,2}$

**P5) Minimize**  $f(X) = 4.0 x_1 + 3.0 x_2$

**Subject to**  $2.0 x_1 + 3.0 x_2 \geq 6.0, 4.0 x_1 + x_2 \geq 4.0, x_i \geq 0, i = \overline{1,2}$

**P6) Minimize**  $f(X) = 3.0 x_1 + 8.0 x_2$

**Subject to**  $3.0 x_1 + 4.0 x_2 \leq 20.0, x_1 + 3.0 x_2 \geq 12.0, x_i \geq 0, i = \overline{1,2}$

Table 2						
Numerical results for polynomial penalty method (Algorithm 1), Karmarkar's Algorithm and PMPQI						
P. NO.	Algorithm 1 ( $\rho = 2$ )		Karmarkar's Algorithm		PMPQI	
	Total It	Time (Secs.)	Total It	Time (Secs.)	Total It	Time (Secs.)
P4	9	3.9	19	3.7	4	1.4634
P5	9	5.8	19	3.7	4	1.5622
P6	10	8.7	18	3.8	6	1.9125

Table 2 reports the results computational for Polynomial penalty method (Algorithm 1) (PPMA1), Karmarkar's Algorithm [13] and PMPQI technique. The first column in Table 1 contains the problem number and the next two columns of each method contain the total iterations and the times (in seconds) of each method.

**6. CONCLUSIONS**

In this paper, a computational approach based on an exterior penalty function MPQI method is given for solving a class of continuous inequality constrained optimization problems. The essential steps of the partial quadratic interpolation technique and its modified are given. The numerical algorithm and the flowchart of the penalty function method combined with the Modified Partial Quadratic Interpolation Technique are given. For illustration, three examples are solved using the proposed method. From the solutions obtained, we observe that the values of their object functions are amongst the smallest when compared with those obtained by other existing methods available in the literature. More importantly, our method finds solution which satisfies the continuous inequality constraints.

## REFERENCES

- [1]Jueyou Li, Zhiyou Wu and Qiang Long, A New Objective Penalty Function Approach for Solving Constrained Minimax Problems, J. Oper. Res. Soc. China, v.2, 2014, pp. 93-108.
- [2]Moengin P., Polynomial Penalty Method for Solving Linear Programming Problems, International Journal of Applied Mathematics, (40)3, 2010, 167-171.
- [3]Moengin P., Exponential Barrier Method in Solving Linear Programming Problems, International Journal of Engineering and Technology, (12)3, 2011, 33-37.
- [4]Yeniay, O., Penalty Function Methods for Constrained Optimization with Genetic Algorithms, Mathematical and Computational Application, 2005, 45-46.
- [5]Salim, M. S., Numerical Studies of Optimal Control Problems and Its Applications, Ph. D. Thesis, Assiut University, 1990.
- [6]Ivan L. Johnson, Lyndon B., The Davidson-Fletcher-Powell penalty function method- a generalized iterative technique for solving parameter optimization problems, National Aer. and Space Administration, 1976,1-21.
- [7]Winkler, M., The Quadratic Penalty Method for Solving Control-Constrained Optimal Control Problems, Universitat der Bundeswehr Munchen, Institut fur Mathematik und Bauinformatik 27th March, 2012, 1-22.
- [8]Changjun Y., Kok L. T., Liansheng Z. and Yanqin B., A new exact penalty function method for continuous inequality optimization problems, J.Industrial and Mangem. optimization, 6(4),2010,895-910
- [9]M. H. Farag, Application of the exterior penalty method for solving constrained optimal control problem, Math. Phys. Soc. Egypt., 1995, 1-12.
- [10] Aliay M. M.,Golalikhanib, Mohsen M. and Zhuang J., A computational study on different penalty approaches for solving constrained global optimization problems with the electromagnetism-like method, Optimization, iFirst, 2012,1-17.
- [11] Joghataie, A. and Takaloozadeh, M., Improving Penalty Functions for Structural Optimization, Sharif University of Technology, Tehran, Iran, 2009.
- [12] Rao, S. S., Optimization: Theory and Applications, Wiley Eastern Limited, (1984).
- [13] Rao, S. S., Engineering Optimization: Theory and Practice, New Age Inter. Limited Publishers, 3rd Ed, (1996).
- [14] El-Gindy, T. M., Numerical Studies in Optimal Control Theory, Ph. D. Thesis, University of Wales, 1977.
- [15] Radwan, A. A., Acceleration of partial quadratic interpolation technique, M. Sc. Thesis, Univ. of Assuit, 1980.
- [16] Jueyou Li, Zhiyou Wu and Qiang Long, A New Objective Penalty Function Approach for Solving Constrained Minimax Problems, J. Oper. Res. Soc. China, v.2, 2014, 93-108.
- [17] Joghataie, A. and Takaloozadeh, M., Improving Penalty Functions for Structural Optimization, Sharif University of Technology, Tehran, Iran, 2009.

**Source of support: Nil, Conflict of interest: None Declared**

***[Copy right © 2014. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]***