



**FUNCTIONS STRONGLY MCSHANE AND KURZWEIL- HENSTOCK INTEGRABLE  
IN BANACH SPACE**

<sup>1</sup>Ismet Temaj\* and <sup>2</sup>Agron Tato

<sup>1</sup>Faculty of Education, University of Prishtina, Kosovo

<sup>2</sup>Faculty of Sciences, University of Elbasan, Albania

E-mail: [itemaj63@yahoo.com](mailto:itemaj63@yahoo.com)

(Received on: 09-05-11; Accepted on: 20-05-11)

**ABSTRACT**

In this article, we characterize the  $S^*M$  integrable functions within the  $S^*HK$  integrable functions in Banach space, relativizing similar propositions in the real line case and generalizing classic results.

**Key Words and Phrases:** Banach space. McShane and Kurzweil-Henstock integrals.

**1. INTRODUCTION AND PRELIMINARIES:**

In this paper, we extend some relations between McShane integrals and Kurzweil- Henstock integrals, known for the real case [2]. We apply the extension for Banach space. All the material pertains to the propositions on mean value or so called "squeeze functions" that are characterized as the Vitali-Caratheodory propositions.

We consider functions  $f : I \rightarrow X$  where  $I \subset \mathbb{R}$  is a compact interval, and  $X$  is a Banach space with the norm  $\| \cdot \|_X$ . Based on the fact that respective definitions on Banach Space are well known if we consider e.g.: [1], [3], [4]. By  $\mu$  let the Lebesgue measure in  $\mathbb{R}$  be denoted.

An interval  $I$  is a compact subinterval of  $\mathbb{R}$ . A collection of intervals is called nonoverlapping if their interiors are disjoint. A partition  $P$  in  $I$  is a collection  $\{(I_i, t_i) : i = 1, 2, \dots, r\}$ , where  $I_1, \dots, I_r$  are nonoverlapping subintervals of  $I$  and  $t_1, \dots, t_r \in I$ . Let a compact interval  $I \subset \mathbb{R}$  be given, we say that  $P$  is

- (i) a partition in  $I$  if  $\bigcup_{i=1}^r I_i \subset I$
- (ii) a partition of  $I$  if  $\bigcup_{i=1}^r I_i = I$
- (iii) a Perron partition (or K – partition ) if  $t_i \in I_i, i = 1, \dots, r$ .
- (iv) a M-partition of  $I$  if  $t_i \in I, i = 1, \dots, r$ .

Given  $f : I \rightarrow X$  and partition  $P = \{(I_i, t_i) : i = 1, \dots, r\}$  in  $I$ , we set  $\sigma(f, P) = \sum_{i=1}^r f(t_i)\mu(I_i)$  and call this number the Riemann sum, of  $f$  associated with  $P$ .

Given  $\delta : I \rightarrow (0, +\infty)$ , called a gauge, a partition  $P = \{(I_i, t_i) : i = 1, \dots, r\}$  in  $I$  is called  $\delta$ – fine if  $I_i \subset (t_i - \delta(t_i), t_i + \delta(t_i))$ ,  $i = 1, 2, \dots, r$ .

A function  $f : I \rightarrow X$  is called - strongly measurable if there exists a sequence  $(f_n)_n$  of simple functions such that

$$f_n(t) \xrightarrow{n} f(t) \text{ a.e.}$$

**\*Corresponding author: Ismet Temaj\*, \*E-mail: [itemaj63@yahoo.com](mailto:itemaj63@yahoo.com)**

**Definition: 1** A function  $f : I \rightarrow X$  is said to be strongly measurable (or Bochner) integrable if there exists a sequence  $(f_n)_n$  of simple functions such that

- (i)  $f_n(t) \rightarrow f(t), \quad a.e.$
- (ii)  $\int_A f_n$  converges in  $X$  for each measurable subset  $A$  of  $I$ .

In this case we put

$$(B) \int_A f = \lim_{n \rightarrow \infty} \int_A f_n$$

**Definition: 2** A function  $f : I \rightarrow X$  is said to be McShane integrable, respectively Kurzweil- Henstock integrable, (briefly *McS*- integrable, respectively *KH*-integrable) on  $I$ , if there exists  $\omega \in X$  satisfying the following property: given  $\varepsilon > 0$  there exists a gauge  $\delta$  on  $I$  such that for each  $\delta$ -fine partition, respectively  $M$ -partition ( $K$ -partition),  $P = \{(I_i, t_i) : i = 1, \dots, r\}$  of  $I$ , we have

$$\| \sigma(f, P) - \omega \|_X < \varepsilon$$

Denote:  $\omega = (M) \int_I f(t) d\mu$  ( $\omega = (KH) \int_I f(t) d\mu$ ) and  $M$  ( $KH$ ) denotes the set of all McShane (Kurzweil- Henstock) integrable functions.

Given a set  $E \subset I$  we denote by  $\chi_E$  its characteristic function ( $\chi_E(t) = 1$  for  $t \in E$ ,  $\chi_E(t) = 0$  otherwise). A function  $f : I \rightarrow X$  is called *McShane (Kurzweil- Henstock) integral over the set  $E \subset I$*  if the function  $f \cdot \chi_E : I \rightarrow X$  is *McShane (Kurzweil- Henstock) intergable*.

In the case we write  $\int_I f \cdot \chi_E = \int_E f$ .

## 2. EXTENSION OF VITALI-CARATHEODORY THEOREM ON BANACH SPACE:

Recall some basic results of the integration on Banach spaces that used for our main proposition. We mainly refer to [1] Let  $Z$  denote the family of all compact subintervals  $J \subset I$ . A function  $F : Z \rightarrow X$  is said to be *additive* if

$$F(J \cup L) = F(J) + F(L)$$

for any nonoverlapping  $J, L \in Z$  such that  $J \cup L \in Z$ .

**Definition: 3** A function  $f : I \rightarrow X$  is said to be *strongly McShane integrable (Kurzweil-Henstock integrable)* on  $I$  if there is an additive function  $F : Z \rightarrow X$  such that for every  $\varepsilon > 0$  there exists a gauge  $\delta$  on  $I$  such that

$$\sum_{i=1}^k \| f(t_i) \mu(J_i) - F(J_i) \|_X < \varepsilon$$

for every  $\delta$ -fine  $M$ -partition ( $K$ -partition)  $P = \{(t_i, J_i) : i = 1, 2, \dots, k\}$  of  $I$ .

**Definition: 4** A function  $f : I \rightarrow X$  has the *property  $S^*M$  ( $S^*HK$ )* if for every  $\varepsilon > 0$  there is a gauge  $\delta$  on  $I$  such that

$$\sum_{i=1}^k \sum_{j=1}^l \| f(t_i) - f(s_j) \|_X \mu(J_i \cap L_j) < \varepsilon$$

for any  $\delta$ -fine  $M$ -partitions ( $K$ -partitions)  $P = \{(t_i, J_i) : i = 1, 2, \dots, k\}$  and  $Q = \{(s_j, L_j) : j = 1, 2, \dots, l\}$  of  $I$ .

**Theorem: 5** [1]. A function  $f : I \rightarrow X$  is Bochner integrable if and only if  $f$  has the property  $S^*M$  or, equivalently, if and only if  $f$  is strongly McShane integrable.

**Lemma: 6** [1]. Assume that  $f : I \rightarrow X$  is Bochner integrable and let  $\varepsilon > 0$  be given. Then there is a gauge  $\delta : I \rightarrow (0, +\infty)$  and  $\eta \in (0, \varepsilon)$  such that the following statement holds.

If  $P = \{(H_m, t_m) : m = 1, 2, \dots, r\}$  is  $\delta$ -fine  $M$ -system ( $K$ -system) for which

$$\sum_{m=1}^r \mu(H_m) < \eta$$

then

$$\sum_{m=1}^r \|f(t_m)\|_X \mu(H_m) < \varepsilon$$

**MAIN RESULT:**

**Theorem: 7** Let  $f : I \rightarrow X$  be a strongly measurable function. The following two statements are equivalent

- (a) function  $f$  is  $S^*M$  integrable on  $I$
- (b) there exists  $S^*HK$  integrable functions  $g(x)$  and  $h(x)$  such that for every  $\eta > 0$

$$\|f(x) - [g(x) + \theta(h(x) - g(x))]\|_X < \eta \tag{1}$$

and for every  $\varepsilon > 0$

$$(KH) \int_I \|h(x) - g(x)\|_X < \varepsilon \tag{2}$$

**Proof:** Assume that  $\varepsilon > 0$  is given. Since function  $f$  is  $S^*M$  integrable then it is Bochner integrable [1, p.146], for which there is an T- Cauchy sequence  $(f_q), q \in N$  of simple functions which converges to  $f$  almost everywhere in  $I$ . i.e.

$$\lim_{q \rightarrow \infty} \|f_q(t) - f(t)\|_X = 0 \tag{3}$$

for almost all  $t \in I$ .

Let  $\alpha > 0$  be real number. Considering [3 p.9], function  $f$  is measurable if and only if function  $f$  has the separable range a.e. on  $I$  so we can find a set  $V$  on  $I$  such that  $\mu(V) < \alpha$  and the range  $f(I \setminus V)$  is separable. Construct the set  $\{x_n \in f(I \setminus V) : n \in N\}$  which is dense everywhere in  $f(I \setminus V)$ . Fix any  $k \in N$  and denote

$$E_n^k = \left\{ s \in I \setminus V : \|f(s) - x_n\|_X < \frac{1}{k} \right\} = (I \setminus V) \cap f^{-1}\left[R_{\frac{1}{k}}(x_n)\right].$$

These sets are measurable. Since for every  $s \in I \setminus V$  and for every  $k$  exists the natural number  $n$  such that

$$\|f(s) - x_n\|_X < 1/k,$$

then we get

$$\bigcup_{n=1}^{\infty} E_n^k = I \setminus V.$$

Construct the sequence of sets

$$B_n^k = E_n^k \setminus (E_1^k \cup E_2^k \cup \dots \cup E_{n-1}^k).$$

We see that these sets are disjoint and

$$\bigcup_{n=1}^{\infty} B_n^k = I \setminus V.$$

Construct now the function with countable range

$$d^k(x) = \sum_{n=1}^{\infty} x_n 1_{B_n^k}(x) + 0 \cdot 1_{S \setminus V}(x) (k \in N).$$

We get that for every  $s \in I \setminus V$  and every  $k$

$$\|f(s) - d^k(x)\|_X < 1/k,$$

it follows that

$$\lim_{n \rightarrow \infty} \|f(s) - d^k(x)\|_X = 0$$

uniformly on  $I \setminus V$ . Since the set  $I \setminus V$  is measurable and its measure is not greater than  $I$ , then exists a number  $n_k \in N$  such that for  $k > n_k$

$$\sum_{n=n_k+1}^{\infty} m(B_n^k) < \frac{1}{k}.$$

Considering the neighborhood  $R_{\frac{1}{k}}(x_n)$  above mentioned, we construct  $R_{\frac{1}{2k}}(x_n) \subset R_{\frac{1}{k}}(x_n)$ .

Let  $y_n$  and  $z_n$  be elements of range  $f(I \setminus V)$  such that  $\|x_n - y_n\|_X < 1/2k$  and  $\|x_n - z_n\|_X < 1/2k$  and  $x_n = y_n + \theta(z_n - y_n)$  where  $\theta$  is real number  $0 \leq \theta \leq 1$ . It easy to see that  $y_n$  and  $z_n$  are inner the  $R_{\frac{1}{k}}(x_n)$

$$\|z_n - y_n\|_X < \|z_n - x_n\|_X + \|x_n - y_n\|_X < 1/k.$$

Construct two measurable functions

$$g(x) = \sum_{n=1}^{\infty} y_n 1_{B_n^k}(x) + 0 \cdot 1_V(x) \quad \text{and} \quad h(x) = \sum_{n=1}^{\infty} z_n 1_{B_n^k}(x) + 0 \cdot 1_V(x).$$

First, we prove the inequality (1). We obtain:

$\|f(s) - [g(s) + \theta(h(s) - g(s))]\|_X \leq \|f(s) - x_n\|_X + \|x_n - [g(s) + \theta(h(s) - g(s))]\|_X < 1/k + 1/k = 2/k$  for every  $B_n^k, n = 1, \dots, n_k$  and inequality may not hold for  $x \notin \bigcup_{n=1}^{n_k} B_n^k$ . Since the function  $f$  is  $S^*M$  integrable, then it is Bochner integrable by [1, p.146]. To show that  $g$  and  $h$  are  $S^*M$  integrable we can prove that  $d^k(x)$  is  $S^*M$  integrable.

Let  $p$  be a natural number  $p \in N$ . By (3) for every  $x \in V$  and  $q > p$  we get  $\|f(x) - f_q(x)\|_X < 1/k$ .

This inequality is satisfied for every  $B_n^k$ . It follows that for  $q > p$

$$\|d^k(x) - f_q(s)\|_X \leq \|d^k(x) - f_q(s)\|_X + \|f(s) - f_q(s)\|_X < 2/k$$

Observing the inequalities

$$\begin{aligned} \sum_{n=1}^{\infty} \|x_n\|_X m(B_n^k) &\leq \sum_{n=1}^{\infty} \|x_n - f_q(x)\|_X m(B_n^k) + \sum_{n=1}^{\infty} \|f_q(x)\|_X m(B_n^k) < \\ &< 2/k \sum_{n=1}^{\infty} m(B_n^k) + \sum_{n=1}^{\infty} \|f_q(x)\|_X m(B_n^k) < (B) \int_S \|f\|_X < \infty \end{aligned}$$

we obtain, by Lemma 1.4.1. [1, p.23], the function  $d^k(x)$  is Bochner absolute integrable and satisfies the condition  $S^*M$  therefore also the  $S^*HK$  condition.

To prove the inequality (2), we can solve the inequality for the Bochner integral and  $x \in I \setminus V$ . We have that

$$(B) \int_{S/V} \|g(x) - h(x)\|_X dm = \sum_{n=1}^{\infty} \|z_n - y_n\|_X m(B_n^k) < \frac{1}{k} m(S).$$

The right side vanish to zero if for every  $\varepsilon > 0$  we choose the number  $k$  such that  $\frac{1}{k} < \frac{\varepsilon}{2(b-a)}$ .

For the second part of theorem, we suppose that (b) holds. Set  $f_1(x) = g(x) + \theta(h(x) - g(x))$ .

By the condition, for every  $\varepsilon > 0$

$$\|f(x) - f_1(x)\| < \varepsilon$$

It follows that

$$\|f(x)\|_X \leq \|f_1(x)\|_X + \|f(x) - f_1(x)\|_X < \varepsilon + \|g(x)\|_X + \|h(x) - g(x)\|_X$$

By the [1, p.2], if  $f$  is measurable then  $\|f\|_X$  is also measurable. By the [3], consequence 1.1.4., p.29], if the function  $f(x)$  and  $g(x)$  are KH absolute integrable then they are Mcshane integrable. This implies  $\|f(x)\|_X$  is Bochner integrable.

According [1], proposition 5.1.2., p. 135] we obtain that  $f(x)$  is  $S^*M$  integrable.

**Corollary: 8 (Theorem Vitaly- Carathedory) [2]**

Let  $f$  be a function  $f : I \rightarrow R$ , the following statements are equivalent

- (a)  $f$  is M-integrable on  $I$ .
- (b)  $f$  is absolutely KH-integrable on  $I$
- (c) For every  $\varepsilon > 0$  there are absolutely KH-integrable functions  $g$  and  $h$  such that

$$g(x) \leq f(x) \leq h(x) \text{ on } I \text{ and } (KH) \int_I (h(x) - g(x)) < \varepsilon$$

**Proof:** In the case where  $X = R$ , it is obvious that equality

$$f(x) = g(x) + \theta(h(x) - g(x)), \quad (0 \leq \theta \leq 1)$$

implies

$$g(x) \leq f(x) \leq h(x).$$

For example, if  $\theta = 1/2$  we have

$$G(x) \leq f(x) = \frac{g(x) + h(x)}{2} \leq h(x).$$

**Lemma: 9** [1, p.133]. Assume that  $f : I \rightarrow X$  is Bochner integrable and let  $\varepsilon > 0$ . Then there is a gauge  $\delta : I \rightarrow (0, \infty)$  and  $\eta \in (0, \varepsilon)$  such that the following statement holds.

If is an  $\{(H_m, t_m), m = 1, 2, 3 \dots r\}$  HK – system (M – system)  $\delta$  – fine for which

$$\sum_{m=1}^r \mu(H_m) < \eta$$

then

$$\sum_{m=1}^r \|f(t_m)\|_X \mu(H_m) < \varepsilon$$

**Theorem: 10**

Let  $f$  be a function  $f : I \rightarrow R$ , the following statements are equivalent:

- (a) function  $f$  is  $S^*M$  -integrable
- (b) For every  $\varepsilon > 0$  there are absolutely  $S^*KH$  -integrable functions  $g$  and  $h$  such that

$$f(x) = g(x) + \phi(x)h(x) \text{ where } \phi(x) : I \rightarrow \{0, 1\} \tag{1}$$

and

$$(KH) \int_I \|H(x) - g(x)\|_X < \varepsilon. \tag{2}$$

**Proof:** Let us choose a gauge  $\delta : I \rightarrow ]0, \infty[$  as in Lemma 9 and  $\eta \in ]0, \varepsilon / 2[$ . Since function  $f$  is  $S^*M$  integrable then it is Bochner integrable and according to definition there exists a consequence of simple functions  $(f_q)$  with converge everywhere on  $I \setminus Z_a$ ,  $\mu(Z_a) = 0$ . By the Egorov theorem, there exists a subsequence of this sequence which is uniformly convergent for every  $x \in I \setminus V$ , when  $I \supset Z_a$  and  $\alpha < \eta / 4$ . This implies, that there exist the measurable disjoint sets  $S_i \subset I$ , such that  $\bigcup_{i=1}^{\infty} S_i = V$  and

$$f(x) = \sum_{i=1}^{\infty} C_i \cdot 1_{S_i}(x).$$

Since function  $f$  is Bochner integrable, it follows that below series is absolute convergent

$$\sum_{i=1}^{\infty} \|C_i\|_X \mu(S_i) = (B) \int_V \|f(x)\|_X < +\infty$$

We obtain

$$\sum_{i=N+1}^{\infty} \|C_i\|_X \mu(S_i) < \frac{\eta}{3}.$$

By the Lesbegue theorem there exists a closed set  $F_i$  and open set  $G_i$  such that

$$F_i \subset S_i \subset G_i$$

and  $\mu(G_i \setminus F_i) < \varepsilon / 2^{i+1}$ .

We observe that for every  $i$  the equality holds

$1_{S_i}(x) = 1_{F_i}(x) + \phi(x) \cdot 1_{G_i}(x)$  where  $\phi : I \rightarrow \{0, 1\}$ . According this equality, we construct the functions

$$g(x) = \sum_{i=1}^{\infty} C_i \cdot 1_{F_i}(x) + 0 \cdot 1_{I \setminus U}(x)$$

with  $U = \bigcup_{i=1}^{\infty} F_i \subset V$  and

$$h(x) = \sum_{i=1}^{\infty} C_i \cdot 1_{G_i}(x) + 0 \cdot 1_{I \setminus T}(x)$$

with  $T = \bigcup_{i=1}^{\infty} G_i \supset V$ . Reviewing the proof arguments of theorem 7, we conclude that these functions are Bochner integrable and it follows that they are absolute KH-integrable.

In order to prove (2) we consider inclusion  $I \setminus U \subset I \setminus T \cup T \setminus I$ .

Since

$$T \setminus U = \bigcup_{i=1}^{\infty} G_i \setminus \bigcup_{i=1}^{\infty} F_i = \bigcup_{i=1}^{\infty} (G_i \setminus F_i)$$

We get

$$m(T \setminus U) = m\left(\bigcup_{i=1}^{\infty} (G_i \setminus F_i)\right) \leq \sum_{i=1}^{\infty} m(G_i \setminus F_i) < \frac{\eta}{2}$$

If above K-system of the set,  $I \setminus U$  has been taken  $\square \square \delta$ -fine and satisfy

$$\sum_{m=1}^r m(H_m) \leq \sum_{i=1}^{\infty} m(G_i \setminus F_i) < \eta$$

then we have

$$\sum_{m=1}^r \|h(t_m) - g(t_m)\|_X m(H_m) < \varepsilon$$

This proves (2).

Second part of proof is the same as in Theorem 7.

#### REFERENCES:

- [1] Shwabik, S. Guoju, Y. Topics in Banach space integration (2005).
- [2] Lee, P.Y., Vyborny, R., The integral, an easy approach after Kurzweil and Hestock, Cambridge University Press, 2000.
- [3] Memetaj, S., Tema integrimi në hapësirat lokalisht konvekse, Mokra, Tiranë 2007.
- [4] Temaj, I., Tato, A., On McShane equiiterable sequences , Journal of Modern Mathematics and Statistics 4 (4): 123-126 2010 , ISSN: 1994-5388.

\*\*\*\*\*