

SOME NEW RELATIONSHIPS BETWEEN THE FRACTIONAL DERIVATIVES
OF FIRST, SECOND, THIRD AND FOURTH CHEBYSHEV WAVELETS

Sameer Qasim Hasan* and Dalia Raad Abd

Department of mathematics, Collage of education, Almustansiriyah University, Iraq.

(Received On: 09-07-14; Revised & Accepted On: 25-07-14)

ABSTRACT

In this paper, the shifted first, second, third and fourth kind Chevelets wavelets $\Psi_{nm}^1(t)$, $\Psi_{nm}^2(t)$, $\Psi_{nm}^3(t)$ and $\Psi_{nm}^4(t)$ properties are presented.

The main aim is:

1. Generalize the first, second operational matrix to the fractional derivatives . In this approach, a truncated first, second matrix of fractional derivatives are used.
2. Presented a new proposal formula expressing of fractional derivative $\alpha > 0$ operational matrix of shifted first kind Chybeshev wavelets $D^\alpha \Psi_{nm}^1(t)$ interms of $D^\alpha \phi(x)$, $\phi(x) = [T_0^*(x), T_1^*(x), \dots, T_n^*(x)]^T$, and a formula expressing the fractional derivative $\alpha > 0$ of second kind Chebyshev wavelets $D^\alpha \Psi_{nm}^2(t)$ interms of $D^\alpha \varphi(x)$, $\varphi(x) = [U_0^*(x), U_1^*(x), \dots, U_n^*(x)]^T$.
3. Presented a new proposal formula expressing of fractional derivative $\alpha > 0$ operational matrix of shifted third and fourth kind Chybeshev wavelets $D^\alpha \Psi_{nm}^3(t)$, $D^\alpha \Psi_{nm}^4(t)$ interms of $\Psi_{nm}^2(t)$ and $\Psi_{nm-1}^2(t)$, and a formula expressing of fractional derivative $\alpha > 0$ of first kind Chebyshev wavelets $D^\alpha \Psi_{nm}^1(t)$ interms of $\Psi_{nm}^2(t)$ and $\Psi_{nm-2}^2(t)$.

All the proposed results are of direct interest in many applications.

1. INTRODUCTION

The Chebyshev polynomials are one of the most useful polynomials, which are suitable in numerical analysis including polynomial approximation, integral and differential equations and spectral methods for partial differential equations [4, 9, 10, 17]. One of the attractive concepts in the initial and boundary value problems is differentiation and integration of fractional order [8, 16, 18, 19]. Many researchers extend classical methods in studies of differential and integral equations of integer order to fractional type of these problems [15, 21]. One of the wide classes of researches focuses on constructing the operational matrix of derivative in some spectral methods. Recently, a lot of attention has been devoted to construct operational matrix of fractional derivative [4, 14, 20]. For example the fractional type first kind chebyshev polynomials are used to solving fractional diffusion equations [3, 7] also are used to solve multi-order fractional equation [12].

In this paper we use shifted chebyshev polynomials of first, second, third and fourth kind and recall some important properties. Next we used obtain the operational matrix of fractional derivative. Wavelets theory is a relatively new emerging in mathematical research [5, 6, 11, 23]. It has been applied in a wide range of engineering disciplines, particularly, shifted first kind chebyshev wavelets play an important role in establishing algebraic methods for the solution of multi-order fractional differential equations, [22] and initial and boundary values problems of fractional order, [13]. New Spectral Second Kind Chebyshev Wavelets Algorithm for Solving Linear and Nonlinear Second-Order Differential Equations, [2].

Corresponding author: Sameer Qasim Hasan*

Department of mathematics, Collage of education, Almustansiriyah University, Iraq.

2. PROPOSAL OPERATIONAL MATRIX OF CHEBYSHEV'S FRACTIONAL DERIVATIVE

A. It well know the first kind chebyshev polynomial $T_n(z)$ of degree n , [22], which defined on $[-1,1]$ by:

$$T_n(z) = \text{Cos}(n \theta) \quad \text{where } z = \text{Cos}(\theta) \theta \in [0, \pi]$$

and can be determined with the aid of the following recurrence formula:

$$T_{n+1}(z) = 2zT_n(z) - T_{n-1}(z). \quad n=1, 2, 3, \dots$$

$$T_0(z) = 1, T_1(z) = z$$

The analytic form of chebyshev polynomial $T_n(x)$ of degree (n) is given by:

$$T_n(z) = \sum_{i=0}^{n/2} (-1)^i 2^{n-2i-1} \frac{n(n-1)!}{i!(n-2i)!} z^{n-2i} \quad n=2, \dots \quad (1)$$

and are orthogonal on $[1,-1]$ with respect to the weight function $\omega(t) = 1/\sqrt{1-z^2}$, that is:

$$\int_{-1}^1 \frac{T_i(z)T_j(z)}{\sqrt{1-z^2}} dz = \begin{cases} \pi & i = j = 0 \\ \frac{\pi}{2} & i = j \neq 0 \\ 0 & i \neq j \end{cases}$$

In order to use these polynomials on the interval $[0, 1]$, we define the shifted chebyshev polynomials by introducing the change variable $z=2x-1$, then $T_n^*(x)$ can be obtained as follows:

$$T_{i+1}^*(x) = 2(2x-1) T_i^*(x) - T_{i-1}^*(x) \quad i=1, 2, \dots$$

where $T_0^*(x)=1, T_1^*(x)=2x-1$ and the analytic form is $T_n^*(x) = \sum_{i=0}^n (-1)^i 2^{2n-2i} \frac{n(2n-i-1)!}{i!(2n-2i)!} x^{n-i}$, $n=2, 3, \dots$ and are orthogonal with respect to the weight function $\omega(x)=1/\sqrt{x-x^2}$, that is:

$$\int_0^1 \frac{T_i^*(x)T_j^*(x)}{\sqrt{x-x^2}} dx = \begin{cases} \pi & i = j = 0 \\ \frac{\pi}{2} & i = j \neq 0 \\ 0 & i \neq j \end{cases}$$

The function $u(x)$ square integrable in $[0, 1]$, may be expressed in the term of shifted first kind chebyshev polynomial as:

$$\sum_{i=0}^{\infty} c_i T_i^*(x)$$

where the coefficients c_i are given by

$$c_i = \frac{\pi \sigma_i}{2} \int_0^1 u(x) T_i^*(x) dx, \quad \sigma_i = \begin{cases} 2 & i = 0 \\ 1 & i \neq 0 \end{cases}$$

In practice, only the first $(m+1)$ -terms shifted first kind chebyshev polynomial are considered. Then we have:

$$u_m(x) = \sum_{i=0}^m c_i T_i^*(x) = C^T \phi(x)$$

where the shifted first kind chebyshev coefficient vector C and the shifted first kind chebyshev vector $\phi(x)$ are given by: $C^T = [c_0(x), c_1(x), \dots, c_m(x)]$, $\phi(x) = [T_0^*(x), T_1^*(x), \dots, T_m^*(x)]^T$.

For the Caputo's derivative we have, [22]: $D^\alpha C = 0$, C is a constant.

$$D^\alpha x^n = \begin{cases} \frac{\Gamma(n+1)}{\Gamma(n+1-\alpha)} x^{n-\alpha} & n \geq [\alpha] \text{ for } n \in \mathbb{N}_0 \\ 0 & n < [\alpha] \text{ for } n \in \mathbb{N}_0 \end{cases} \quad (2)$$

for $\mathbb{N}_0 = \{0, 1, 1, \dots\}$.

In the following theorem we will define the fractional derivative of the vector $\phi(x)$.

Theorem 2.1: Let $\phi(x)$ be shifted first kind chebyshev vector defined as $\phi(x) = [T_0^*(x), T_1^*(x), \dots, T_m^*(x)]^T$ and also suppose $\alpha > 0$ then $D^\alpha \phi(x) = \Delta^{(\alpha)} \phi(x)$

where $\Delta^{(\alpha)}$ is $(m+1) \times (m+1)$ is an operational matrix of fractional derivative of order $\alpha > 0$ in the caputo sense and is defined as follows :

$$\Delta^\alpha = \begin{bmatrix} 0 & 0 & 0 \\ \vdots & \vdots & \vdots \\ 0 & 0 & 0 \\ w_{0,0,i}^{(1)} & w_{0,1,i}^{(1)} & \dots & w_{0,m,i}^{(1)} \\ \vdots & \vdots & \vdots & \vdots \\ \sum_{i=0}^{n-[\alpha]} w_{n-[\alpha],0,i}^{(1)} & \sum_{i=0}^{n-[\alpha]} w_{n-[\alpha],1,i}^{(1)} & \dots & \sum_{i=0}^{n-[\alpha]} w_{n-[\alpha],m,i}^{(1)} \\ \vdots & \vdots & \vdots & \vdots \\ \sum_{i=0}^m w_{m,0,i}^{(1)} & \sum_{i=0}^m w_{m,1,i}^{(1)} & \dots & \sum_{i=0}^m w_{m,m,i}^{(1)} \end{bmatrix}$$

and

$w_{n-[\alpha],j,i}^{(1)}$ is given by:

$$w_{n-[\alpha],j,i}^{(1)} = \frac{\sigma_j}{\sqrt{\pi}} \sum_{k=0}^j (-1)^{k+i} 2^{2j-2k+2n-2i} \frac{n(2n-i-1)j!(2j-k-1)! [(n-i-\alpha+j-k+\frac{1}{2})]}{i!(2n-2i)! [(n-i-\alpha+1)k!(2j-2k)! [(n-i-\alpha+j-k+1)]}$$

where $n=[\alpha] \dots m$ and $\sigma_j = \begin{cases} 1 & j = 0 \\ 2 & j \neq 0 \end{cases}$

Note that in Δ^α , the first $[\alpha]$ rows, are all zero.

Proof: Let $T_n^*(x)$ be shifted first kind chebyshev polynomial then by using (1) and (2) we can find that: $D^\alpha T_n^*(x) = 0, n < [\alpha]$ and for $n \geq [\alpha]$

$$D^\alpha T_n^*(x) = \sum_{i=0}^n (-1)^i 2^{2n-2i} \frac{n(2n-i-1)!}{i!(2n-2i)!} D^\alpha X^{n-i} \\ = \sum_{i=0}^{n-[\alpha]} (-1)^i 2^{2n-2i} \frac{n(2n-i-1)!}{i!(2n-2i)! [(n-i-\alpha+1)]} X^{n-i-\alpha}$$

Now, approximate $(x^{n-i-\alpha})$ by $(m+1)$ -terms of shifted first kind chebyshev polynomial, we have

$$x^{n-i-\alpha} = \sum_{j=0}^m d_{n-i,j} T_j^*(x) \text{ where } d_{n-i,j} = \frac{\sigma_j}{\pi} \int_0^1 \frac{x^{n-i-\alpha}}{\sqrt{x-x^2}} T_j^*(x) dx$$

$$T_j^*(x) = \sum_{k=0}^j (-1)^k 2^{2j-2k} \frac{j(2j-k-1)!}{k!(2j-2k)!} x^{j-k}, \text{ then}$$

$$d_{n-i,j} = \frac{\sigma_j}{\pi} \sum_{k=0}^j (-1)^k 2^{2j-2k} \frac{j(2j-k-1)!}{k!(2j-2k)!} \int_0^1 \frac{x^{n-i-\alpha+j-k}}{\sqrt{x-x^2}} dx \\ = \frac{\sigma_j}{\pi} \sum_{k=0}^j (-1)^k 2^{2j-2k} \frac{j(2j-k-1)! [(n-i-\alpha+j-k+\frac{1}{2})\sqrt{\pi}]}{k!(2j-2k)! [(n-i-\alpha+j-k+1)]}$$

where,

$$\sigma_j = \begin{cases} 1 & j = 0 \\ 2 & j \neq 0 \end{cases} \text{ then}$$

$$D^\alpha T_n^*(x) = \sum_{i=0}^{n-[\alpha]} \sum_{j=0}^m (-1)^i 2^{2n-2i} \frac{n(2n-i-1)!(n-i)!}{i!(2n-2i)! [(n-i-\alpha+1)]} d_{n-i,j} T_n^*(x) \\ = \sum_{j=0}^m [\sum_{i=0}^{n-[\alpha]} w_{n,j,i}^{(1)}] T_n^*(x), \text{ for } n \geq [\alpha]. \\ = [\sum_{i=0}^{n-[\alpha]} w_{n-[\alpha],i,0}^{(1)}, \sum_{i=0}^{n-[\alpha]} w_{n-[\alpha],i,1}^{(1)}, \dots, \sum_{i=0}^{n-[\alpha]} w_{n-[\alpha],i,m}^{(1)}] \emptyset(x), \text{ for } n \geq [\alpha]$$

and $D^\alpha T_n^*(x) = [0, \dots, 0] \emptyset(x), n < [\alpha]$.

B. The second kind of degree (n), [6], which defined on the interval $[-1, 1]$ as:

$$U_n(z) = \frac{\sin(n+1)\theta}{\sin \theta} \text{ where } z = \cos \theta, \theta \neq n\pi + 2k\pi. B$$

These polynomials satisfy the following recurrence relation

$$U_0(z) = 1, U_1(z) = 2z, \\ U_n(z) = 2zU_{n-1}(z) - U_{n-2}(z), n = 2, 3, \dots$$

For using this polynomials on interval (0,1) which called shifted chebyshev polynomials,[6], by introducing the change variable $(2x - 1)$ and satisfy the following $U_n^*(x) = (4x - 2)U_{n-1}^*(x) - U_{n-2}^*(x), n = 2, 3, \dots$

where $U_0^*(x) = 1, U_1^*(x) = 4x - 2$

and the analytic form of shifted chebyshev polynomials $u_n^*(x)$ of degree (n) is given by

$$U_n^*(x) = \sum_{r=0}^{n+1} r (-1)^{n+1-r} \frac{(n+r)! 2^{2r-1}}{(n+1-r)! 2r!} x^{r-1} \quad (3)$$

and are orthogonal with respect to the weight function $\omega(x) = 1/\sqrt{x-x^2}$, that is:

$$\int_0^1 \frac{U_m^*(x) \cdot U_n^*(x)}{\sqrt{x-x^2}} dx = \begin{cases} \frac{\pi}{8} & m = n \\ 0 & m \neq n \end{cases}$$

The function $f(x)$ square integrable in $[0, 1]$, may be expressed in the term of shifted first kind chebyshev polynomial as:

$$\sum_{i=0}^{\infty} c_i U_i^*(x)$$

where the coefficients c_i are given by

$$c_i = \frac{8}{\pi} \int_0^1 f(x) T_i^*(x) dx,$$

In practice, only the first (m+1)-terms shifted first kind chebyshev polynomial are considered. Then we have:

$$f_m(x) = \sum_{i=0}^m c_i T_i^*(x) = C^T \varphi(x)$$

where the shifted first kind chebyshev coefficient vector C and the shifted first kind chebyshev vector $\varphi(x)$ are given by: $C^T = [c_0(x), c_1(x), \dots, c_m(x)]$, $\varphi(x) = [U_0^*(x), U_1^*(x), \dots, U_m^*(x)]^T$.

In the following theorem we will define the fractional derivative of the vector $\varphi(x)$.

Theorem 2.2: Let $\varphi(x)$ be shifted second kind chebyshev vector defined in $\varphi(x) = [U_0^*(x), U_1^*(x), \dots, U_n^*(x)]^T$ also suppose $\alpha > 0$ then

$$D^\alpha \varphi(x) = \Delta^{(\alpha)} \varphi(x)$$

where Δ^α is the $(m_1 + 1) \times (m_1 + 1)$ operational Matrix of fractional derivative of order α in the Caputo sense and defined as follows:

$$\Delta^\alpha = \begin{bmatrix} 0 & 0 & 0 \\ \vdots & \vdots & \vdots \\ 0 & 0 & 0 \\ \sum_{r=[\alpha]+1}^{[\alpha]+1} w_{[\alpha]+1,0,r}^{(2)} & \sum_{r=2}^2 w_{[\alpha]+1,1,r}^{(2)} & \dots & \sum_{r=2}^2 w_{[\alpha]+1,m_1,r}^{(2)} \\ \vdots & \vdots & \vdots & \vdots \\ \sum_{r=[\alpha]+1}^{n+1} w_{n+1,0,r}^{(2)} & \sum_{r=[\alpha]+1}^{n+1} w_{n+1,1,r}^{(2)} & \dots & \sum_{r=[\alpha]+1}^{n+1} w_{n+1,m_1,r}^{(2)} \\ \vdots & \vdots & \vdots & \vdots \\ \sum_{r=[\alpha]+1}^{m_1} w_{m_1,0,r}^{(2)} & \sum_{r=[\alpha]+1}^{m_1} w_{m_1,1,r}^{(2)} & \dots & \sum_{r=[\alpha]+1}^{m_1} w_{m_1,m_1,r}^{(2)} \end{bmatrix}$$

and

$$w_{n+1,p,r}^{(2)} = \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r \cdot \ell (-1)^{n+p+2-(\ell+r)} (n+r)! (p+\ell)! (r-1)! 2^{2(\ell+r)-2} \left[(r-\alpha+\ell+\frac{1}{2}) \right]}{(n+1-r)! 2r! [(r-\alpha)(p+1-\ell)! 2\ell! [(r-\alpha+\ell+2)]}$$

Note that in Δ^α , the first $[\alpha] + 1$ rows, are all zero

Proof: Let $U_m^*(x)$ be shifted sccondt kind chebyshev polynomial then by using (2) and (3) we can find that:

$D^\alpha U_n^*(x) = 0n \leq [\alpha]$ and for $(n = [\alpha] + 1 \dots m)$ we have

$$\begin{aligned} D^\alpha U_n^*(x) &= \sum_{r=0}^{n+1} r (-1)^{n+1-r} \frac{(n+r)! 2^{2r-1}}{(n+1-r)! 2r!} D^\alpha x^{r-1} \\ &= \sum_{r=[\alpha]+1}^{n+1} r (-1)^{n+1-r} \frac{(n+r)! 2^{2r-1} (r-1)!}{(n+1-r)! 2r! [(r-\alpha)]} x^{r-\alpha-1} \end{aligned}$$

Now, approximate $(x^{r-\alpha-1})$ by $(m + 1)$ -terms of shifted second kind chebyshev series, we have

$$x^{r-\alpha-1} = \sum_{p=0}^{m_1} d_{r-1,p} U_p^*(x)$$

$$\begin{aligned} d_{r-1,p} &= \frac{8}{\pi} \int_0^1 x^{r-\alpha-1} \sqrt{x-x^2} U_p^*(x) dx \\ &= \frac{8}{\pi} \sum_{\ell=0}^{p+1} \frac{\ell(-1)^{p+1-\ell} (p+\ell)! 2^{2\ell-1}}{(p+1-\ell)! 2\ell!} \int_0^1 \sqrt{x-x^2} x^{r-\alpha-1} dx \\ &= \frac{8}{\pi} \sum_{\ell=0}^{p+1} \frac{\ell(-1)^{p+1-\ell} (p+\ell)! 2^{2\ell-1} \Gamma(r-\alpha+\ell+\frac{1}{2}) \cdot \sqrt{\pi}}{(p+1-\ell)! 2\ell! \Gamma(r-\alpha+\ell+2)} \end{aligned}$$

$$\begin{aligned} D^\alpha U_n^*(x) &= \sum_{r=[\alpha]+1}^{n+1} \sum_{p=0}^{m_1} r(-1)^{n+1-r} \frac{(n+r)! 2^{2r-1} (r-1)!}{(n+1-r)! 2r! [(r-\alpha)]} d_{r-1,p} U_n^*(x) \\ &= \sum_{p=0}^{m_1} \left[\sum_{r=[\alpha]+1}^{n+1} w_{n+1,p,r}^{(2)} \right] U_n^*(x) \\ &= \left[\sum_{r=[\alpha]+1}^{n+1} w_{n+1,0,r}^{(2)}, \sum_{r=[\alpha]+1}^{n+1} w_{n+1,1,r}^{(2)}, \dots, \sum_{r=[\alpha]+1}^{n+1} w_{n+1,m_1,r}^{(2)} \right] \varphi(x), \end{aligned}$$

for $n=[\alpha] + 1 \dots m_1$ and $D^\alpha U_n^*(x) = [0, \dots, 0] \varphi(x)$ $n \leq [\alpha]$.

3. PROPOSAL OPERATIONAL MATRIX OF CHEBYSHEV WAVELETS FRACTIONAL DERIVATIVE

A. This shifted first kind chebyshev wavelets $\Psi_{\tilde{n}m}^1(t) = \Psi^1(k, \tilde{n}, m, t)$ have four arguments; $k \in \mathbb{N}$, $n=1,2,\dots,2^{k-1}$ and $\tilde{n} = 2n - 1$; moreover, m is the order of the chebyshev polynomials of the first kind and t is the normalized time, and they are defined on the interval $[0,1)$ as, [12]:

$$\Psi_{\tilde{n}m}^1(t) = \begin{cases} 2^{k/2} T_m^*(2^k t - \tilde{n}) & \frac{\tilde{n}-1}{2^k} \leq t \leq \frac{\tilde{n}}{2^k} \\ 0 & \text{o.w} \end{cases}$$

where $T_m^* = \begin{cases} \frac{1}{\sqrt{\pi}} T_m & m = 0 \\ \sqrt{\frac{2}{\pi}} T_m & m > 0 \end{cases}$

$m = 0, 1 \dots M-1$, $n = 1, 2, \dots, 2^{k-1}$ and $\tilde{n}=0,1,\dots,2^k-1$, the weight function

$$\tilde{w} = w(2t - 1) \text{ and } w_{\tilde{n}}(t) = w(2^k t - \tilde{n}) \text{ where } w(2^k t - \tilde{n}) = \frac{1}{\sqrt{(2^k t - \tilde{n}) - (2^k t - \tilde{n})^2}}$$

A function $f(t)$ defined over $[0,1)$ may be expanded as follows

$$f(t) = \sum_{\tilde{n}=0}^{\infty} \sum_{m=0}^{\infty} c_{\tilde{n}m} \Psi_{\tilde{n}m}^1(t) \text{ where}$$

$$c_{nm} = (f(t), \Psi_{\tilde{n}m}^1(t))_w = \int_0^1 \tilde{w}(t) f(t) \cdot \Psi_{\tilde{n}m}^1(t) dt \text{ and } f(t) = \sum_{\tilde{n}=1}^{\infty} \sum_{m=0}^{\infty} c_{\tilde{n}m} \Psi_{\tilde{n}m}^1(t) = C^T \Psi_{\tilde{n}m}^1(t)$$

where $C = [c_{00}, c_{01}, \dots, c_{2^k-1,M}, \dots, c_{2^k-1,1}, \dots, c_{2^k-1,M}]^T$

Thus,

$$\Psi_{\tilde{n}m}^1(t) = [\Psi_{0,0}^1, \Psi_{0,1}^1, \dots, \Psi_{0,M}^1, \dots, \Psi_{2^k-1,M}^1, \dots, \Psi_{2^k-1,1}^1, \dots, \Psi_{2^k-1,M}^1]^T$$

Theorem 3.1: Let $\Psi_{\tilde{n}m}^1(t)$ be shifted of first kind chebyshev vector and also suppose $\alpha > 0$) then

$D^\alpha \Psi_{\tilde{n}m}^1(t)(t) = D^\alpha \left(\frac{c_m \cdot 2^{k/2}}{\sqrt{\pi}} T_m^*(2^k t - \tilde{n}) \right) = \Delta^{(\alpha)} \Psi_{\tilde{n}m}^1(t)$, such that $C_m = \begin{cases} 1 & m = 0 \\ \sqrt{2} & m > 0 \end{cases}$, where $\Delta^{(\alpha)}$ is the $(m+1) \times (m+1)$ operational matrix derivative of order (α) in the caputo sense and is defined as follows:

$$\Delta^{(\alpha)} = \begin{bmatrix} 0 & 0 & 0 \\ \vdots & \vdots & \vdots \\ 0 & 0 & 0 \\ \vdots & w_{0,0,i} \tilde{w}_{0,1,i} & \dots & w_{0,M,i} \\ \vdots & \vdots & \vdots & \vdots \\ \sum_{i=0}^{m-[\alpha]} w_{m-[\alpha],0,i} \tilde{w}_{m-[\alpha],1,i} & \sum_{i=0}^{m-[\alpha]} w_{m-[\alpha],1,i} & \dots & \sum_{i=0}^{m-[\alpha]} w_{m-[\alpha],M,i} \\ \vdots & \vdots & \vdots & \vdots \\ \sum_{i=0}^{m_1} w_{M,0,i} \tilde{w}_{M,1,i} & \sum_{i=0}^{m_1} w_{M,1,i} & \dots & \sum_{i=0}^{m_1} w_{M,M,i} \end{bmatrix}$$

And $w_{m-[\alpha]j,i}$ is given by

$$w_{m-[\alpha],j,i}^{\sim} = \frac{\sigma_j}{\pi} \sum_{k=0}^j (-1)^{k+i} 2^{2(j+m)-2(k+i)} \cdot \frac{m(2m-i-1)!(m-i)!j(2j-k-1)! [(m-i-\alpha+j-k+\frac{1}{2})]}{i!(2m-2i)! [(m-i-\alpha+1)k!(2j-2k)! [(m-i-\alpha+k+1)]}$$

where,

$$\sigma_j = \begin{cases} 1 & j = 0 \\ 2 & j \neq 0 \end{cases}$$

Note that in Δ^α , the first $[\alpha]$ rows, are all zero.

Proof: $D^\alpha T_m^* (2^k t - \tilde{n}) = 0 \quad m < [\alpha]$ and for $(m \geq [\alpha])$

$$\begin{aligned} D^\alpha T_m^* (2^k t - \tilde{n}) &= \sum_{i=0}^m (-1)^i 2^{2m-2i} \frac{m(2m-i-1)!}{i!(2m-2i)!} D^\alpha (2^k t - \tilde{n})^{m-i} \\ &= \sum_{i=0}^{m-[\alpha]} (-1)^i 2^{2m-2i} \frac{m(2m-i-1)!}{i!(2m-2i)!} (2^k t - \tilde{n})^{m-i-\alpha} \end{aligned}$$

Now, approximate $(2^k t - \tilde{n})^{m-i-\alpha}$ by $(m+1)$ terms of shifted first kind chebyshev wavelet, we have

$$(2^k t - \tilde{n})^{m-i-\alpha} = \sum_{j=0}^m d_{m-i,j} T_j^* (2^k t - \tilde{n})$$

$$\begin{aligned} d_{m-i,j} &= \frac{\sigma_j}{\pi} \int_0^1 (2^k t - \tilde{n})^{n-i-\alpha} \frac{T_j^* (2^k t - \tilde{n})}{\sqrt{(2^k t - \tilde{n}) - (2^k t - \tilde{n})^2}} dt \\ &= \frac{\sigma_j}{\pi} \sum_{k=0}^j (-1)^k 2^{2j-2k} \frac{j(2j-k-1)!}{k!(2j-2k)!} \int_0^1 \frac{(2^k t - \tilde{n})}{\sqrt{(2^k t - \tilde{n}) - (2^k t - \tilde{n})^2}} dt \end{aligned}$$

where,

$$\sigma_j = \begin{cases} 1 & j = 0 \\ 2 & j \neq 0 \end{cases}, \text{ then}$$

$$d_{m-i,j} = \frac{\sigma_j}{\pi} \sum_{k=0}^j (-1)^k 2^{2j-2k} \frac{j(2j-k-1)! [(m-i-\alpha+j-k+\frac{1}{2})]}{k!(2j-2k)! [(m-i-\alpha+k+1)]}$$

Therefore,

$$\begin{aligned} D^\alpha T_m^* (2^k t - \tilde{n}) &= \sum_{i=0}^{m-[\alpha]} \sum_{j=0}^{m-1} (-1)^i 2^{2m-2i} \frac{m(2m-i-1)!(m-i)!}{i!(2m-2i)! [(m-i-\alpha+1)]} d_{m-i,j} T_m^* (2^k t - \tilde{n}) \\ &= \sum_{j=0}^{m-1} [\sum_{i=0}^{m-[\alpha]} w_{m,j,i}] T_m^* (2^k t - \tilde{n}). \end{aligned}$$

$$D^\alpha \Psi_{\tilde{n}m}^1(t) = [\sum_{i=0}^{m-[\alpha]} w_{m-[\alpha],i,0}^{\sim}, \sum_{i=0}^{m-[\alpha]} w_{m-[\alpha],i,1}^{\sim}, \dots, \sum_{i=0}^{m-[\alpha]} w_{m-[\alpha],i,M}^{\sim}] \Psi_{\tilde{n}m}^1(t), \text{ for } m \geq [\alpha]$$

$$\text{and } D^\alpha \Psi_{\tilde{n}m}^1(t) = [0, \dots, 0] \Psi_{\tilde{n}m}^1(t) \quad \text{for } m < [\alpha].$$

B. Second kind chebyshev wavelets $\Psi_{nm}^2(t) = \Psi^2(k, n, m, t)$ have four arguments k, n can assume any positive integer, m is the order of second kind chebyshev polynomials, and t is the normalized time. They are defined on the interval $[0, 1]$ by: [2]

$$\Psi_{nm}^2(t) = \begin{cases} \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} u_m^*(2^k t - n) & t \in [\frac{n}{2^k}, \frac{n+1}{2^k}] \quad m=0,1,\dots,M, n=0,1,\dots,2^k-1 \\ 0 & \text{o.w} \end{cases}$$

and $w(2t-1)$ has to be dilated and translated as follows:

$$w_n(2^k t - n) = \sqrt{(2^k t - n) - (2^k t - n)^2} \text{ and the function approximation}$$

A function $f(t)$ defined over $[0, 1]$ may be expanded in terms of second kind Chebyshev wavelets as

$$f(t) = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} c_{nm} \Psi_{nm}^2(t)$$

$$c_{nm} = (f(t), \Psi_{nm}^2(t)) = \int_0^1 w(t) \cdot f(t) \cdot \Psi_{nm}^2(t) dt \text{ and}$$

$(t) = \sqrt{t - t^2}$ If the infinite series is truncated, then (t) can be approximated as

$$f(t) = \sum_{n=0}^{2^k-1} \sum_{m=0}^M c_{nm} \Psi_{nm}^2(t) = C^T \Psi_{nm}^2(t)$$

where C and $\Psi(t)$ are $2^k (M+1) \times 1$ matrices defined by

$$c = [c_{00}, c_{01}, \dots, c_{2^k-1,M}, \dots, c_{2^k-1,1}, \dots, c_{2^k-1,M}]^T$$

$$\Psi_{nm}^2(t) = [\Psi_{0,0}^2, \Psi_{0,1}^2, \dots, \Psi_{0,M}^2, \dots, \Psi_{2^k-1,M}^2, \dots, \Psi_{2^k-1,1}^2, \dots, \Psi_{2^k-1,M}^2]$$

Theorem 3.2: Let $\Psi_{nm}^2(t)$ be second kind chebyshev wavelets and suppose

$$\alpha > 0 \text{ then } D^\alpha \Psi_{nm}^2(t) = \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} D^\alpha (u_m^*(2^k t - n)) = \Delta^{(\alpha)} \Psi_{nm}^2(t)$$

where Δ^α is $(M+1) \times (M+1)$ operational matrix derivative of order α in the caputo sense and defined as follow:

$$\Delta^\alpha = \begin{bmatrix} 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \\ \sum_{r=|\alpha|+1}^{|\alpha|+1} \tilde{w}_{|\alpha|+1,0,r} & \sum_{r=2}^2 \tilde{w}_{|\alpha|+1,1,r} & \dots & \sum_{r=2}^2 \tilde{w}_{|\alpha|+1,M,r} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{r=|\alpha|+1}^{m+1} \tilde{w}_{m+1,0,r} & \sum_{r=|\alpha|+1}^{n+1} \tilde{w}_{m+1,1,r} & \dots & \sum_{r=|\alpha|+1}^{n+1} \tilde{w}_{m+1,M,r} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{r=|\alpha|+1}^M \tilde{w}_{M,0,r} & \sum_{r=|\alpha|+1}^M \tilde{w}_{M,1,r} & \dots & \sum_{r=|\alpha|+1}^M \tilde{w}_{M,M,r} \end{bmatrix}$$

and

$$\tilde{w}_{m+1,p,r} = \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r^\ell (-1)^{m+p+2-(\ell+r)} (r-1)! (p+\ell)! 2^{2(\ell+r)-2} \Gamma(r-\alpha+\ell+\frac{1}{2})}{(m+1-r)! 2r! [(r-\alpha)(p+1-\ell)] 2^\ell \Gamma(r-\alpha+\ell+2)}$$

Proof: From (), we have that,

$$U_m^*(2^k t - n) = \sum_{r=0}^{m+1} r (-1)^{m+1-r} \frac{(m+r)! 2^{2r-1}}{(m+1-r)! 2r!} (2^k t - n)^{r-1}$$

Also, we have that

$$D^\alpha U_m^*(2^k t - n) = n \leq |\alpha| \text{ and for } (n = |\alpha| + 1 \dots M) \text{ we have}$$

$$\begin{aligned} D^\alpha U_m^*(2^k t - n) &= \sum_{r=0}^{m+1} r (-1)^{m+1-r} \frac{(m+r)! 2^{2r-1}}{(m+1-r)! 2r!} D^\alpha (2^k t - n)^{r-1} \\ &= \sum_{r=|\alpha|+1}^{m+1} r (-1)^{m+1-r} \frac{(m+r)! 2^{2r-1} (r-1)! (2^k t - n)^{r-\alpha-1}}{(m+1-r)! 2r! [(r-\alpha)]} \end{aligned}$$

Now, approximate $(2^k t - n)^{r-\alpha-1}$ by $(M+1)$ - term of shifted second kind chebyshev wavelets, we have:

$$(2^k t - n)^{r-\alpha-1} = \sum_{p=0}^M d_{r-1,p} U_p^*(2^k t - n)$$

$$\begin{aligned} d_{r-1,p} &= \frac{8}{\pi} \int_0^1 (2^k t - n) U^*(2^k t - n) \sqrt{(2^k t - n) - (2^k t - n)^2} dt \\ &= \frac{8}{\pi} \sum_{\ell=0}^{p+1} \frac{\ell (-1)^{p+1-\ell} (p+\ell)! 2^{2\ell-1}}{(p+1-\ell)! 2^\ell [(r+\ell-\alpha)]} \int_0^1 \sqrt{(2^k t - n) - (2^k t - n)^2} (2^k t - n) dt \\ &= \frac{8}{\pi} \sum_{\ell=0}^{p+1} \frac{\ell (-1)^{p+1-\ell} (p+\ell)! 2^{2\ell-1} [(r-\alpha+\ell+\frac{1}{2})\sqrt{\pi}]}{(p+1-\ell)! 2^\ell [(r+\ell-\alpha+2)]} \end{aligned}$$

$$\begin{aligned} D^\alpha U_m^*(2^k t - n) &= \sum_{r=|\alpha|+1}^{m+1} \sum_{p=0}^M r (-1)^{m+1-r} \frac{(m+r)! 2^{2r-1} (r-1)!}{(m+1-r)! 2r! [(r-\alpha)]} \cdot d_{r-1,p} \cdot U_m^*(2^k t - n) \\ &= \sum_{p=0}^M \left[\sum_{r=|\alpha|+1}^{m+1} \tilde{w}_{n+1,p,r} \right] U_m^*(2^k t - n) \end{aligned}$$

$$D^\alpha \Psi_{nm}^2(t) = \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} \sum_{p=0}^M \left[\sum_{r=|\alpha|+1}^{m+1} \tilde{w}_{n+1,p,r} \right] U_m^*(2^k t - n), \text{ thus}$$

$$D^\alpha \Psi_{nm}^2(t) = \left[\sum_{r=|\alpha|+1}^{m+1} \tilde{w}_{m+1,0,r}, \sum_{r=|\alpha|+1}^{m+1} \tilde{w}_{m+1,1,r}, \dots, \sum_{r=|\alpha|+1}^{m+1} \tilde{w}_{m+1,M,r} \right] \cdot \Psi_{nm}^2(t)$$

$$\text{and } D^\alpha \Psi_{nm}^2(t) = \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} D^\alpha U_m^*(2^k t - n) = [0, 0, \dots, 0] \Psi_{nm}^2(t) \quad n \leq |\alpha|.$$

4. Operational Matrices of Fractional Derivative Chebyshev wavelets

The third and fourth – kind chebyshev wavelets $\Psi_{nm}^3(t) = \Psi_{nm}^4(t) = \Psi(k, n, m, t)$ has four argument, $k, n \in \mathbb{N}$, m is the order of the polynomial $V_m^*(t)$ or $W_m^*(t)$ and t is the normalized time . they are defined explicitly on the interval $[0,1]$ as: [1]

$\Psi_{nm}^3 = \frac{2^{\frac{k+1}{2}}}{\sqrt{\pi}} v_n^*(2^k t - n)$, $\Psi_{nm}^4 = \frac{2^{\frac{k+1}{2}}}{\sqrt{\pi}} w_n^*(2^k t - n)$ for $t \in [\frac{n}{2^k}, \frac{n+1}{2^k}]$, $m=0,1,\dots,M$, $n=0,1,\dots,2^k - 1$ and $\Psi_{nm}^3 = \Psi_{nm}^4 = 0$ otherwise. and the weight function:

$$w_1^* = \sqrt{\frac{(2^k t - n)}{1 - (2^k t - n)}}, \quad w_2^* = \sqrt{\frac{1 - (2^k t - n)}{(2^k t - n)}}$$

A function and the function $f(t)$ defined over $[0,1]$ may be expanded in terms of chebyshev wavelets as:

$$f(t) = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} c_{nm} \Psi_{nm}^{3,4}(t),$$

where

$$c_{nm} = \int_0^1 w_{1,2}^*(t) \cdot \Psi_{nm}^{3,4}(t) dt$$

$$f(t) = \sum_{n=0}^{2^k-1} \sum_{m=0}^M c_{nm} \Psi_{nm}^{3,4}(t)$$

$$c = [c_{00}, c_{01}, \dots, c_{0,M}, \dots, c_{2^k-1,1}, \dots, c_{2^k-1,M}]^T$$

$$\Psi_{nm}^{3,4}(t) = [\Psi_{0,0}^{3,4}, \Psi_{0,1}^{3,4}, \dots, \Psi_{2^k-1,M}^{3,4}, \dots, \Psi_{2^k-1,0}^{3,4}, \Psi_{2^k-1,1}^{3,4}, \dots, \Psi_{2^k-1,M}^{3,4}]^T$$

4.1. New Relation between Operational Matrices of Fractional Derivative for $\Psi_{nm}^2(t)$ and $\Psi_{nm}^3(t)$

$$\Psi_{nm}^2(t) = \begin{cases} \frac{2^{\frac{k+1}{2}}}{\sqrt{\pi}} U_m^*(2^k t - n) & t \in [\frac{n}{2^k}, \frac{n+1}{2^k}] \\ 0 & \text{w} \end{cases}$$

$$m=0, 1, \dots, m, n=0, 1, \dots, 2^k - 1, U_m^*(t) = \sqrt{\frac{2}{\pi}} U_m(t)$$

$$\Psi_{nm}^3(t) = \begin{cases} \frac{2^{\frac{k+1}{2}}}{\sqrt{\pi}} V_m^*(2^k t - n) & t \in [\frac{n}{2^k}, \frac{n+1}{2^k}] \\ 0 & \text{w} \end{cases}$$

$$m=0, 1, \dots, M, n=0, 1, \dots, 2^k - 1$$

$$V_m^*(t) = \frac{1}{\sqrt{\pi}} V_m(t)$$

$$V_m(2^k t - n) = U_m(2^k t - n) - U_{m-1}(2^k t - n)$$

$$\sqrt{\frac{2}{\pi}} V_m(2^k t - n) = \sqrt{\frac{2}{\pi}} U_m(2^k t - n) - \sqrt{\frac{2}{\pi}} U_{m-1}(2^k t - n)$$

$$\sqrt{2} V_m^*(2^k t - n) = U_m^*(2^k t - n) - U_{m-1}^*(2^k t - n)$$

$$\sqrt{2} \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} V_m^*(2^k t - n) = \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} U_m^*(2^k t - n) - \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} U_{m-1}^*(2^k t - n)$$

$$2\sqrt{2} \Psi_{nm}^3(t) = \Psi_{nm}^2(t) - \Psi_{nm-1}^2(t)$$

Theorem 4.1.1: Let $\Psi_{nm}^3(t)$ be third kind chebyshev vector and suppose $(\alpha > 0)$ then:

$$(\sum_{r=|\alpha|+1}^{m+1} W_{m+1,M,r}^{\sim} - \sum_{r=|\alpha|+1}^m W_{m,M,r}^{\approx}) D^{\alpha} \Psi_{nm}^3(t) = \frac{2^{\frac{k+3}{2}}}{2\sqrt{2}\sqrt{\pi}} D^{\alpha} (U_m^*(2^k t - n) - U_{m-1}^*(2^k t - n)) = \Delta^{\alpha} \Psi_{nm}^3(t)$$

where Δ^{α} is the $(M+1) \times (M+1)$ operational matrix derivative of order α in the caputo sense and defined as follow

$$\Delta^\alpha = \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 \\ \vdots & \vdots & \vdots \\ 0 & 0 & 0 \\ \left(\sum_{r=[\alpha]+1}^{[\alpha]+1} \tilde{w}_{[\alpha]+1,0,r} - \sum_{r=[\alpha]+1}^{[\alpha]+1} \tilde{w}_{[\alpha]+1,0,r} \right) & \left(\sum_{r=[\alpha]+1}^{[\alpha]+1} \tilde{w}_{[\alpha]+1,1,r} - \sum_{r=[\alpha]+1}^{[\alpha]+1} \tilde{w}_{[\alpha]+1,1,r} \right) \cdots & \left(\sum_{r=[\alpha]+1}^{[\alpha]+1} \tilde{w}_{[\alpha]+1,M,r} - \sum_{r=[\alpha]+1}^{[\alpha]+1} \tilde{w}_{[\alpha]+1,M,r} \right) \\ \vdots & \vdots & \vdots \\ \left(\sum_{r=[\alpha]+1}^{m+1} \tilde{w}_{m+1,0,r} - \sum_{r=[\alpha]+1}^m \tilde{w}_{m,0,r} \right) & \left(\sum_{r=[\alpha]+1}^{m+1} \tilde{w}_{m+1,1,r} - \sum_{r=[\alpha]+1}^m \tilde{w}_{m,1,r} \right) \cdots & \left(\sum_{r=[\alpha]+1}^{m+1} \tilde{w}_{m+1,M,r} - \sum_{r=[\alpha]+1}^m \tilde{w}_{m,M,r} \right) \\ \vdots & \vdots & \vdots \\ \left(\sum_{r=[\alpha]+1}^M \tilde{w}_{M,0,r} - \sum_{r=[\alpha]+1}^M \tilde{w}_{M,0,r} \right) & \left(\sum_{r=[\alpha]+1}^M \tilde{w}_{M,1,r} - \sum_{r=[\alpha]+1}^M \tilde{w}_{M,1,r} \right) \cdots & \left(\sum_{r=[\alpha]+1}^M \tilde{w}_{M,M,r} - \sum_{r=[\alpha]+1}^M \tilde{w}_{M,M,r} \right) \\ \vdots & \vdots & \vdots \end{pmatrix}$$

where

$$\tilde{w}_{m+1,p,r} = \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r \cdot \ell (-1)^{p+m+2-(\ell+r)} (r-1)! (m+r)! (p+\ell)! 2^{2(r+\ell)-2} \left[(r-\alpha+\ell+\frac{1}{2}) \right]}{(m+1-r)! 2r! [(r-\alpha)(p+1-\ell)! 2\ell! [(r+\ell-\alpha+2)]}$$

$$\tilde{w}_{m,p,r} = \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r \cdot \ell (-1)^{p+m+1-(\ell+r)} (r-1)! (m-1+r)! (p+\ell)! 2^{2(r+\ell)-2} \left[(r-\alpha+\ell+\frac{1}{2}) \right]}{(m-r)! 2r! [(r-\alpha)(p+1-\ell)! 2\ell! [(r+\ell-\alpha+2)]}$$

Note that in Δ^α , the first $[\alpha] + 1$ rows, are all zero.

Proof: From theorem(2.2), we have that

$$D^\alpha U_m^*(2^k t - n) = \sum_{r=[\alpha]+1}^{m+1} \sum_{p=0}^M \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r \cdot \ell (-1)^{p+m+2-(\ell+r)} (r-1)! (m+r)! (p+\ell)! 2^{2(r+\ell)-2} \left[(r-\alpha+\ell+\frac{1}{2}) \right]}{(m+1-r)! 2r! [(r-\alpha)(p+1-\ell)! 2\ell! [(r+\ell-\alpha+2)]} \text{ and}$$

$$D^\alpha U_{m-1}^*(2^k t - n) = \sum_{r=[\alpha]+1}^m \sum_{p=0}^M \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r \cdot \ell (-1)^{p+m+1-(\ell+r)} (r-1)! (m-1+r)! (p+\ell)! 2^{2(r+\ell)-2} \left[(r-\alpha+\ell+\frac{1}{2}) \right]}{(m-r)! 2r! [(r-\alpha)(p+1-\ell)! 2\ell! [(r+\ell-\alpha+2)]}$$

Then

$$D^\alpha U_m^*(2^k t - n) - D^\alpha U_{m-1}^*(2^k t - n) = \left(\sum_{p=0}^M \sum_{r=[\alpha]+1}^{m+1} \tilde{w}_{m+1,p,r} \right) U_m^*(2^k t - n) - \left(\sum_{p=0}^M \sum_{r=[\alpha]+1}^m \tilde{w}_{m,p,r} \right) U_m^*(2^k t - n),$$

Thus,

$$D^\alpha \Psi_{nm}^3(t) = \frac{2^{\frac{k+3}{2}}}{2\sqrt{2} \cdot \sqrt{\pi}} \sum_{p=0}^M \left[\sum_{r=[\alpha]+1}^{m+1} \tilde{w}_{m+1,p,r} - \sum_{r=[\alpha]+1}^m \tilde{w}_{m,p,r} \right] U_m^*(2^k t - n), \text{ then}$$

$$D^\alpha \Psi_{nm}^3(t) = \frac{1}{2\sqrt{2}} \left(\sum_{r=[\alpha]+1}^{m+1} \tilde{w}_{m+1,0,r} - \sum_{r=[\alpha]+1}^m \tilde{w}_{m,0,r} \right), \left(\sum_{r=[\alpha]+1}^{m+1} \tilde{w}_{m+1,1,r} - \sum_{r=[\alpha]+1}^m \tilde{w}_{m,1,r} \right), \dots, \left(\sum_{r=[\alpha]+1}^{m+1} \tilde{w}_{m+1,M,r} - \sum_{r=[\alpha]+1}^m \tilde{w}_{m,M,r} \right) \cdot \Psi_{nm}^3(t) \text{ and}$$

$$D^\alpha \Psi_{nm}^3(t) = \frac{2^{\frac{k+3}{2}}}{2\sqrt{2} \cdot \sqrt{\pi}} (D^\alpha U_m^*(2^k t - n) - D^\alpha U_{m-1}^*(2^k t - n))$$

$$= \frac{1}{2\sqrt{2}} [0, 0, \dots, 0] \Psi_{nm}^3(t) \quad n \leq [\alpha]$$

4.2. New Relation between Operational Matrices of Fractional Derivative for $\Psi_{nm}^2(t)$ and $\Psi_{nm}^4(t)$

$$\Psi_{nm}^2(t) = \begin{cases} \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} U_m^*(2^k t - n) & t \in \left[\frac{n}{2^k}, \frac{n+1}{2^k} \right] \\ 0 & o.w \end{cases}$$

$$m = 0, 1, \dots, M, n = 0, 1, \dots, 2^k - 1,$$

$$U_m^*(t) = \sqrt{\frac{2}{\pi}} U_m(t)$$

$$\Psi_{nm}^4(t) = \begin{cases} \frac{2^{\frac{k+1}{2}}}{\sqrt{\pi}} W_m^*(2^k t - n) & t \in \left[\frac{n}{2^k}, \frac{n+1}{2^k} \right] \\ 0 & o.w. \end{cases}$$

$$m = 0, 1, \dots, M, n = 0, 1, \dots, 2^k - 1,$$

$$W_m^*(t) = \frac{1}{\sqrt{\pi}} W_m(t)$$

$$W_m(2^k t - n) = U_m(2^k t - n) + U_{m-1}(2^k t - n)$$

$$\sqrt{\frac{2}{\pi}} W_m(2^k t - n) = \sqrt{\frac{2}{\pi}} U_m(2^k t - n) + U_{m-1}(2^k t - n)$$

$$\sqrt{2} W_m^*(2^k t - n) = U_m^*(2^k t - n) + U_{m-1}^*(2^k t - n)$$

$$\sqrt{2} \cdot \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} W_m^*(2^k t - n) = \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} U_m^*(2^k t - n) + \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} U_{m-1}^*(2^k t - n)$$

$$2\sqrt{2} \cdot \frac{2^{\frac{k+1}{2}}}{\sqrt{\pi}} W_m^*(2^k t - n) = \Psi_{nm}^2(t) - \Psi_{nm-1}^2(t)$$

$$2\sqrt{2} \Psi_{nm}^4(t) = \Psi_{nm}^2(t) - \Psi_{nm-1}^2(t)$$

Theorem 4.2.1: Let $\varphi_{nm}^4(t)$ be fourth kind chebyshev vector and suppose $\alpha > 0$, Then:

$$D^\alpha \Psi_{nm}^4(t) = \frac{2^{\frac{k+3}{2}}}{2\sqrt{2} \cdot \sqrt{\pi}} D^\alpha (U_m^*(2^k t - n) + U_{m-1}^*(2^k t - n)) = \Delta^{(\alpha)} \Psi_{nm}^4(t)$$

where $\Delta^{(\alpha)}$ is the $(M+1) \times (M+1)$ operational Matrix derivative of order α in the Caputoense and defined as follow:

$$\Delta^{(\alpha)} = \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & & 0 & & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & & 0 & & 0 \\ \left(\sum_{r=|\alpha|+1}^{|\alpha|+1} \tilde{w}_{|\alpha|+1,0,r} - \sum_{r=|\alpha|+1}^{|\alpha|+1} \tilde{w}_{|\alpha|+1,0,r} \right) & \dots & \left(\sum_{r=|\alpha|+1}^{|\alpha|+1} \tilde{w}_{|\alpha|+1,1,r} - \sum_{r=|\alpha|+1}^{|\alpha|+1} \tilde{w}_{|\alpha|+1,1,r} \right) & \dots & \left(\sum_{r=|\alpha|+1}^{|\alpha|+1} \tilde{w}_{|\alpha|+1,M,r} - \sum_{r=|\alpha|+1}^{|\alpha|+1} \tilde{w}_{|\alpha|+1,M,r} \right) \\ \vdots & & \vdots & & \vdots \\ \left(\sum_{r=|\alpha|+1}^{m+1} \tilde{w}_{m+1,0,r} - \sum_{r=|\alpha|+1}^m \tilde{w}_{m,0,r} \right) & \dots & \left(\sum_{r=|\alpha|+1}^{m+1} \tilde{w}_{m+1,1,r} - \sum_{r=|\alpha|+1}^m \tilde{w}_{m,1,r} \right) & \dots & \left(\sum_{r=|\alpha|+1}^{m+1} \tilde{w}_{m+1,M,r} - \sum_{r=|\alpha|+1}^m \tilde{w}_{m,M,r} \right) \\ \vdots & & \vdots & & \vdots \\ \left(\sum_{r=|\alpha|+1}^M \tilde{w}_{M,0,r} - \sum_{r=|\alpha|+1}^M \tilde{w}_{M,0,r} \right) & \dots & \left(\sum_{r=|\alpha|+1}^M \tilde{w}_{M,1,r} - \sum_{r=|\alpha|+1}^M \tilde{w}_{M,1,r} \right) & \dots & \left(\sum_{r=|\alpha|+1}^M \tilde{w}_{M,M,r} - \sum_{r=|\alpha|+1}^M \tilde{w}_{M,M,r} \right) \\ \vdots & & \vdots & & \vdots \end{pmatrix}$$

where

$$\tilde{w}_{m+1,p,r} = \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r \cdot \ell (-1)^{p+m+2-(\ell+r)} (r-1)! (m+r)! (p+\ell)! 2^{2(r+\ell)-2} \left[(r-\alpha+\ell+\frac{1}{2}) \right]}{(m+1-r)! 2r! [(r-\alpha)(p+1-\ell)! 2\ell! [(r+\ell-\alpha+2)]}$$

$$\tilde{w}_{m,p,r} = \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r \cdot \ell (-1)^{p+m+1-(\ell+r)} (r-1)! (m-1+r)! (p+\ell)! 2^{2(r+\ell)-2} \left[(r-\alpha+\ell+\frac{1}{2}) \right]}{(m-r)! 2r! [(r-\alpha)(p+1-\ell)! 2\ell! [(r+\ell-\alpha+2)]}$$

Note that in Δ^α , the first $[\alpha] + 1$ rows, are all zero .

4.3. New Relation between Operational Matrices of Fractional Derivative for $\Psi_{nm}^2(t)$, $\Psi_{nm}^3(t)$ and $\Psi_{nm}^4(t)$

$$\Psi_{nm}^2(t) = \begin{cases} \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} U_m^*(2^k t - n) & t \in \left[\frac{n}{2^k}, \frac{n+1}{2^k} \right] \\ 0 & o. w. \end{cases}$$

$$m = 0, 1, \dots, M, n = 0, 1, \dots, 2^k - 1, U_m^*(t) = \sqrt{\frac{2}{\pi}} U_m(t)$$

$$\Psi_{nm}^3(t) = \begin{cases} \frac{2^{\frac{k+1}{2}}}{\sqrt{\pi}} V_m^*(2^k t - n) & t \in \left[\frac{n}{2^k}, \frac{n+1}{2^k} \right] \\ 0 & o. w. \end{cases}$$

$$m = 0, 1, \dots, M, n = 0, 1, \dots, 2^k - 1, V_m^*(t) = \frac{1}{\sqrt{\pi}} V_m(t)$$

$$\Psi_{nm}^4(t) = \begin{cases} \frac{2^{\frac{k+1}{2}}}{\sqrt{\pi}} W_m^*(2^k t - n) & t \in \left[\frac{n}{2^k}, \frac{n+1}{2^k} \right] \\ 0 & O. W. \end{cases}$$

$$m = 0, 1, \dots, M, n = 0, 1, \dots, 2^k - 1, W_m^*(t) = \frac{1}{\sqrt{\pi}} W_m(t)$$

$$2U_m(2^k t - n) = V_m(2^k t - n) + W_m(2^k t - n)$$

$$\sqrt{2} \cdot \sqrt{\frac{2}{\pi}} U_m(2^k t - n) = \frac{1}{\sqrt{\pi}} V_m(2^k t - n) + \frac{1}{\sqrt{\pi}} W_m(2^k t - n)$$

$$\sqrt{2} \cdot \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} U_m^*(2^k t - n) = \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} V_m^*(2^k t - n) + \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} W_m^*(2^k t - n)$$

$$\frac{2^{\frac{k+3}{2}}}{\sqrt{2}\sqrt{\pi}} U_m^*(2^k t - n) = \frac{2^{\frac{k+1}{2}}}{\sqrt{\pi}} V_m^*(2^k t - n) + \frac{2^{\frac{k+1}{2}}}{\sqrt{\pi}} W_m^*(2^k t - n)$$

$$\frac{1}{\sqrt{2}} \Psi_{nm}^2(t) = \Psi_{nm}^3(t) + \Psi_{nm}^4(t)$$

Theorem 4.3.1: Let $\Psi_{nm}^2(t), \Psi_{nm}^3(t)$ and $\Psi_{nm}^4(t)$ are shifted second, third and fourth kind chebyshev vector respectively and suppose $\alpha > 0$, Then:

$$\begin{aligned} D^\alpha \Psi_{nm}^2(t) &= \Delta^{(\alpha)} \Psi_{nm}^2(t) = \frac{2^{\frac{k+1}{2}}}{2\sqrt{2} \cdot \sqrt{\pi}} D^\alpha (v_m^*(2^k t - n) + w_m^*(2^k t - n)) \\ &= \frac{1}{2\sqrt{2}} (\Delta^{(\alpha)} \varphi_{nm}^3(t) + \Delta^{(\alpha)} \Psi_{nm}^4(t)) \end{aligned}$$

where $\Delta^{(\alpha)}$ is the $(M+1) \times (M+1)$ operational Matrix derivative of order α in the Caputo derivative.

4.4. New Relation between Operational Matrices of Fractional Derivative for $\Psi_{\hat{n}m}^1(t)$ and $\Psi_{nm}^2(t)$

$$\Psi_{\hat{n}m}^1(t) = \begin{cases} \frac{2^{k/2}}{\sqrt{\pi}} T_m^*(2^k t - \hat{n}) & t \in \left[\frac{\hat{n}-1}{2^k}, \frac{\hat{n}}{2^k} \right] \\ 0 & o. w. \end{cases}$$

where

$$T_m^*(t) = \begin{cases} \frac{1}{\sqrt{\pi}} T_m(t) & m = 0 \\ \sqrt{\frac{2}{\pi}} T_m(t) & m > 0 \end{cases}$$

$$\Psi_{nm}^2(t) = \begin{cases} \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} U_m^*(2^k t - n) & t \in \left[\frac{n}{2^k}, \frac{n+1}{2^k} \right] \\ 0 & \text{o.w.} \end{cases}$$

$$U_m^*(t) = \sqrt{\frac{2}{\pi}} U_m(t)$$

$$2 T_m(2^k t - n) = U_m(2^k t - n) + U_{m-2}(2^k t - n) \quad m = 2, 3, \dots$$

$$2 \sqrt{\frac{2}{\pi}} T_m(2^k t - n) = \sqrt{\frac{2}{\pi}} U_m(2^k t - n) - \sqrt{\frac{2}{\pi}} U_{m-2}(2^k t - n)$$

$$2 T_m^*(2^k t - n) = U_m^*(2^k t - n) + U_{m-2}^*(2^k t - n)$$

$$2 \cdot \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} T_m^*(2^k t - n) = \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} U_m^*(2^k t - n) + \frac{2^{\frac{k+3}{2}}}{\sqrt{\pi}} U_{m-2}^*(2^k t - n)$$

$$c_m = \begin{cases} 1 & m = 0 \\ \sqrt{2} & m \neq 0 \end{cases}$$

$$\frac{4\sqrt{2}}{C_m} \Psi_{nm}^1(t) = \Psi_{nm}^2(t) - \Psi_{nm-2}^2(t) \quad m = 2, 3, \dots$$

Theorem 4.4.1: Let $\Psi_{nm}^1(t)$ be fourth kind chebyshev vector and suppose $\alpha > 0$, Then:

$$D^\alpha \Psi_{nm}^1(t) = \frac{C_m 2^{\frac{k+3}{2}}}{4\sqrt{2} \cdot \sqrt{\pi}} D^\alpha (U_m^*(2^k t - n) + U_{m-1}^*(2^k t - n)) = \Delta^{(\alpha)} \Psi_{nm}^2(t)$$

where $\Delta^{(\alpha)}$ is the $(M+1) \times (M+1)$ operational Matrix derivative of order α in the Caputo sense and defined as follow:

$$\Delta^{(\alpha)} = \begin{pmatrix} 0 & & 0 & & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & & 0 & & 0 \\ \left(\sum_{r=[\alpha]+1}^{[\alpha]+1} w_{[\alpha]+1,0,r} - \sum_{r=[\alpha]+1}^{[\alpha]+1} w_{[\alpha]+1,0,r} \right) & \left(\sum_{r=[\alpha]+1}^{[\alpha]+1} w_{[\alpha]+1,1,r} - \sum_{r=[\alpha]+1}^{[\alpha]+1} w_{[\alpha]+1,1,r} \right) & \dots & \left(\sum_{r=[\alpha]+1}^{[\alpha]+1} w_{[\alpha]+1,M,r} - \sum_{r=[\alpha]+1}^{[\alpha]+1} w_{[\alpha]+1,M,r} \right) \\ \vdots & \vdots & & \vdots \\ \left(\sum_{r=[\alpha]+1}^{m+1} w_{m+1,0,r} - \sum_{r=[\alpha]+1}^{m-1} w_{m-1,0,r} \right) & \left(\sum_{r=[\alpha]+1}^{m+1} w_{m+1,1,r} - \sum_{r=[\alpha]+1}^{m-1} w_{m-1,1,r} \right) & \dots & \left(\sum_{r=[\alpha]+1}^{m+1} w_{m+1,M,r} - \sum_{r=[\alpha]+1}^{m-1} w_{m-1,M,r} \right) \\ \vdots & \vdots & & \vdots \\ \left(\sum_{r=[\alpha]+1}^M w_{M,0,r} - \sum_{r=[\alpha]+1}^M w_{M,0,r} \right) & \left(\sum_{r=[\alpha]+1}^M w_{M,1,r} - \sum_{r=[\alpha]+1}^M w_{M,1,r} \right) & \dots & \left(\sum_{r=[\alpha]+1}^M w_{M,M,r} - \sum_{r=[\alpha]+1}^M w_{M,M,r} \right) \\ \vdots & \vdots & & \vdots \end{pmatrix}$$

$$= \frac{C_m}{4\sqrt{2}}$$

where

$$w_{m+1,p,r} = \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r \cdot \ell (-1)^{p+m+2-(\ell+r)} (r-1)! (m+r)! (p+\ell)! 2^{2(r+\ell)-2} \left[(r-\alpha+\ell+\frac{1}{2}) \right]}{(m+1-r)! 2r! [(r-\alpha)(p+1-\ell)! 2\ell! [(r+\ell-\alpha+2)]}$$

$$w_{m-2,p,r} = \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r \cdot \ell (-1)^{p+m-(\ell+r)} (r-1)! (m-2+r)! (p+\ell)! 2^{2(r+\ell)-2} \left[(r-\alpha+\ell+\frac{1}{2}) \right]}{(m-1-r)! 2r! [(r-\alpha)(p+1-\ell)! 2\ell! [(r+\ell-\alpha+2)]}$$

Note that in Δ^α , the first $[\alpha]+1$ rows, are all zero.

Proof:

$$D^\alpha U_m^*(2^k t - n) = \sum_{r=|\alpha|+1}^{m+1} \sum_{p=0}^M \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r \cdot \ell (-1)^{p+m+2-(\ell+r)} (r-1)! (m+r)! (p+\ell)! 2^{2(r+\ell)-2} \left[(r-\alpha+\ell+\frac{1}{2}) \right]}{(m+1-r)! 2r! [(r-\alpha)(p+1-\ell)! 2^\ell! [(r+\ell-\alpha+2)]} \text{ and}$$

$$D^\alpha U_{m-2}^*(2^k t - n) = \sum_{r=|\alpha|+1}^{m-2} \sum_{p=0}^M \frac{8}{\sqrt{\pi}} \sum_{\ell=0}^{p+1} \frac{r \cdot \ell (-1)^{p+m-(\ell+r)} (r-1)! (m-2+r)! (p+\ell)! 2^{2(r+\ell)-2} \left[(r-\alpha+\ell+\frac{1}{2}) \right]}{(m-1-r)! 2r! [(r-\alpha)(p+1-\ell)! 2^\ell! [(r+\ell-\alpha+2)]}$$

Then

$$D^\alpha U_m^*(2^k t - n) - D^\alpha U_{m-2}^*(2^k t - n) = \left(\sum_{p=0}^M \sum_{r=|\alpha|+1}^{m+1} w_{m+1,p,r} \right) U_m^*(2^k t - n) - \left(\sum_{p=0}^M \sum_{r=|\alpha|+1}^{m-1} w_{m-1,p,r} \right) U_m^*(2^k t - n),$$

Thus,

$$D^\alpha \Psi_{nm}^1(t) = \frac{C_m 2^{\frac{k+3}{2}}}{4\sqrt{2}\sqrt{\pi}} \sum_{p=0}^M \left[\sum_{r=|\alpha|+1}^{m+1} w_{m+1,p,r} - \sum_{r=|\alpha|+1}^{m-1} w_{m-1,p,r} \right] U_m^*(2^k t - n), \text{ then}$$

$$D^\alpha \Psi_{nm}^1(t) = \frac{C_m}{4\sqrt{2}} \left[\left(\sum_{r=|\alpha|+1}^{m+1} w_{m+1,0,r} - \sum_{r=|\alpha|+1}^{m-1} w_{m-1,0,r} \right), \left(\sum_{r=|\alpha|+1}^{m+1} w_{m+1,1,r} - \sum_{r=|\alpha|+1}^{m-1} w_{m-1,1,r} \right), \dots, \left(\sum_{r=|\alpha|+1}^{m+1} w_{m+1,M,r} - \sum_{r=|\alpha|+1}^{m-1} w_{m-1,M,r} \right) \right] \cdot \Psi_{nm}^2(t) \text{ and}$$

$$D^\alpha \Psi_{nm}^1(t) = \frac{C_m 2^{\frac{k+3}{2}}}{4\sqrt{2}\sqrt{\pi}} (D^\alpha U_m^*(2^k t - n) - D^\alpha U_{m-1}^*(2^k t - n)) = \frac{C_m}{4\sqrt{2}} [0, 0, \dots, 0] \Psi_{nm}^2(t) \quad n \leq [\alpha]$$

REFERENCES

- [1] Abd-Elhameed. W. M., Doha. E. H., and Youssri. Y. H., "New Wavelets Collocation Method for Solving Second-Order Multipoint Boundary Value problems Using Chebyshev Polynomials of Third and Fourth Kinds", Hindawi Publishing Corporation Abstract and Applied Analysis, Vol 2013, Article ID 542839,9 pages.
- [2] Abd-Elhameed. W. M., Doha. E. H., and Youssri. Y. H., "New Spectral Second Kind Chebyshev Wavelets Algorithm for Solving Linear and Nonlinear Second- Order Differential Equations Involving Singular and Bratu Type Equations", Hindawi Publishing Corporation Abstract and Applied Analysis, Vol 2013, Article ID 715756, 9 pages.
- [3] Azizi. H., Loghmani. G. B, "Numerical Approximation For space Fractional Diffusion equations Via Chebyshev Finite Difference Method", Journal of Fractional Calculus and Applications, vol. 4(2) July 2013, pp, 303-311.
- [4] Bhrawy.A.H., Alofi. A.S., "The operational matrix of fractional integration for shifted Chebyshev polynomials", Applied Mathematics Letters, 26(2013) 25-31.
- [5] Biazar. J., Ebrahimi. H., "A Strong Method for Solving System of Integro-Differential Equations, "Applied Mathematical, Vol.2,(2011), 1105-1113.
- [6] Chi.K& Liu. M., "A Wavelet approach to fast approximation of power Electronics circuits", International Journal of Circuit theory and Applications, 31(2003), 591-610.
- [7] Darani. M. A., Nasiri. M, "A fractional type of the Chebyshev polynomials for approximation of solution of linear fractional differential equations", Vol. 1, No. 2, 2013, pp. 96-107
- [8] Diethelm. K., "The analysis of fractional differential equations", Berlin: Springer-Verlag, 2010.
- [9] Doha. E. H., Bhrawy A.H, Ezz-Eldien S.S., "A Chebyshev spectral method based on operational matrix for initial and boundary value problems of fractional order", Comput Math Appl., 62(2011)2364-2373.
- [10] Fox, Lslie and Ian Bax Parker, "Chebyshev Polynomials in Numerical Analysis", Oxford university press, London, vol 29, 1968.
- [11] Jie. W.,Liu. M., "Application of Wavelet Transform to steady-state Approximation of power electronics Waveforms", Department of Electronic, Hong Kong polytechnic University, (2003).
- [12] Heydari. M.H.,Hooshmandasl. M.R., MaalekGhaini. F. M., and Mohammadi. F., "Wavelet Collocation Method for Solving Multiorder Fractional Differential Equations", Hindawipublishing Corporation, journal of applied mathematics, vol 2012/Article ID 542401, 19.

- [13] KazemiNasab., Kilicman A., pashazadehAtabakan. Z., and Abbasbandy. S., "Chebyshev Wavelet Finit Difference Method: A New Approach for Solving Initial and Boundary Value Problem of Fractional Order", Hindawi Publishing Corporation Abstract and Applied Analysis, vol 2013, Article ID 916456, 15 pages.
- [14] Lakestani.M., Dehghan.M., Irandoust-pakchin. S., "The construction of operational matrix of fractional derivatives using B-spline functions", Commun Nonlinear Sci Numer Simulat, 17 (2012) 1149-1162.
- [15] Li. X., "Numerical solution of fractional differential equations using cubic –spline wavelet collocation method, Commun Nonlinear Sci Numer Simulat., 17(2012) 3934-3946.
- [16] Miller. K.S., Ross. B., "An Introduction to the Fractional calculus and Fractional Differential Equations", Wiley, New York, 1993.
- [17] Nkwanta.A. and Barnes. E.R., "Two Catalan-type Riordan arrays and their connections to the Chebyshev polynomials of the first kind", Jornal of Integer Sequences, 15(2012)1-19.
- [18] Oldham. K.B.,Spanier. J., "The Fractional Calculus", Academic press. New York, 1974.
- [19] Podlubny. I., "Fractional Differential Equations". Academic Press, San Diego, 1999.
- [20] Saadatmandi. A., Dehghan. M., "A new operational matrix for solving fractional- order differential equations", Comput. Math. Appl., 59 (2010) 1326-1336.
- [21] Saadatmandi. A., Dehghan. M.,zizi. M R., "The Sine-Legendre collocatio n method for a class of fractional convection-diffusion equations with variable coefficients, Commun Nonliner Sci Numer Simulat., 17(2012)4125-4136.
- [22] Seifollahi. M.,Shamloo. A. S., "Numerical Solution of Nonlinear Multi-order Fractional Differential EQUATIONS BY Operational Matrix of Chebyshev Polynomials", World Applied Programming, Vol(3), Issue(3), March 2013. 85-92.
- [23] Sohrabi. S., "Comparison Chebyshev Wavelets Method with BPFs method For Solving Abel's Integral equation", in Shams Engineering Journal, Vol.2, (2011).

Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2014. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]