

**ADJACENT VERTEX SUM POLYNOMIAL**

**E. SUKUMARAN\***

Associate professor in Mathematics,  
 Lekshmpuram College of Arts & Science, Neyyoor, Kanyakumari District, Tamil Nadu, India.

(Received On: 10-05-15; Revised & Accepted On: 20-06-15)

**ABSTRACT**

Let  $G$  be a graph. The adjacent vertex sum polynomial is defined as  $S(G, x) = \sum_{i=0}^{\Delta(G)} n_{\Delta(G)-i} \cdot x^{\alpha_{\Delta(G)-i}}$  where  $\Delta(G) = \max \{ \deg v / v \in G \}$ ,  $n_{\Delta(G)-i}$  is the sum of the number of adjacent vertices of all the vertices of degree  $\Delta(G) - i$  and  $\alpha_{\Delta(G)-i}$  is the sum of the degree of adjacent vertices of all the vertices of degree  $\Delta(G) - i$ . In this paper I find the adjacent vertex sum polynomial of Cyclic graph, Complete graph, Generalized Peterson graph, Complete bipartite graph, Anti regular graph, Gear graph, Barbell graph and Book graph.

**2010 AMS Subject Classification:** 05 c x x

**Key words:** Maximum degree, Adjacent vertices, Cyclic graph, Complete graph, Generalised Peterson graph, Complete bipartite graph, Anti regular graph, Gear graph, Barbell graph and Book graph.

I define the adjacent vertex sum polynomial as follows.

Let  $G$  be a graph. The adjacent vertex sum polynomial is defined as  $S(G, x) = \sum_{i=0}^{\Delta(G)} n_{\Delta(G)-i} \cdot x^{\alpha_{\Delta(G)-i}}$  where  $\Delta(G) = \max \{ \deg v / v \in G \}$ ,  $n_{\Delta(G)-i}$  is the sum of the number of adjacent vertices of all the vertices of degree  $\Delta(G) - i$  and  $\alpha_{\Delta(G)-i}$  is the sum of the degree of adjacent vertices of all the vertices of degree  $\Delta(G) - i$ .

**Result: Adjacent vertex sum polynomial of Cyclic graph**

For cyclic graph  $C_n (n \geq 3)$  there are  $n$  vertices and each vertex is of degree 2 also for each vertex 2 vertices are adjacent vertices. Therefore the sum of the number of adjacent vertices of all the vertices of degree 2 is  $2n$  and the sum of the degree of adjacent vertices of all the vertices of degree 2 is  $4n$ .

Hence adjacent vertex sum polynomial of cyclic graph is  $S(C_n, x) = 2n x^{4n}$  for all  $n \geq 3$ .

**Result: Adjacent vertex sum polynomial of Complete graph**

For complete graph  $K_n (n \geq 1)$  there are  $n$  vertices and each vertex is of degree  $n - 1$  also for each vertex  $n - 1$  vertices are adjacent vertices. Therefore the sum of the number of adjacent vertices of all the vertices of degree  $n - 1$  is  $n(n - 1)$  and the sum of the degree of adjacent vertices of all the vertices of degree  $n - 1$  is  $(n - 1)^2 n$ .

Hence adjacent vertex sum polynomial of complete graph  $K_n$  is  $S(K_n, x) = n(n - 1) x^{(n-1)^2 n}$  for all  $n \geq 1$ .

**Result: Adjacent vertex sum polynomial of generalized Peterson graph**

For generalized Peterson graph  $P(n, k)$  there are  $2n$  vertices and  $3n$  edges. Also for this graph each vertex has degree 3. For this graph each vertex has 3 vertices are adjacent. Therefore the sum of the number of adjacent vertices of all the vertices of degree 3 is  $3 \times 2n = 6n$  and the sum of the degree of adjacent vertices of all the vertices of degree 3 is  $9 \times 2n = 18n$ . Hence adjacent vertex sum polynomial of generalized Peterson graph  $P(n, k)$  is  $S(P(n, k); x) = 6nx^{18n}$ .  
 $\forall n = 2, 3 \dots$

**Corresponding Author: Dr. E. Sukumaran\***

**Result: Adjacent vertex sum polynomial of Complete bipartite graph  $K_{m,n}$  ( $m \neq n$ )**

Note that the Complete bipartite graph  $K_{m,n}$  has  $m + n$  vertices and  $mn$  edges. Here  $m$  vertices have degree  $n$  and  $n$  vertices have degree  $m$ . Here for  $m$  vertices each has  $n$  vertices are adjacent. Therefore the sum of the number of adjacent vertices of all the vertices of degree  $n$  is  $mn$  and the sum of the degree of adjacent vertices of all the vertices of degree  $n$  is  $m^2n$ . Also here for  $n$  vertices each has  $m$  vertices are adjacent. Therefore the sum of the number of adjacent vertices of all the vertices of degree  $m$  is  $mn$  and the sum of the degree of adjacent vertices of all the vertices of degree  $m$  is  $mn^2$ . Hence adjacent vertex sum polynomial of complete bipartite graph  $K_{m,n}$  ( $m \neq n$ ) is

$$S(K_{m,n}; x) = mn x^{m^2n} + mn x^{mn^2} \quad \forall m \neq n$$

$$S(K_{m,n}; x) = mn (x^{m^2n} + x^{mn^2}) \quad \forall m \neq n$$

**Result: Adjacent vertex sum polynomial of antiregular graph  $A_{2n}$**

Vertex polynomial of antiregular graph  $A_{2n}$  is

$$V(A_{2n}; x) = x^{2n-1} + x^{2n-2} + \dots + x^{n+1} + 2x^n + x^{n-1} + \dots + x^3 + x^2 + x \quad \forall n = 1, 2, 3 \dots [1]$$

For  $A_{2n}$  the sum of the number of adjacent vertices of the vertex of degree  $2n - 1$  is  $2n - 1$  and the sum of the degree of adjacent vertices of the above vertex is  $(2n - 2) + (2n - 3) + \dots + (n + 1) + 2n + (n - 1) + \dots + 3 + 2 + 1$ . For  $A_{2n}$  the sum of the number of adjacent vertices of the vertex of degree  $2n - 2$  is  $2n - 2$  and the sum of the degree of adjacent vertices of the vertex of degree  $2n - 2$  is  $(2n - 1) + (2n - 3) + \dots + (n + 1) + 2n + (n - 1) + \dots + 3 + 2$ . For  $A_{2n}$  the sum of the number of adjacent vertices of the vertex of degree  $2n - 3$  is  $2n - 3$  and the sum of the degree of adjacent vertices of the vertex of degree  $2n - 3$  is  $(2n - 1) + (2n - 2) + (2n - 4) + \dots + (n + 1) + 2n + (n - 1) + \dots + 4 + 3$ . For  $A_{2n}$  the sum of the number of adjacent vertices of the vertex of degree  $n + 1$  is  $n + 1$  and the sum of the degree of adjacent vertices of the vertex of degree  $n + 1$  is  $(2n - 1) + (2n - 2) + \dots + (n + 3) + (n + 2) + 2n + (n - 1)$ . In  $A_{2n}$  there are two vertices of degree  $n$ . Therefore the sum of the number of adjacent vertices of the above two vertices of degree  $n$  is  $2n$  and the sum of the degree of adjacent vertices of the above two vertices of degree  $n$  is  $2(2n - 1) + 2(2n - 2) + 2(2n - 3) + \dots + 2(n + 1) + 2n$ . For  $A_{2n}$  the sum of the number of adjacent vertices of the vertex of degree  $n - 1$  is  $n - 1$  and the sum of the degree of adjacent vertices of the vertex of degree  $n - 1$  is  $(2n - 1) + (2n - 2) + \dots + (n + 2) + (n + 1)$ . For  $A_{2n}$  the sum of the number of adjacent vertices of the vertex of degree  $2$  is  $2$  and the sum of the degree of adjacent vertices of the vertex of degree  $2$  is  $(2n - 1) + (2n - 2)$ . For  $A_{2n}$  the number of adjacent vertex of the vertex of degree  $1$  is  $1$ . In this case there is only one vertex of degree  $2n - 1$  is adjacent to  $1$ .

$$S(A_{2n}; x) = (2n - 1) x^{(2n-2) + (2n-3) + \dots + (n+1) + 2n + (n-1) + \dots + 3 + 2 + 1}$$

$$+ (2n - 2) x^{(2n-1) + (2n-3) + \dots + (n+1) + 2n + (n-1) + \dots + 3 + 2}$$

$$+ (2n - 3) x^{(2n-1) + (2n-2) + (2n-4) + \dots + (n+1) + 2n + (n-1) + \dots + 4 + 3}$$

$$+ \dots + (n + 1) x^{(2n-1) + (2n-2) + \dots + (n+3) + (n+2) + 2n + (n-1)}$$

$$+ 2n x^{2(2n-1) + 2(2n-2) + 2(2n-3) + \dots + 2(n+1) + 2n} + (n - 1) x^{(2n-1) + (2n-2) + \dots + (n+1)}$$

$$+ \dots + 2x^{(2n-1) + (2n-2)} + x^{2n-1}$$

$$= (2n - 1) x^{2n^2 - 2n + 1} + (2n - 2) x^{2n^2 - 2n + 1} + (2n - 3) x^{2n^2 - 2n}$$

$$+ (2n - 4) x^{2n^2 - 2n - 2} + (2n - 5) x^{2n^2 - 2n - 5} + (2n - 6) x^{2n^2 - 2n - 9}$$

$$+ (2n - 7) x^{2n^2 - 2n - 14} + \dots + (n + 1) x^{2n^2 - 2n - \frac{n^2 - 5n + 4}{2}} + 2n x^{n(3n-1)}$$

$$+ (n - 1) x^{(2n-1) + (2n-2) + (2n-3) + \dots + (n+2) + (n+1)}$$

$$+ (n - 2) x^{(2n-1) + (2n-2) + (2n-3) + \dots + (n+3) + (n+2)}$$

$$+ (n - 3) x^{(2n-1) + (2n-2) + (2n-3) + \dots + (n+4) + (n+3)}$$

$$+ \dots + 3x^{(2n-1) + (2n-2) + (2n-3)} + 2x^{(2n-1) + (2n-2)} + x^{(2n-1)}$$

$$\therefore S(A_{2n}; x) = (2n - 1) x^{2n^2 - 2n + 1} + (2n - 2) x^{2n^2 - 2n + 1} + (2n - 3) x^{2n^2 - 2n}$$

$$+ (2n - 4) x^{2n^2 - 2n - 2} + (2n - 5) x^{2n^2 - 2n - 5} +$$

$$+ (2n - 6) x^{2n^2 - 2n - 9} + (2n - 7) x^{2n^2 - 2n - 14}$$

$$+ (n + 1) x^{2n^2 - 2n - \frac{n^2 - 5n + 4}{2}} + 2n x^{n(3n-1)}$$

$$+ (n - 1) x^{\sum_{i=1}^{n-1} (2n-i)} + (n - 2) x^{\sum_{i=1}^{n-2} (2n-i)} + (n - 3) x^{\sum_{i=1}^{n-3} (2n-i)}$$

$$+ \dots + 3x^{\sum_{i=1}^3 (2n-i)} + 3x^{\sum_{i=1}^2 (2n-i)} + x^{2n-1}$$

**Result: Adjacent vertex sum polynomial of antiregular graph  $A_{2n+1}$**

Vertex polynomial of antiregular graph  $A_{2n+1}$  is

$$V(A_{2n+1}; x) = x^{2n} + x^{2n-1} + x^{2n-2} + \dots + x^{n+1} + 2x^n + x^{n-1} + x^{n-2} + \dots + x^3 + x^2 + x \quad \forall n = 1, 2, 3 \dots [1]$$

Similar to previous argument adjacent vertex sum polynomial of  $A_{2n+1}$  is

$$S(A_{2n+1}; x) = 2nx^{(2n-1)+(2n-2)+\dots+(n+1)+2n+(n-1)+\dots+3+2+1} \\ + (2n-1)x^{2n+(2n-2)+(2n-3)+\dots+(n+1)+2n+(n-1)+(n-2)+\dots+3+2} \\ + (2n-2)x^{2n+(2n-1)+(2n-3)+\dots+(n+1)+2n+(n-1)+(n-2)+\dots+4+3} \\ + (2n-3)x^{2n+(2n-1)+(2n-2)+(2n-4)+\dots+(n+1)+2n+(n-1)+(n-2)+\dots+5+4} \\ + \dots + (n+1)x^{2n+(2n-1)+(2n-2)+\dots+(n+3)+(n+2)+2n} \\ + 2nx^{2 \times 2n + 2(2n-1) + 2(2n-2) + \dots + 2(n+2) + 2(n+1)} \\ + (n-1)x^{2n+(2n-1)+(2n-2)+\dots+(n+3)+(n+2)} \\ + \dots + 3x^{2n+(2n-1)+(2n-2)} + 2x^{2n+(2n-1)} + x^{2n}.$$

$$\therefore S(A_{2n+1}; x) = 2n x^{2n^2} + (2n-1)x^{2n^2} + (2n-2)x^{2n^2-1} + (2n-3)x^{2n^2-3} \\ + (2n-4)x^{2n^2-6} + (2n-5)x^{2n^2-10} + (2n-6)x^{2n^2-15} \\ + (n+1)x^{2n^2 - \frac{n^2-3n+2}{2}} + 2n x^{n(3n+1)} \\ + (n-1)x^{\sum_{i=0}^{n-2} (2n-i)} + (n-2)x^{\sum_{i=0}^{n-3} (2n-i)} + (n-3)x^{\sum_{i=0}^{n-4} (2n-i)} \\ + \dots + 3x^{\sum_{i=0}^2 (2n-i)} + 2x^{\sum_{i=0}^1 (2n-i)} + x^{2n}.$$

**Result: Adjacent vertex sum polynomial of Gear graph**

Vertex polynomial of Gear graph  $G_n$  is

$$V(G_n; x) = x^{n-1} + (n-1)x^3 + (n-1)x^2 \quad \forall n = 2, 3, \dots [1]$$

For Gear graph  $G_n$  has  $2n - 1$  vertices and  $3(n - 1)$  edges. In this graph one vertex has  $n - 1$  vertices are adjacent. Therefore the sum of the number of adjacent vertices of the above vertex is  $n - 1$  and the sum of the degree of adjacent vertices of the above vertex is  $3(n - 1)$ . For  $G_n$  the sum of the number of adjacent vertices of all the vertices of degree 3 is  $3(n - 1)$  and the sum of the degree of adjacent vertices of all the vertices of degree 3 is  $(n - 1)(n + 3)$ . For  $G_n$  the sum of the number of adjacent vertices of all the vertices of degree 2 is  $2(n - 1)$  and the sum of the degree of adjacent vertices of all the vertices of degree 2 is  $6(n - 1)$ . Hence adjacent vertex sum polynomial of gear graph is  $S(G_n, x) = (n - 1)x^{3(n-1)} + 3(n - 1)x^{(n-1)(n+3)} + 2(n - 1)x^{6(n-1)} \quad \forall n = 2, 3, \dots$

**Result: Adjacent vertex sum polynomial of Barbell graph**

Vertex polynomial of Barbell graph  $B_n$  is  $V(B_n; x) = 2[x^n + (n - 1)x^{n-1}] \quad \forall n = 1, 2, 3, \dots [1]$

For Barbell graph  $B_n$  has  $2n$  vertices and  $n^2 - n + 1$  edges. In this graph each vertex of degree  $n$  is adjacent to  $n$  vertices. Therefore the sum of the number of adjacent vertices of all the vertices of degree  $n$  is  $2n$  and the sum of the degree of adjacent vertices of all the vertices of the degree  $n$  is  $2(n^2 - n + 1)$ . For  $B_n$  the sum of the number of adjacent vertices of all the vertices of degree  $n - 1$  is  $2(n - 1)$  and the sum of the degree of adjacent vertices of all the vertices of degree  $n - 1$  is  $2(n - 1)[n + (n - 1)(n - 2)]$ . Hence adjacent vertex sum polynomial of Barbell graph is

$$S(B_n, x) = 2nx^{2(n^2-n+1)} + 2(n-1)^2 x^{2(n-1)[n+(n-1)(n-2)]} \quad \forall n = 1, 2, 3, \dots$$

**Result: Adjacent vertex sum polynomial of Book graph.**

Vertex polynomial of Book graph  $B'_n$  is

$$V(B'_n; x) = 2x^{n+1} + 2nx^2 \quad \forall n = 1, 2, 3, \dots [1]$$

For Book graph  $B'_n$  has  $2(n+1)$  vertices and  $3n + 1$  edges. In this graph 2 vertices have degree  $n + 1$ . Therefore the sum of the number of adjacent vertices of the above two vertices is  $2(n+1)$  and the sum of the degree of adjacent vertices of the above two vertices is  $2(3n+1)$ . For  $B'_n$  the sum of the number of adjacent vertices of all the vertices of degree 2 is  $4n$  and the sum of the degree of adjacent vertices of all the vertices of degree 2 is  $2n(n+3)$ . Hence adjacent vertex sum polynomial of Book graph is

$$S(B'_n; x) = 2(n+1)x^{2(3n+1)} + 4nx^{2n(n+3)} \quad \forall n = 2, 3, 4, \dots$$

## REFERENCES

1. Devaraj, J and Sukumaran, E., On Vertex Polynomial, International Journal of Math.Scis and Engg. Applications, IJMSEA, 6(V) (January 2012), 371-380.
2. Devaraj, J and Sukumaran, E., Vertex Polynomial of well known graphs and newly constructed graph and their vertex polynomial, International Journal of Math. Scis and Engg. Applications, IJMSEA Vol. 6 No. III (May 2012), 443-459.
3. Emanuele, Munarini, Characteristic, Admittance and matching Polynomials of an Antiregular Graph, Applicable Analysis and Discrete Mathematics (19<sup>th</sup> Dec 2008).
4. Gutman, I., A real graph Polynomial? Graph Th. Notes, New York, 22 (1992), 33 – 37.
5. Harary, F., Graph Theory, Narose Publishing House (1969).

**Source of support: Nil, Conflict of interest: None Declared**

***[Copy right © 2015. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]***