

A STUDY ON THE DOMINATING TRANSVERSAL CHROMATIC NUMBER

P. S. S. R. SUJATHA*

Department of Mathematics, Sri Aditya Engineering College, Surampalem, India.

P. S. BALARAM

Department of Management Studies, Aditya P.G. College, Kakinada, India.

(Received On: 15-04-15; Revised & Accepted On: 21-07-15)

ABSTRACT

Domination is a fast developing area in Graph Theory. The concept of domination leads to very interesting research work with good applications.


In this paper three areas of Graph Theory namely Domination, Coloring and Transversal theory are combined judiciously to define a new domination parameter. A result on std-set is given which can be used to verify whether the given std-set is minimal or not. Also, γ_{st} i.e., the dominating transversal chromatic number for different graphs such as Petersen graph, Wheel, Herschel graph and Grotzsch graph are given.

Key words: *Dominating colour transversal set, χ -partition, Herschel graph, Grotzsch graph, Petersen Graph and Wheel.*

BASIC DEFINITIONS AND RESULTS

Definition 1.1: Let $G = (V, E)$ be a graph. A non-empty set $D \subseteq V$ is called a dominating set of G if every vertex in $V-D$ is adjacent to some vertex of D . Let \hat{C} be the collection of all dominating sets of G . The member of \hat{C} whose cardinality is minimum is called a γ -set and its cardinality is defined to be the domination number of G and it is denoted by γ .

Example: In the following graph, The Red Vertices denote the Dominating sets.



Definition 1.2: A partition of the vertex set V into a minimum number of disjoint equivalence classes of independent sets, called colour classes, is said to be a χ -partition of G and this minimum number is denoted by χ .

Definition 1.3: A dominating set is called a dominating colour transversal set (std-set) if it has a non-empty intersection with every colour class of some χ -partition of G . This set is called an std-set because it is transversal of at least one (single) χ -partition of G .

Definition 1.4: An std-set with minimum cardinality is called a γ_{st} -set and its cardinality is denoted by γ_{st} .

Definition 1.5: Let Π be a χ -partition of a graph G . A set $D \subseteq V$ is said to be a transversal of Π if D intersects every colour class of Π .

Corresponding Author: P. S. S. R. Sujatha
 Department of Mathematics, Sri Aditya Engineering College, Surampalem, India.*

Definition 1.6: A dominating set $D \subseteq V$ is called a dominating colour transversal set (*std-set*) of a graph G if D is a transversal of at least one χ -partition of G . We call this set an *std-set* because it is a dominating colour transversal of at least one (single) χ -partition. An *std-set* is minimal if none of its proper subsets is an *std-set*.

Definition 1.7: Let $\hat{C} = \{D \subseteq V \mid D \text{ is an } \text{std-set of } G\}$. \hat{C} is non empty as $v \in \hat{C}$. Let $D^1 \in \hat{C}$ whose cardinality is minimum. D^1 is called a γ_{st} -set and its cardinality is called the dominating colour transversal number denoted by γ_{st} .

Result 1.8: For any graph G , $1 \leq \gamma \leq \gamma_{st}$

Result 1.9: For any Bipartite graph G , γ_{st} is either γ or γ_{+1} since by adding if necessary a single vertex to a γ_{st} -set, we get a γ_{st} -set for G .

Theorem 1.10: For any *std-set* D is minimal \Leftrightarrow for every $u \in D$, any of the following holds:

- (a) u is an isolate of D
- (b) There exist a vertex $v \in V - D \ni N(v) \cap D = \{u\}$.
- (c) For every χ -partition, $\Pi = \{V_1, V_2, V_3, \dots, V_k\}$, There exist one $V_i \ni V_i \cap D = \{u\}$ or \emptyset .

Proof: Let D be an *std-set*.

Suppose D is minimal.

Then $D - \{u\}$ is not an *std-set* for every $u \in D$.

\Rightarrow either $D - \{u\}$ is not a dominating set or not a transversal of every χ -partition of G .

Case-1: Suppose $D - \{u\}$ is not a dominating set.

Then there exists a vertex $v \in (V - D) \cup \{u\}$ that is not adjacent to any vertex of $D - \{u\}$.

If $u = v$, then u is an isolate of D .

If $u \neq v$, then v is adjacent to u but not to any other vertex of D .

Hence $N(v) \cap D = \{u\}$.

Case-2: Suppose $D - \{u\}$ is not a transversal for every χ -partition of $\{V_1, V_2, V_3, \dots, V_k\}$.

$\Rightarrow D - \{u\} \cap V_i = \emptyset$, for some i .

i.e., $V_i \cap D = \{u\}$ or \emptyset for some i

Hence (c) is satisfied.

Conversely, assume any one of the three conditions.

We prove that D is a minimal *std-set*.

Suppose D is not a minimal *std-set*.

Then D is an *std-set* but not minimal.

$\Rightarrow D$ and $D - \{u\}$ are *std-sets* for some $u \in D$.

Let $\{V_1, V_2, V_3, \dots, V_k\}$ be a χ -partition of V for which $D - \{u\}$ and D are transversals.

Then $D - \{u\} \cap V_i \neq \emptyset$ and $D \cap V_i \neq \emptyset$ for every i .

$\Rightarrow D \cap V_i \neq \{u\}$ or \emptyset , contradicting condition (c)

Result 1.11: For any graph G , $\gamma \leq \gamma_g \leq \gamma_{st}$.


Result 1.12: If G has K -components G_1, G_2, \dots, G_k such that $\chi(G_1) \geq \chi(G_i)$, for $i=1, 2, \dots, k$, then $\gamma_{st(G)} \leq \gamma_{st(G_1)} + \sum_{i=2}^k \gamma(G_i)$.

THE DOMINATING COLOUR TRANSVERSAL NUMBER FOR SOME STANDARD GRAPHS

1. PETERSEN GRAPH

In the mathematical field of graph theory, the **Petersen graph** is an undirected graph with 10 vertices and 15 edges. It is a small graph that serves as a useful example and counterexample for many problems in graph theory. The Petersen graph is named for Julius Petersen, who in 1898 constructed it to be the smallest bridgeless cubic graph with no three-edge-coloring.


For the Petersen graph P , $\gamma_{st} = 4$.



2. HERSCHEL GRAPH

In graph theory, a branch of mathematics, the **Herschel graph** is a bipartite undirected graph with 11 vertices and 18 edges, the smallest non-Hamiltonian polyhedral graph. It is named after British astronomer Alexander Stewart Herschel, who wrote an early paper concerning William Rowan Hamilton's icosian game: the Herschel graph describes the smallest convex polyhedron for which this game has no solution. However, Herschel's paper described solutions for the icosian game only on the graphs of the regular tetrahedron and regular icosahedron; it did not describe the Herschel graph.


The Herschel graph H as shown below is bipartite since it contains no odd cycles. Hence $\chi(H) = 2$ and $\gamma_{st}=3$.



3. GROTZSCH GRAPH

In the mathematical field of graph theory, the **Grötzsch graph** is a triangle-free graph with 11 vertices, 20 edges. It is named after German mathematician Herbert Grötzsch.


G_4 is a triangle free 4-chromatic graph we can easily verify that $\gamma_{st} = 4$.



4. WHEEL GRAPH

In the mathematical discipline of graph theory, a **wheel graph** W_n is a graph with n vertices ($n \geq 4$), formed by connecting a single vertex to all vertices of an $(n-1)$ -cycle. The numerical notation for wheels is used inconsistently in the literature: some authors instead use n to refer to the length of the cycle, so that their W_n is the graph we denote W_{n+1} .^[1] A wheel graph can also be defined as the 1-skeleton of an $(n-1)$ -gonal pyramid.

$$\gamma_{st}(W_p) = \begin{cases} 3, & \text{if } p \text{ is odd} \\ 4, & \text{if } p \text{ is even} \end{cases}$$



APPLICATIONS

1. Dominating sets are useful in routing problems. How many internet routers do you need so that every computer in your business has internet access?
2. It is also useful in scheduling problems. Suppose that some people are attending a conference. Each person writes down a list of events they want to attend. What is the smallest number of time slots needed so that everyone can attend all sessions that they want to attend (assuming there are enough rooms)? Make a graph whose vertices are the different events, with an edge between two vertices x and y if there is at least one person who wants to attend events x and y . The smallest number of events is the chromatic number of the graph.
3. Suppose you have hazardous chemicals that cannot be stored together. To find the minimum number of storage units you need to store all your chemicals?
4. suppose you have a set of volunteers and certain volunteers who won't work with each other. How do you group them to minimize the number of groups

REFERENCES

1. Math. Phy. Sci. 14 (1980), 471–475.
2. S. Arumugam and R. Kala, Domsaturation number of a graph, Ind. J.Pure. Appl. Math. 33 (2002), 1671–1676.
3. B.D. Acharya, The Strong Domination Number of a graph and related concepts, Jour. Connected domsaturation number of a graph, Ind. J. Pure. Appl. Math. 35 (2004), 1215–1221.
4. R. Balakrishnan and K. Ranganathan, A Textbook of Graph Theory, Springer, New York, 2000.
5. C. Berge, Graphs and Hypergraphs, North-Holland, Amsterdam, 1973.6. G. Chartrand and L. Lesniak, Graphs and Digraphs, Chapman and Hall, CRC, 2005.
6. G. Chartrand and Pingzhang, Introduction to Graph Theory, Tata Mc-Graw Hill, 2006.
7. E.J. Cockayne and S.T. Hedetniemi, Towards a Theory of Domination in Graphs, Networks 7 (1977), 247–261.
8. Total Domination in Graphs, Networks 10 (1980), 211 –219.
9. G.S. Domke, J.E. Dunbar, and L. Markus, Gallai-type theorems and domination parameters, Disc. Maths., To appear.
10. J.E. Dunbar, T.W. Haynes, and M.A. Henning, The Domatic Number of a Graph and its Complement, To appear.
11. J.F. Fink, M.S. Jacobson, L.F. Kinch, and J. Roberts, The Bondage Number of a Graph, Disc. Math. 86 (1990), 47–57.
12. F. Harary, Graph Theory, Addison-Wesley, Reading, Mass, 1969.
13. T.W. Haynes, S.T. Hedetniemi, and P.J. Slater, Domination in Graphs: Advanced Topics, Marcel Dekker Inc., 1997.
14. Fundamentals of Domination in Graphs, Marcel Dekker Inc.,
15. S.T. Hedetniemi and R. Laskar, Connected Domination in Graphs,
16. Graph Theory and Combinatorics (B. Bollobas, ed.), Academic Press, 1984, pp. 209–218.
17. F. Jaeger and C. Payan, Relations du type Nordhaus Gaddum pour lenombre d'absorption d'un graphe simple, vol. 274, pp. 728–730, C.R.Acad. Sci., Paris A, 1972.
18. S.R. Jayaram, The Domatic Number of a Graph, Nat. Acad. Sci. Lett.10 (1987), 23–25.
19. L. Mirsky, Transversal Theory, Academic Press, New York, 1971.
20. E.A. Nordhaus and J.W. Gaddum, On Complementary Graphs, Amer. Math. Monthly 63 (1956), 175–177.
21. O. Ore, Theory of Graphs, vol. 38, Amer. Math. Soc. Colloq. Publ., Providence, 1962.
22. J. Paulraj Joseph and S. Arumugam, On the Connected Domatic Number of a Graph, J. Ramanujan Math. Soc. 9 (1994), 69–77.
23. <http://en.wikipedia.org/wiki/Petersengraph>
24. [wikipedia.org/wiki/Grötzsch graph](http://en.wikipedia.org/wiki/Grötzsch_graph).

Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2015. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]