

A NOTE ON B_1 -NEAR-FIELD SPACES OVER REGULAR δ -NEAR-RINGS (B_1 - NFS-R- δ -NR)

Dr. N. V. NAGENDRAM*

Professor, Department of Mathematics(S&H)
Kakinada Institute of Technology & Science (KITS)
Tirupathi (v), Divili 533 433, East Godavari District, Andhra Pradesh. India.

(Received On: 10-08-15; Revised & Accepted On: 31-08-15)

ABSTRACT

The aim of Dr N V Nagendram in this paper is to extend the concepts of B_1 near-rings and after in depth study to introduce the concepts of B_1 - near-field spaces and strong B_1 - near-field spaces. We say that a right near-field space N is a B_1 - near-field space if for every $a \in N$, there exists $x \in N^*$ where $N^* = N - \{0\}$, such that $Nax = Nxa$. By way of generalization, we define N as a strong B_1 - near-field space if $Nab = Nba \forall a, b \in N$. Dr. N. V. Nagendram discuss some of their properties and obtain a characterization and also a structure theorem.

Keywords: B_1 -near-ring, strong B_1 -near-ring, near-field, near-field space, sub near-field space, B_1 -near-field spaces, B_1 - sub near-field spaces.

Subject Classification Code: MSC (2010):16D25, 54G05, 54C40, 16Y30.

SECTION 1: INTRODUCTION

Throughout, this paper N stands for a right near-field space $(N, +, \cdot)$, with at least two elements and “0” denotes the identity element of the near-field $(N, +)$. Obviously $0n = 0 \forall n \in N$. If in addition, $n0 = 0 \forall n \in N$ then we say that N is zero symmetric near-field. A sub near-field space $(M, +)$ of $(N, +)$ is called an N - sub near-field space of N if $NM \subset M$ and an invariant N - sub near-field space of N if $MN \subset M$ as well. N is called weak commutative near-field space if $abc = acb \forall a, b, c \in N$ (by G Pilz [4]). N is said to be regular near-field space if for every $a \in N$ there exists $b \in N$ such that $a = aba$. An element a is said to be nilpotent if $a^k = 0$ for some positive integer k . N is called nil near-field space if every element of N is nilpotent. N is called integral near-field space if N has no non-zero zero divisors. N is called a P_k near-field space if there exists a positive integer k such that $x^k N = x N x \forall x \in N$. For any sub near-field space A of near-field space N , we denote by A^* the set of all non-zero elements of A . In particular $N^* = N - \{0\}$. N is called a strong B_1 near-field space if $N^* = N_{B_1(a)} \forall a \in N$ where $N_{B_1(a)} = \{x \in N^* / axa = xa\}$. For basic concepts and terms used but left undefined in this paper we refer to Gunter Pilz [4].

Near-field over a regular delta Near-Ring is a generalized structure of a near-field. Dr. N.V. Nagendram introduced the notion of B_1 - near-field spaces over a regular delta near-ring.

Dr. N.V. Nagendram after in depth study of near-rings, regular near-rings, regular δ -near-ring, various types of regular δ -near-rings, near-fields, semi near-fields and near-field space over regular delta near-rings extended the concept to B_1 - near-field spaces over a regular delta near-ring and characterization of B_1 - near-field spaces over a regular delta near-ring.

As in near-ring theory it is interesting to fuzzify some sub structures of near-field spaces over regular delta near-rings of a near-ring. Hence our aim in this paper is to study B_1 - sub-near-field spaces and B_1 - near-field spaces.

The organization of this paper is as follows:

In section 2, some fundamental and preliminary definitions and results are given.

In section 3, B_1 - near-field spaces of a near-field over regular delta near-ring are defined and obtained some results in relation B_1 - near-field spaces.

*Corresponding Author: Dr. N. V. Nagendram**

In section 4, Study of Strong B_1 – near-field spaces of a near-field over regular delta near-ring and are defined. Also obtain some results in relation Strong B_1 – near-field spaces.

Section 5 concludes the paper. In depth study of various types of near-fields [10], Dr. N V Nagendram extended the B_1 – near-fields and strong B_1 -near-fields over regular delta near-ring in a near-field space by applying the topological applications making it into algebraic topology in modern and abstract algebra for large scale scope of applications in algebra of mathematics. Algebraic topology is a branch of mathematics that uses tools from abstract algebra to study topological spaces. The basic goal is to find algebraic invariants that classify topological spaces up to homeomorphism, though usually most classify up to homotopy equivalence. Although algebraic topology primarily uses algebra to study topological problems, using topology to solve algebraic problems is sometimes also possible. Algebraic topology, for example, allows for a convenient proof that any subgroup of a free group is again a free group.

SECTION 2: FUNDAMENTAL DEFINITIONS AND RESULTS

In this section we recall some fundamental definitions and results for the sake of completeness in the sense of topology:

2.1 Definition: zero symmetric near-field. N stands for a right near-field space $(N, +, \cdot)$, with at least two elements and “0” denotes the identity element of the near-field $(N, +)$.

Obviously $0n = 0 \forall n \in N$. If in addition, $n0 = 0 \forall n \in N$ then we say that N is zero symmetric near-field.

2.2 Definition: sub near-field space. A sub near-field space $(M, +)$ of $(N, +)$ is called an N - sub near-field space of N if $NM \subset M$ and an invariant N - sub near-field space of N if $MN \subset M$ as well.

2.3 Definition: weak commutative near-field space [G Pilz4]. N is called weak commutative near-field space if $abc = acb \forall a, b, c \in N$.

2.4 Definition: regular near-field space. N is said to be regular near-field space if for every $a \in N$ there exists $b \in N$ such that $a = aba$.

2.5 Definition: nil near-field space. An element a is said to be nilpotent if $a^k = 0$ for some positive integer k .

N is called nil near-field space if every element of N is nilpotent.

2.6 Definition: Integral near-field space. N is called integral near-field space if N has no non-zero zero divisors.

2.7 Definition: P_k near-field space. N is called a P_k near-field space if there exists a positive integer k such that $x^k N = x N x \forall x \in N$.

Here is an extension of the results from [3], [4] and [5] mentioned remarks as below:

2.8 Result: a B_1 – near-field space N has no non-zero nilpotent elements $\Leftrightarrow a^2 = 0 \Rightarrow a = 0 \forall a \in N$.

2.9 Result: a B_1 – near-field space N is zero symmetric \Leftrightarrow left ideal of B_1 – N is an B_1 – N -sub near-field space of N .

2.10 Result: Let a B_1 – near-field space N be zero symmetric then the following are equivalent:

- (i) a B_1 – near-field space N has no non-zero nilpotent elements
- (ii) a B_1 – near-field space N is a sub direct product of integral near-fields say N_i for all $i = 1, 2, 3, \dots$

2.11 Result: If N is a strong B_1 – near-field space then N is zero symmetric.

2.12 Result: N is a strong B_1 – near-field space $\Leftrightarrow axa = xa \forall a, x \in N$.

SECTION 3: B_1 - NEAR-FIELD SPACES OVER REGULAR DELTA NEAR - RINGS

In this section, I discuss about a right near-field space N is said to be left bi-potent if $Na = Na^2 \forall a \in N$.

3.1 Definition: B_1 near-field space. A right near-field space N is said to be left nilpotent if $Na = Na^2 \forall a \in N$. Then N is a B_1 – near-field space if for every $a \in N$, there exists $x \in N^*$ such that $Nax = Nxa$.

3.2 Example: Every constant near-field space is a B_1 – near-field space.

3.3 Example: The near-field space $(Z_4, +, \cdot)$ where $(Z_4, +)$ is the group of integers modulo “4” and “ \cdot ” is defined as follows (Scheme (4), p.407 of Gunter Pilz [4]).

.	0	1	2	3
0	0	0	0	0
1	0	0	1	0
2	0	0	3	0
3	0	0	2	0

This is a B_1 - near-field space over a regular δ -near-ring.

3.4 Theorem: Let N be a near-field space. Each of the following implies that N is a B_1 - near-field space over a regular δ -near-ring.

- (i) N is a zero symmetric nil near-field space over a regular δ -near-ring.
- (ii) N is a weak commutative.
- (iii) N has identity “1”.
- (iv) N is a near-field space over a regular δ -near-ring.

Proof: To prove (i): Let $a \in N$. if $a = 0$, then $\forall x \in N^*$, $Nax = Nxa = N0 = \{0\}$. If $a \in N^*$, since N is nil near-field space over a regular δ -near-ring, there exists a positive integer k such that $a^k = 0$. Put $x = a^{k-1} \neq 0$.

Now $Nax = Naa^{k-1} = Na^k = Na^{k-1}a = Nxa = N0 = \{0\}$, Thus N is a B_1 - near-field space over a regular δ -near-ring.

To prove (ii): Let $a \in N$. $\forall x \in N^*$, $y \in Nax \Rightarrow y = nax$ where $n \in N$. Since N is weak commutative, $y = nxa \in Nxa$. Therefore $Nax \subset Nxa$. Similarly we can show that $Nxa \subset Nax$ and (ii) follows.

To prove (iii): Follows by taking $x = 1$ in the definition 3.1.

To prove (iv): Follows from (iii).

This completes the proof of the theorem.

3.5 Theorem: Let N be a B_1 - near-field space over a regular δ -near-ring. If N is a strong B_1 - near-field space over a regular δ -near-ring without non-zero zero divisors then the following are true:

- (i) Every non-zero N -sub near-field space over a regular δ -near-ring of N is a B_1 - near-field space over a regular δ -near-ring.
- (ii) Every non-zero ideal of near-field space over a regular δ -near-ring N is an B_1 - near-field space over a regular δ -near-ring.

Proof: Since N is a strong B_1 - near-field space over a regular δ -near-ring, by result 2.11 N is zero symmetric near-field space and by result 2.12

$$aba = ba \quad \forall a, b, c \in N \tag{1}$$

(i) Let M be an N - sub near-field space of N and let $m \in M$. If $m = 0$ then $\forall x \in N^*$, $Nmx = N0 = \{0\}$ (Since N is zero symmetric near-field space) = Nxm . For $m \neq 0$, since N is a B_1 - near-field space over a regular δ -near-ring, $\exists y \in N^*$ such that $Nm = Nym$ (2)

Let $n = ym$.

$$\Rightarrow n \in M^*. \text{ Now } Mmn = Mm(ym) \subset Nm(ym) \subset (Nmy)m = (Nym)m \text{ [by (2)]}$$

$$\Rightarrow \hspace{10em} = N(mym)m \text{ [by (1)]}$$

$$\Rightarrow \hspace{10em} = Nm(ym)m \subset M(ym)m = Mnm.$$

$$\text{i.e., } Mmn \subset Mnm \tag{3}$$

$$\text{In a similar fashion we get } Mnm \subset Mmn \tag{4}$$

(3) and (4) $\Rightarrow Mmn = Mnm$.

Consequently, M is a B_1 - near-field space over a regular δ -near-ring.

(ii) Since N is zero symmetric near-field space, by result 2.9 demands that every ideal of near-field space is an N -sub near-field space of N and now (ii) follows from (i).

This completes the proof of the theorem.

3.6 Proposition: Let N be a B_1 -near-field space over a regular δ -near-ring. Then for every $a \in N$, there exists $x \in N^*$ such that the following are true.

- (i) $\exists n \in N \ni axa = nax$.
- (ii) $Nax \subset Na \cap Nx$.
- (iii) If N is a Boolean near-field space over a regular δ -near-ring then $Naxa = Nxa$.
- (iv) If N is a strong B_1 -near-field space over a regular δ -near-ring then $\exists n \in N \ni xa = nax$.

Proof: Let $a \in N$.

Since N is a B_1 -near-field space over a regular δ -near-ring, $\exists x \in N^* \ni Nax = Nxa$ (1)

- (i) Since $axa \in Nxa$, by using (1) we get $axa = nax$ for some $n \in N$ and (i) follows.
- (ii) (1) $\Rightarrow Nax = Nxa \subset Na$. Obviously, $Nax \subset Nx$.
 $\therefore Nax \subset Na \cap Nx$.
- (iii) When N is Boolean near-field space over a regular δ -near-ring, $Nxa = Nxa^2 = (Nxa)a = (Nax)a$ [by (1)] and (iii) follows.
- (iv) Since N is a strong B_1 -near-field space over a regular δ -near-ring, the result follows from result 2.12 and (i). This completes the proof of proposition.

SECTION 4: STRONG B_1 - NEAR-FIELD SPACES OVER REGULAR DELTA NEAR-RINGS

In this section, By generalizing the concept of B_1 -near-field space over a regular δ -near-ring, Dr. N.V. Nagendram introduce strong B_1 -near-field space over a regular δ -near-ring. Also study some of its important properties, obtain a simple characterization under a condition and also a structure theorem.

4.1 Definition: strong B_1 near-field space. For any sub near-field space A of near-field space N , we denote by A^* the set of all non-zero elements of A . In particular $N^* = N - \{0\}$. N is called a strong B_1 near-field space if $N^* = N_{B_1}(a) \forall a \in N$ where $N_{B_1}(a) = \{x \in N^* / axa = xa\}$.

In other words, we say that N is a strong B_1 -near-field space over a regular δ -near-ring if $Nab = Nba \forall a, b \in N$.

4.2 Example: Every commutative near-field space over a regular δ -near-ring is a strong B_1 -near-field space over a regular δ -near-ring.

4.3 Example: Consider the near-field space $(N, +, \cdot)$ where $(N, +)$ is the Klein’s four group $\{0, a, b, c\}$ and “ \cdot ” Is defined as follows (scheme p.408 Gunter Pilz[4])

.	0	a	b	c
0	0	0	0	0
a	a	a	a	a
b	0	a	b	c
c	a	0	c	b

This is a strong B_1 -near-field space over a regular δ -near-ring.

4.4 Proposition: Every strong B_1 -near-field space over a regular δ -near-ring is a B_1 -near-field space over a regular δ -near-ring.

Proof: Obvious.

4.5 Note: Converse of proposition 4.4 is not valid. For example we consider the near-field space $(N, +, \cdot)$ where $(N, +)$ is the Klein’s four group $\{0, a, b, c\}$ and “ \cdot ” Is defined as follows (Scheme (14), p.408 of Gunter Pilz [4])

.	0	a	b	c
0	0	0	0	0
a	a	a	a	a
b	0	a	b	c
c	a	0	c	b

This is a B_1 -near-field space over a regular δ -near-ring. But it is not a strong B_1 -near-field space over a regular δ -near-ring since, $Nac \neq Nca$.

4.6 Note: It is clear that the property of N being strong B_1 -near-field space over a regular δ -near-ring is preserved under near-field homo-morphisms.

Consequently, the following theorem can be deduced.

4.7 Theorem: Every strong B_1 -near-field space over a regular δ -near-ring is isomorphic to a sub direct product of sub directly irreducible strong B_1 -near-field space over a regular δ -near-rings.

Proof: by theorem 1.62, p.26 of Gunter Pilz [4] we get, N is isomorphic to a sub direct product of sub directly irreducible near-fields N_i 's, say, and each N_i is a homomorphic image of N under the usual projection map π_i . The desired result now follows from note 4.6. This completes the proof of the theorem.

4.8 Lemma: If N is a strong B_1 -near-field space over a regular δ -near-ring $\Leftrightarrow \forall a, b, c \in N, \exists n \in N \ni abc = ncb$.

Proof: \Rightarrow (If part): Let $a, b, c \in N$. Now $abc \in Nbc$.

Since N is a strong B_1 -near-field space over a regular δ -near-ring, $Nbc = Ncb$.

$\therefore abc \in Ncb$.

$\Rightarrow abc = ncb$ for some $n \in N$.

\Leftarrow (IFF Part): Let $a, b, c \in N$.

Now $abc \in Nbc$.

$\exists n \in N \ni abc = ncb \in Ncb$.

$\therefore Nbc \subset Ncb$.

In similar manner we get $Ncb \subset Nbc$. Thus N is a strong B_1 -near-field space over a regular δ -near-ring. This completes the proof of the lemma.

4.9 Theorem: Let N be a strong B_1 -near-field space over a regular δ -near-ring. If N is regular then we have the following:

- (i) $\forall a \in N, \exists x \in N$ such that $a = a^2x$.
- (ii) N has no non-zero nilpotent elements.
- (iii) Any two principal N -sub near-field spaces of N commute with each other.
- (iv) N is a P_1 -near-field space over a regular δ -near-ring.
- (v) N is left bi-potent.

Proof: Obvious.

4.10 Corollary: Let N be a zero symmetric strong B_1 -near-field space over a regular δ -near-ring. If N is regular near-field space then N is the sub-direct product of integral near-field spaces over a regular δ -near-ring.

4.11 Theorem: Let N be a strong B_1 -near-field space over a regular δ -near-ring. If N is Boolean near-field space over a regular δ -near-ring then the following are true:

- (i) $NaNb = Nab \forall a, b \in N$.
- (ii) All principal N -sub near-field spaces of N commute with one another.
- (iii) Every ideal of N is a strong near-field space over a regular δ -near-ring.
- (iv) Every N -sub near-field space of N is a strong near-field space over a regular δ -near-ring.
- (v) Every N - sub near-field space of N is an invariant N -sub near-field space of N over a regular δ -near-ring.

Proof: Obvious.

Here the discussion with the following characterization of strong near-field space over a regular δ -near-ring.

4.12 Theorem: Let N be a Boolean B_1 -near-field space over a regular δ -near-ring. Then N is a strong B_1 -near-field space over a regular δ -near-ring $\Leftrightarrow Na \cap Nb = Nab \forall a, b \in N$.

Proof: \Rightarrow (If Part) Let $y \in Na \cap Nb$.

$\therefore y = na = n'b$ for some $n, n' \in N$.

Now by lemma 4.8, $\exists z \in N \ni y^2 = (na)(n'b) = (nan')b = (zn'a)b = (zn')ab \in Nab$.

Since, N is Boolean, this yields $y \in Nab$.

Thus $Na \cap Nb \subset Nab$ (1)

Since N is a strong B_1 - near-field space over a regular δ -near-ring,
 $Nab = Nba$.

But $Nba \subset Na$ and $Nab \subset Nb$.

Hence, $Nab \subset Na \cap Nb$ (2)

(1), (2) \Rightarrow we get $Na \cap Nb = Nab$.

\Leftarrow (only IF Part): Let $a, b \in N$. now $Nab = Na \cap Nb = Nb \cap Na = Nba$. Thus N is a strong B_1 - near-field space over a regular δ -near-ring.

This completed the proof of the theorem.

SECTION 5: CONCLUSION

Near-field space theory over regular delta near-rings under algebra of mathematics has many applications in the study of permutation groups, block schemes and projective geometry. Near-field spaces over regular delta near-rings provide a non-linear analogue to the development of Linear Algebra, combinatorial problems and useful for agricultural experiments. In this paper Dr. N.V. Nagendram presented the notion of B_1 - near-field spaces over regular delta near-rings and derived the properties and characterization of the strong B_1 - near-field spaces over regular delta near-rings.

REFERENCES

1. R.Balakrishnan and S.Suryanarayanan, On P_k and P_{-k} near-rings, Bull. Malaysian Math., Sc. Soc. (Second Series) 23 (2000), 9-24.
2. J.L.Jat and S.C.Choudhary, On left bipotent near-rings, Proc. Edinburgh Math., Soc. 22(1979), no.2, 99-107.
3. Neal.H. McCoy, Theory of Rings, MacMillan & Co., 1970.
4. G.Pilz, near-rings, North Holland, Amsterdam, 1983.
5. S.Silviya, R.Balakrishnan, T.Tamizh Chelvam, Strong S_1 near-rings, International Journal of Algebra, Vol.4, 2010, no. 14, 685 - 691.
6. N V Nagendram, T V Pradeep Kumar and Y V Reddy On "Semi Noetherian Regular Matrix δ -Near Rings and their extensions", International Journal of Advances in Algebra (IJAA), Jordan, ISSN 0973-6964, Vol.4, No.1, (2011), pp.51-55.
7. N V Nagendram, T V Pradeep Kumar and Y V Reddy "A Note on Bounded Matrices over a Noetherian Regular Delta Near Rings", (BMNR-delta-NR) published in International Journal of Contemporary Mathematics, Vol.2, No.1, June 2011, Copyright@MindReaderPublications, ISSNNo:0973-6298, pp.13-19.
8. N V Nagendram, T V Pradeep Kumar and Y V Reddy "A Note on Boolean Regular Near-Rings and Boolean Regular δ -Near Rings", (BR-delta-NR) published in International Journal of Contemporary Mathematics, IJCM Int. J. of Contemporary Mathematics, Vol. 2, No. 1, June 2011, Copyright @ Mind Reader Publications, ISSN No: 0973-6298, pp. 29 - 34.
9. N V Nagendram, T V Pradeep Kumar and Y V Reddy "on p -Regular δ -Near-Rings and their extensions", (PR-delta-NR) accepted and to be published in int. J. Contemporary Mathematics (IJCM), 0973-6298, vol.1, no.2, pp.81-85, June 2011.
10. N V Nagendram, T V Pradeep Kumar and Y V Reddy "On Strongly Semi -Prime over Noetherian Regular δ -Near Rings and their extensions", (SSPNR-delta-NR) published in International Journal of Contemporary Mathematics, Vol.2, No.1, June 2011, pp.83-90.
11. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Structure Theory and Planar of Noetherian Regular δ -Near-Rings (STPLNR-delta-NR)", International Journal of Contemporary Mathematics, IJCM, published by IJSMA, pp.79-83, Dec, 2011.
12. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Matrix's Maps over Planar of Noetherian Regular δ -Near-Rings (MMPLNR-delta-NR)", International Journal of Contemporary Mathematics, IJCM, published by IJSMA, pp.203-211, Dec, 2011.
13. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On IFP Ideals on Noetherian Regular- δ - Near Rings (IFPINR-delta-NR)", Int. J. of Contemporary Mathematics, Copyright @ Mind Reader Publications, ISSN No: 0973-6298, Vol. 2, No. 1, pp.53-58, June 2011.
14. N V Nagendram, B Ramesh paper "A Note on Asymptotic value of the Maximal size of a Graph with rainbow connection number $2*(AVM-SGR-CN2^*)$ " published in an International Journal of Advances in Algebra (IJAA) Jordan @ Research India Publications, Rohini, New Delhi, ISSN 0973-6964 Volume 5, Number 2 (2012), pp. 103-112.

15. N V Nagendram and B Ramesh on "Polynomials over Euclidean Domain in Noetherian Regular Delta Near Ring Some Problems related to Near Fields of Mappings(PED-NR-Delta-NR & SPR-NF)" Accepted and published in an International Journal of Mathematical Archive (IJMA), An International Peer Review Journal for Mathematical, Science & Computing Professionals ISSN 2229-5046,vol.3,no.8,pp no. 2998-3002, 2012.
16. N V Nagendram research paper on "Near Left Almost Near-Fields (N-LA-NF)" communicated to for 2nd international conference by International Journal of Mathematical Sciences and Applications, IJMSA@mindreader publications, New Delhi on 23-04-2012 also for publication.
17. N V Nagendram, T Radha Rani, Dr T V Pradeep Kumar and Dr Y V Reddy "A Generalized Near Fields and (m, n) Bi-Ideals over Noetherian regular Delta-near rings (GNF-(m, n) BI-NR-delta-NR)", published in an International Journal of Theoretical Mathematics and Applications (TMA), Greece, Athens, dated 08-04-2012.
18. N V Nagendram, Smt.T.Radha Rani, Dr T V Pradeep Kumar and Dr Y V Reddy "Applications of Linear Programming on optimization of cool freezers(ALP-on-OCF)" Published in International Journal of Pure and Applied Mathematics, IJPAM-2012-17-670 ISSN-1314-0744 Vol-75 No-3(2011).
19. N V Nagendram "A Note on Algebra to spatial objects and Data Models(ASO-DM)" Published in international Journal American Journal of Mathematics and Mathematical Sciences, AJMMS,USA, Copyright @ Mind Reader Publications, Rohini, New Delhi, ISSN. 2250-3102, Vol.1, No.2 (Dec. 2012), pp. 233 – 247.
20. N V Nagendram, Ch Padma, Dr T V Pradeep Kumar and Dr Y V Reddy "A Note on Pi-Regularity and Pi-S-Unitality over Noetherian Regular Delta Near Rings(PI-R-PI-S-U-NR-Delta-NR)" Published in International Electronic Journal of Pure and Applied Mathematics, IeJPAM-2012-17-669 ISSN-1314-0744 Vol-75 No-4 (2011).
21. N V Nagendram, Ch Padma, Dr T V Pradeep Kumar and Dr Y V Reddy "Ideal Comparability over Noetherian Regular Delta Near Rings(IC-NR-Delta-NR)" Published in International Journal of Advances in Algebra (IJAA, Jordan), ISSN 0973-6964 Vol:5, NO:1(2012), pp.43-53@Research India publications, Rohini, New Delhi.
22. N. V. Nagendram, S. Venu Madava Sarma and T. V. Pradeep Kumar, "A NOTE ON SUFFICIENT CONDITION OF HAMILTONIAN PATH TO COMPLETE GRPHS (SC-HPCG)", IJMA-2(11), 2011, pp.1-6.
23. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Noetherian Regular Delta Near Rings and their Extensions(NR-delta-NR)",IJCMS,Bulgaria,IJCMS-5-8-2011, Vol.6, 2011, No.6, 255-262.
24. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Semi Noetherian Regular Matrix Delta Near Rings and their Extensions (SNRM-delta-R)", Jordan,@ResearchIndiaPublications,AdvancesinAlgebraISSN 0973-6964 Volume 4, Number 1 (2011), pp.51-55© Research India Publicationspp.51-55
25. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Boolean Noetherian Regular Delta Near Ring(BNR-delta-NR)s", International Journal of Contemporary Mathematics, IJCM Int. J. of Contemporary Mathematics ,Vol. 2, No. 1-2, Jan-Dec 2011 , Mind Reader Publications, ISSN No: 0973-6298,pp. 23-27.
26. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Bounded Matrix over a Noetherian Regular Delta Near Rings (BMNR-delta-NR)", Int. J. of Contemporary Mathematics,Vol. 2, No. 1-2, Jan-Dec 2011, Copyright @ Mind Reader Publications, ISSN No: 0973-6298,pp.11-16
27. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Strongly Semi Prime over Noetherian Regular Delta Near Rings and their Extensions(SSPNR-delta-NR)", Int. J. of Contemporary Mathematics,Vol. 2, No. 1, Jan-Dec 2011, Copyright @ Mind Reader Publications, ISSN No: 0973-6298, pp.69-74.
28. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On IFP Ideals on Noetherian Regular Delta Near Rings (IFPINR-delta-NR)", Int. J. of Contemporary Mathematics,Vol. 2, No. 1-2, Jan-Dec 2011, Copyright @ Mind Reader Publications, ISSN No: 0973-6298, pp.43-46.
29. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Structure Thoery and Planar of Noetherian Regular delta-Near-Rings (STPLNR-delta-NR)", International Journal of Contemporary Mathematics, IJCM, accepted for international conference conducted by IJSMA, New Delhi December 18,2011, pp:79-83, Copyright @ Mind Reader Publications and to be published in the month of Jan 2011.
30. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Matrix's Maps over Planar of Noetherian Regular delta-Near-Rings (MMPLNR-delta-NR)", International Journal of Contemporary Mathematics, IJCM, accepted for international conference conducted by IJSMA, New Delhi December 18,2011,pp:203-211,Copyright @ Mind Reader Publications and to be published in the month of Jan 2011.
31. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "Some Fundamental Results on P- Regular delta-Near-Rings and their extensions (PNR-delta-NR)", International Journal of Contemporary Mathematics, IJCM,Jan-December'2011,Copyright@MindReader Publications,ISSN:0973-6298, vol.2,No.1-2, PP.81-85.
32. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "A Generalized ideal based-zero divisor graphs of Noetherian regular Delta-near rings (GIBDNR- d-NR)" , International Journal of Theoretical Mathematics and Applications (TMA)accepted and published by TMA, Greece, Athens,ISSN:1792- 9687 (print),vol.1, no.1, 2011, 59-71, 1792-9709 (online),International Scientific Press, 2011.
33. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "Inversive Localization of Noetherian regular Delta-near rings (ILNR- Delta-NR)" , International Journal of Pure And Applied Mathematics published by IJPAM-2012-17-668, ISSN.1314-0744 vol-75 No-3,SOFIA, Bulgaria.

34. N V Nagendram, S V M Sarma, Dr T V Pradeep Kumar "A note on Relations between Barnette and Sparse Graphs" published in an International Journal of Mathematical Archive (IJMA), An International Peer Review Journal for Mathematical, Science & Computing Professionals, 2(12), 2011, pg no.2538-2542, ISSN 2229 – 5046.
35. N V Nagendram "On Semi Modules over Artinian Regular Delta Near Rings(S Modules-AR-Delta-NR) Accepted and published in an International Journal of Mathematical Archive (IJMA)", An International Peer Review Journal for Mathematical, Science & Computing Professionals ISSN 2229-5046, IJMA-3-474, 2012.
36. N V Nagendram¹, N Chandra Sekhara Rao² "Optical Near field Mapping of Plasmonic Nano Prisms over Noetherian Regular Delta Near Fields (ONFMPN-NR-Delta-NR)" accepted for 2nd international Conference by International Journal of Mathematical Sciences and Applications, IJMSA @ mind reader publications, New Delhi going to conduct on 15 – 16 th December 2012 also for publication.
37. N V Nagendram, K V S K Murthy(Yoga), "A Note on Present Trends on Yoga Apart From Medicine Usage and Its Applications (PTYAFMUIA)" published by the International Association of Journal of Yoga Therapy, IAYT 18 th August, 2012.
38. N V Nagendram, B Ramesh, Ch Padma, T Radha Rani and S V M Sarma research article "A Note on Finite Pseudo Artinian Regular Delta Near Fields (FP AR-Delta-NF)" communicated to International Journal of Advances in Algebra, IJAA, Jordan on 22 nd August 2012.
39. Dr. N.V. Nagendram advanced research article "A note on TL-ideals of near-field spaces over a regular δ -near-rings" communicated to International Journal of Mathematical Archive IJMA on 08th August 2015.
40. Dr. N.V. Nagendram advanced research article "a note on divided near-field spaces and ϕ -pseudo – valuation near-field spaces over regular δ -near-rings (dnf- ϕ pvnfs-o- δ -nr)" published in a reputed International Journal of Mathematical Archive IJMA on 2015.
41. Dr. N.V. Nagendram advanced research article accepted for IV international conference as Invited talk at Thailand "Characterization of near-fields over Baer-ideals" International Journal of Science and Mathematical Applications(IJSMA), sponsored by Kakinada Institute of technology and Science, Tirupathi(v), DIVILI 533 433 on 15-19 December, 2015.

Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2015. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]