

#g-PRECLOSED SETS IN A TOPOLOGICAL SPACES

K. ALLI*

Department of Mathematics, The M. D. T. Hindu College, Tirunelveli, Tamil Nadu, India.

(Received On: 12-08-15; Revised & Accepted On: 10-09-15)

ABSTRACT

In this paper, we introduce a new class of #gp- closed sets, and their properties. Applying #g-preclosed sets, we also introduce and study some new classes of spaces, namely, Td# spaces, #Td spaces, Td## spaces, αTd# spaces, αTd## spaces and #sTd spaces and some interrelationships between these spaces.

1991 AMS Classification: 54A05, 54D10.

Keywords & Phrases: Topological space, preclosed set and ag-closed set.

1. INTRODUCTION

N. Levine [11] introduced the class of g-closed sets. M. K. R. S. Veerakumar introduced several generalized closed sets namely, g*-closed sets, g#-closed sets, g* p- closed sets and their properties. In this paper we introduce #gp-closed sets and their properties. Applying g#-preclosed sets, we introduce and study some new classes of spaces, namely, Tp# spaces, #T_d spaces, T_d## spaces, αT_d# spaces, αT_d## spaces and #sTp spaces. We obtained some interrelationships between these spaces.

2. PRELIMINARIES

Throughout this paper (X, τ) (or X) represent topological spaces on which no separation axioms are assumed unless otherwise mentioned. For a subset A of a space (X, τ), cl(A), int(A) and A^c denote the closure of A, the interior of A and the complement of A respectively.

We recall the following definitions which are useful in the sequel.

Definition 2.1: A subset A of a space (X, τ) is called

1. a preopen set [16] if $A \subseteq \text{int}(\text{cl}(A))$ and a preclosed set if $\text{cl}(\text{int}(A)) \subseteq A$.
2. a semi-open set [12] if $A \subseteq \text{cl}(\text{int}(A))$ and a semi-closed set if $\text{int}(\text{cl}(A)) \subseteq A$.
3. an α-open set [17] if $A \subseteq \text{int}(\text{cl}(\text{int}(A)))$ and an α-closed set if $\text{cl}(\text{int}(\text{cl}(A))) \subseteq A$.
4. a semi-preopen [2] set (=β-open [1]) if $A \subseteq \text{cl}(\text{int}(\text{cl}(A)))$ and a semi-preclosed set [2] (=β-closed[1]) if $\text{int}(\text{cl}(\text{int}(A))) \subseteq A$.
5. a regular open set [20] if $A = \text{int}(\text{cl}(A))$ and a regular closed set [20] if $\text{cl}(\text{int}(A)) = A$.

The semi-closure (resp. preclosure, α-closure, semi-preclosure) of a subset A of (X, τ) is the intersection of all semi-closed (resp. preclosed, α-closed, semi-preclosed) sets that contain A and is denoted by scl(A) (resp. pcl(A), αcl(A), splc(A)).

Definition 2.2: A subset A of a space (X, τ) is called

1. a generalized closed (briefly g-closed) set [11] if $\text{cl}(A) \subseteq U$ whenever $A \subseteq U$ and U is open in (X, τ).
2. semi-generalized closed [4] (briefly sg-closed) set if $\text{scl}(A) \subseteq U$ whenever $A \subseteq U$ and U is semi-open in (X, τ). The complement of a sg-closed set is called a sg-open set.
3. a generalized semi-closed (briefly gs-closed) set [3] if $\text{scl}(A) \subseteq U$ whenever $A \subseteq U$ and U is open in (X, τ).
4. an α-generalized closed (briefly ag-closed) set [13] if $\alpha\text{cl}(A) \subseteq U$ whenever $A \subseteq U$ and U is open in (X, τ).

Corresponding Author: K. Alli*

Department of Mathematics, The M. D. T. Hindu College, Tirunelveli, Tamil Nadu, India.

5. a generalized α -closed (briefly $g\alpha$ -closed) set [14] if $\alpha cl(A) \subseteq U$ whenever $A \subseteq U$ and U is α -open in (X, τ) . The complement of a $g\alpha$ -closed set is called a $g\alpha$ -open [5] set.
6. a generalized semi-preclosed (briefly gsp -closed) set [7] if $spcl(A) \subseteq U$ whenever $A \subseteq U$ and U is open in (X, τ) .
7. a generalized preregular closed (briefly gpr -closed) set [10] if $pcl(A) \subseteq U$ whenever $A \subseteq U$ and U is regular open in (X, τ) .
8. a g^* -pre closed [18] (briefly g^*p -closed) set if $pcl(A) \subseteq U$ whenever $A \subseteq U$ and U is g -open in (X, τ) .
9. $g\#$ -closed set [19] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is αg -open in (X, τ) .
10. a generalized pre closed (briefly gp -closed) set [15] if $pcl(A) \subseteq U$ whenever $A \subseteq U$ and U is open in (X, τ) .

Definition 2.3: A space (X, τ) is called a,

1. $T1/2$ space [11] if every g -closed set is closed,
2. semi- $T1/2$ space [4] if every sg -closed set is semi-closed,
3. semi-pre- $T1/2$ space [7] if every gsp -closed set is semi-preclosed,
4. preregular $T1/2$ space [10] if every gpr -closed set is preclosed.l.

BASIC PROPERTIES OF #g-PRECLOSED SETS

We introduce the following definition.

Definition 3.1: A subset A of a space (X, τ) is called a # g -preclosed set (briefly # gp -closed set) if $pcl(A) \subseteq U$ whenever $A \subseteq U$ and U is a αg -open set of (X, τ) .

Theorem 3.2: Let (X, τ) be a topological space. Then

1. Every preclosed set is # gp -closed, and thus every $g\alpha$ -closed set, and every α -closed set and every closed set is # gp -closed.
2. Every # gp -closed set is gp -closed, and hence gsp -closed and gpr -closed too.
3. Every $g\#$ -closed set is # gp -closed.

Proof:

1. Since $pcl(A) = A$ for any preclosed set A of (X, τ) , we have that every preclosed set is # gp -closed. Since every closed (α -closed) set is α -closed ($g\alpha$ -closed) and every $g\alpha$ -closed set is preclosed (Theorem 2.4(ii) of [11]), then every $g\alpha$ -closed set, every α -closed set and every closed set is also # gp -closed.
2. Follows from the fact that every open (gp -open) set is g -open (gsp -closed and gpr -closed) (see [13]).
3. Let A be a $g\#$ -closed set and $A \subseteq U$ and U is a αg -open, $cl(A) \subseteq U$ and $pcl(A) \subseteq U$. Hence A is # gp -closed.

The reverse implications in the above theorem are not true as we see the following examples.

Example 3.3: Let $X = \{a, b, c\}$ and $\tau = \{\emptyset, X, \{a\}, \{a, c\}\}$. The set $A = \{a, b\}$ is # gp -closed but not even a preclosed set of (X, τ) .

Example 3.4: Let $X = \{a, b, c\}$ and $\tau = \{\emptyset, X, \{a\}\}$. The set $B = \{a, b\}$ is gp -closed and thus gsp -closed and gpr -closed. But B is not a # gp -closed set of (X, τ) .

Example 3.5: Let $X = \{a, b, c\}$ and $\tau = \{\emptyset, X, \{a, c\}\}$. The set $C = \{c\}$ is # gp -closed but it is not a $g\#$ -closed set of (X, τ) .

Thus the class of # gp -closed sets properly contains the classes of preclosed sets, $g\alpha$ -closed set, α -closed set, $g\#$ -closed set and the class of closed sets. Moreover the class of # gp -closed set is properly contained in the classes of gp -closed sets, gsp -closed sets and in the class of gpr -closed sets.

Theorem 3.6:

1. # gp -closed sets are independent of semi-closed sets and semi-preclosed sets.
2. # gp -closed sets are independent of g -closed sets, αg -closed sets, gs -closed sets and sg -closed sets.

Proof: Follows from the following examples.

Example 3.7: Let X, τ and A be as in the example 3.3. A is a # gp -closed set but not even a semi-preclosed set of (X, τ) .

Example 3.8: Let $X = \{a, b, c\}$ and $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$. Let $C = \{a\}$. C is semi-closed and hence a semi-preclosed set of (X, τ) . But C is not a # gp -closed set of (X, τ) . Moreover C is a sg -closed set.

Example 3.9: Let $X = \{a, b, c\}$ and $\tau = \{\emptyset, X, \{a, b\}\}$. Let $D = \{a\}$. D is #gp -closed. D is neither sg-closed nor even a gs-closed set of (X, τ) .

Example 3.10: Let X, τ and B be as in the example 3.4. B is g-closed and hence α g-closed and gs-closed set in (X, τ) . But B is not a #gp -closed set of (X, τ) .

Remark 3.11: Union of two #gp -closed sets need not be #gp -closed set again.

Proof: Let (X, τ) be as in the example 3.9. $A = \{a\}$ and $B = \{b\}$ are #gp -closed sets but $A \cup B = \{a, b\}$, their union is not a #gp -closed set of (X, τ) .

Theorem 3.12: If A is α g-open and #gp -closed set of (X, τ) , then A is a preclosed set of (X, τ) .


Theorem 3.13: Let A be a #gp -closed set of (X, τ) . Then

1. $\text{pcl}(A) - A$ does not contain any non-empty α g-closed set.
2. If $A \subseteq B \subseteq \text{pcl}(A)$, then B is also a #gp -closed set of (X, τ) .

Proof:

1. Let F be a g-closed set contained in $\text{pcl}(A) - A$. $\text{pcl}(A) \subseteq X - F$ since $X - F$ is g-open set with $A \subseteq X - F$ and A is a #gp -closed. Then $F \subseteq (X - \text{pc}(A)) \cap (\text{pc}(A) - A) \subseteq (X - \text{pcl}(A)) \cap \text{pcl}(A) = \emptyset$. Therefore $F = \emptyset$.
2. Let U be a g-open set of (X, τ) such that $B \subseteq U$. Then $A \subseteq U$. Since $A \subseteq U$ and A is #gp -closed, $\text{pcl}(A) \subseteq U$. Then $\text{pcl}(B) \subseteq \text{pcl}(\text{pcl}(A)) = \text{pcl}(A)$ since $B \subseteq \text{pcl}(A)$. Thus $\text{pcl}(B) \subseteq \text{pcl}(A) \subseteq U$. Hence B is also a #gp -closed set of (X, τ) .

Theorem 3.14: The following diagram shows the relationships between #gp -closed sets and some other sets.



where $A \xrightarrow{\quad} B$ ($A \xleftarrow{\quad} B$) represents A implies B and B need not imply A (A and B are independent of each other).

4. APPLICATIONS OF #g-PRECLOSED SETS

We now introduce and study some new spaces, namely $T_d^\#$ spaces, ${}^\#T_d$ spaces, $T_d^{\#\#}$ spaces, ${}_\alpha T_d^\#$ spaces, ${}_\alpha T_d^{\#\#}$ spaces and ${}^\#_s T_d$ spaces.

Definition 4.1: A topological space (X, τ) is said to be

1. a $T_d^\#$ space if every #gp -closed set in it is closed.
2. a ${}^\#T_d$ space if every gp-closed set in it is #gp -closed.
3. a $T_d^{\#\#}$ space if every #gp -closed set in it is $g\alpha$ -closed.
4. a ${}_\alpha T_d^\#$ space if every #gp -closed set in it is preclosed.
5. a ${}_\alpha T_d^{\#\#}$ space if every #gp -closed set in it is α -closed.
6. a ${}^\#_s T_d$ space if every gsp-closed set in it is #gp -closed.

Theorem 4.2:

1. Every ${}_{\alpha}T_d^{##}$ space is ${}_{\alpha}T_d^{\#}$ space and $T_d^{##}$ space.
2. Every $T_d^{\#}$ space is ${}_{\alpha}T_d^{##}$ space and hence $T_d^{##}$ space ${}_{\alpha}T_d^{\#}$ space.
3. Every $T_d^{##}$ space is ${}_{\alpha}T_d^{\#}$ space.

Proof:

1. Follows from the fact that every α -closed set is $g\alpha$ -closed and hence a preclosed set.
2. Follows from (1) and the fact that every closed set is an α -closed set.
3. Follows from the fact that every $g\alpha$ -closed set is preclosed.

Following examples show that the reverse implications in the above theorem need not be true in general.

Example 4.3: Let $X = \{a, b, c\}$ and $\tau = \{\phi, X, \{a\}, \{b, c\}\}$. $\{b\}$ is a [#]gp-closed set but not even an α -closed set. So (X, τ) is neither $T_d^{\#}$ nor ${}_{\alpha}T_d^{##}$. However (X, τ) is a $T_d^{##}$ space.

Example 4.4: Let $X = \{a, b, c\}$ and $\tau = \{\phi, X, \{a\}\}$. (X, τ) is not a $T_d^{\#}$ space since $\{b\}$ is a [#]gp -closed set but not a closed set. However (X, τ) is an ${}_{\alpha}T_d^{\#}$ space.

Example 4.5: Let $X = \{a, b, c\}$ and $\tau = \{\phi, X, \{a, b\}\}$. (X, τ) is not even a $T_d^{##}$ space since $\{a\}$ is a [#]gp -closed set but not a $g\alpha$ -closed set. However (X, τ) is an ${}_{\alpha}T_d^{\#}$ space.

Thus $T_d^{\#}$ spaces is properly contained in the classes of ${}_{\alpha}T_d^{##}$ spaces, $T_d^{##}$ spaces and in the class of ${}_{\alpha}T_d^{\#}$ spaces. The class of ${}_{\alpha}T_d^{##}$ spaces is properly contained in the classes of ${}_{\alpha}T_d^{\#}$ spaces and $T_d^{##}$ spaces. Moreover the class of ${}_{\alpha}T_d^{\#}$ spaces properly contains the class of $T_d^{##}$ spaces.

Theorem 4.6:

1. Every sT_d space is a [#] T_d space.
2. Every semi-pre- $T_{1/2}$ space is an ${}_{\alpha}T_d^{\#}$ space and a [#] T_d space.

Proof:

1. Follows from the fact that every gp-closed set is gsp-closed.
2. Follows from the fact that every [#]gp -closed (resp. preclosed) set is gp-closed (resp. [#]gp -closed) set and the Lemma 2.4 of [3].

Following examples show that the reverse implications in the above theorem need not be true in general.

Example 4.7: Let $X = \{a, b, c\}$ and $\tau = \{\phi, X, \{a\}, \{b\}, \{a, b\}\}$. $\{a, b\}$ is a gsp-closed set but not a [#]gp -closed set of (X, τ) . Thus (X, τ) is not a sT_d space. However (X, τ) is a [#] T_d space.

Example 4.8: Let (X, τ) be as in the example 4.4. (X, τ) is not a semi-pre- $T_{1/2}$ space since $\{a, b\}$ is a gsp-closed set but not a semi-preclosed set. However (X, τ) is an ${}_{\alpha}T_d^{\#}$ space.

Example 4.9: Let (X, τ) be as in the example 3.3. (X, τ) is not a semi-pre- $T_{1/2}$ space since $\{a, b\}$ is a gsp-closed set but not a semi-preclosed set. However (X, τ) is a [#] T_d space.

Thus the class of [#] T_d spaces properly contains the class of sT_d spaces. Moreover the class of semi-pre- $T_{1/2}$ spaces is properly contained in the class of ${}_{\alpha}T_d^{\#}$ spaces and in the class of [#] T_d spaces.

Dontchev [10] proved that the dual of the class of semi- $T_{1/2}$ spaces to the class of $T_{1/2}$ spaces is the class of semi-pre- $T_{1/2}$ spaces.

Theorem 4.10: A space (X, τ) is semi-pre- $T_{1/2}$ if and only if it is ${}_{\alpha}T_d^{\#}$ and [#] T_d .

Proof: Follows from the theorem 2.4 of [4].

Thus the class of [#] T_d spaces is the dual of the class of semi-pre- $T_{1/2}$ spaces to the class of ${}_{\alpha}T_d^{\#}$ spaces.

The following theorem gives a tridecomposition for $T_{1/2}$ spaces.

Theorem 4.11: A space (X, τ) is $T_{1/2}$ if and only if it is semi- $T_{1/2}$, ${}_{\alpha}T_d^{\#}$ and [#] T_d .

Proof: Follows from the above theorem 4.10 and the theorem 4.5 of Dontchev [10].

Theorem 4.12:

1. $\alpha T_d^{\#}$ spaces are independent of ${}^{\#}T_d$ spaces and ${}_s T_d^{\#}$ spaces.
2. ${}^{\#}T_d$ spaces are independent of $T_d^{\#\#}$ spaces and $\alpha T_d^{\#\#}$ spaces.
3. ${}_s T_d^{\#}$ spaces are independent of $T_d^{\#}$ spaces, $T_d^{\#\#}$ spaces, $\alpha T_d^{\#}$ spaces, $\alpha T_d^{\#\#}$ spaces and semi-pre- $T_{1/2}$ spaces.
4. $\alpha T_d^{\#\#}$ spaces are independent of semi-pre- $T_{1/2}$ spaces and preregular $T_{1/2}$ spaces.
5. $T_d^{\#}$ spaces are independent of preregular $T_{1/2}$ spaces.

Proof: The space (X, τ) in the example 3.8 is a $T_d^{\#}$ space but it is not ${}_s T_d^{\#}$. Moreover (X, τ) is not preregular $T_{1/2}$. The space (X, τ) in the example 4.3 is ${}_s T_d^{\#}$, preregular $T_{1/2}$, semi-pre- $T_{1/2}$, ${}^{\#}T_d$ but not even an $\alpha T_d^{\#\#}$ space. The space (X, τ) in the example 4.4 is an $\alpha T_d^{\#\#}$ space and thus $\alpha T_d^{\#}$ and also $T_d^{\#\#}$ but it is not even a ${}^{\#}T_d$ space. The space (X, τ) in the example 4.5 is a ${}_s T_d^{\#}$ space and thus a ${}^{\#}T_d$ space but it is not even a $T_d^{\#\#}$ space. The space in the example 4.9 is ${}^{\#}T_d$ but not $\alpha T_d^{\#}$.

Theorem 4.13:

1. Every preregular $T_{1/2}$ space is a ${}^{\#}T_d$ space but not conversely.
2. Every preregular $T_{1/2}$ space is an $\alpha T_d^{\#}$ space but not conversely.

Proof:

1. The first assertion follows from the fact that every preclosed set is a ${}^{\#}gp$ -closed set. The space (X, τ) in the example 3.3 is a ${}^{\#}T_d$ space. But (X, τ) is not a preregular $T_{1/2}$ space.
2. The first assertion follows from the theorem 4.06(2) and the corollary 5.8 of [13]. The space (X, τ) in the example 3.8 supports the second assertion.

Thus the class of preregular $T_{1/2}$ spaces is properly contained in the class of $\alpha T_d^{\#}$ spaces and in the class of ${}^{\#}T_d$ spaces.

Definition 4.14: A subset A of a space (X, τ) is called a ${}^{\#}gp$ -open if $C(A)$ is ${}^{\#}gp$ -closed.


Theorem 4.15: The following statements are true but the respective converses are not true in general.

1. If (X, τ) is a $T_d^{\#}$ space, then every singleton of X is either αg -closed or open.
2. If (X, τ) is a ${}^{\#}T_d$ space, then every singleton of X is either closed or ${}^{\#}gp$ -open.
3. If (X, τ) is a $T_d^{\#\#}$ space, then every singleton of X is either αg -closed or $g\alpha$ -open.
4. If (X, τ) is an $\alpha T_d^{\#}$ space, then every singleton of X is either αg -closed or pre-open.
5. If (X, τ) is an $\alpha T_d^{\#\#}$ space, then every singleton of X is either αg -closed or α -open.
6. If (X, τ) is a ${}_s T_d^{\#}$ space, then every singleton of X is either closed or ${}^{\#}gp$ -open.

Proof: (1): Let $x \in X$ and suppose that $\{x\}$ is not a g -closed set of (X, τ) . This implies $X - \{x\}$ is not a g -open set. So X is the only g -open set such that $X - \{x\} \subseteq X$. Then $X - \{x\}$ is a ${}^{\#}gp$ -closed set of (X, τ) . Since (X, τ) is a $T_d^{\#}$ space, then $X - \{x\}$ is closed or equivalently $\{x\}$ is open. The space in the example 4.03 shows that the converse need not be true.

The proofs for the first assertions of (2) to (6) are similar to as that of the first assertion of (1). The space (X, τ) in the example 4.4 shows that converse of (2) need not be true. The space (X, τ) in the example 4.3 shows that the converse of (4) need not be true. The space (X, τ) in the example 4.0 shows that the converses of (3), (5) and (6) need not be true.

Theorem 4.16: The following diagram shows some relationships among the spaces considered in this paper.



where $A \longrightarrow B$ (resp. $A \longleftarrow \text{---} B$) represents A implies B but B need not imply A (resp. A and B are independent of each other).

REFERENCES

1. Abd El-Monsef, M.E., El-Deeb, S.N. and Mahmoud, R.A.: β -open sets and β -continuous mappings, Bull. Fac. Sci. Assiut Univ., 12(1983), 77-90.
2. Andrijević, D.: Semi-preopen sets, Mat. Vesnik, 38(1) (1986), 24-32.
3. Arya, S.P. and Nour, T.: Characterizations of s -normal spaces, Indian J. Pure. Appl. Math., 21(8) (1990), 717-719.
4. Bhattacharya, P. and Lahiri, B.K.: Semi-generalized closed sets in topology, Indian J. Math., 29(3)(1987), 375-382.
5. Devi, R., Balachandran, K. and Maki, H.: Generalized α -closed maps and α -generalized closed maps, Indian J. Pure. Appl. Math., 29(1) (1998), 37-49.
6. Dontchev, J.: On door spaces, Indian J. Pure. Appl. Math., 26(9) (1995), 873-881.
7. Dontchev, J.: On generalizing semi-preopen sets, Mem. Fac. Sci. Kochi Univ. Ser. A. Math., 16(1995), 35-48.
8. Dontchev, J.: On some separation axioms associated with the α -topology, Mem. Fac. Sci. Kochi Univ. Ser. A. Math, 18(1997), 31-35.
9. Dunham, W.: $T_{1/2}$ spaces, Kyungpook Math. J., 17(1977), 161-169.
10. Gnanambal, Y.: On generalized preregular closed sets in topological spaces, Indian J. Pure. Appl. Math., 28(3) (1997), 351-360.
11. Levine, N.: Generalized closed sets in topology, Rend. Circ. Math. Palermo, 19(2) (1970), 89-96.
12. Levine. Semi-open sets and semi-continuity in topological spaces, Amer. Math. Monthly, 70(1963), 36-41.
13. Maki, H., Devi, R. and Balachandran, k.: Associated topologies of generalized α -closed sets and α -generalized closed sets, Mem. Fac. Sci. Kochi Univ. Ser. A. Math., 15(1994), 51-63.
14. Maki, H., Devi, R. and Balachandran, k.: Generalized α -closed sets in topology, Bull. Fukuoka Univ. Ed. Part III, 42(1993), 13-21.
15. Maki, H., Umehara, J. and Noiri, T.: Every topological space is pre- $T_{1/2}$ space, Mem. Fac. Sci. Kochi Univ. Ser. A. Math., 17(1996), 33-42.
16. Mashhour, A.S., Abd El-Monsef, M.E. and El-Deeb, S.N.: On pre-continuous and weak pre-continuous mappings, Proc. Math. and Phys. Soc. Egypt, 53(1982), 47-53.
17. Njåstad, O.: On some classes of nearly open sets, Pacific J. Math., 15(1965), 961-970.
18. Veera kumar, M.K.R.S.: g^* -preclosed sets Acta ciencia Indica (Maths) Meerut XXVIII (M) (1) 2002, 51-60.
19. Veera kumar, M.K.R.S.: $g\#$ -closed sets in topological spaces, Mem. Fac. Sci. Kochi Univ. Ser. A. Math 24 (2003), 1-13.
20. Kuratowski, C.: Topology I, Hafner, New York, 1958.

Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2015. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]