

ACCURATE INDEPENDENT DOMINATION IN FUZZY GRAPHS

*P. GLADYIS, **C. V. R. HARINARAYANAN

*Assistant Professor of Mathematics,
Bharathidasan University Model College, Aranthangi, Pudukkottai, (T.N.), India.

**Research Supervisor and Assistant Professor of Mathematics,
Govt. Arts College, Paramakudi, (T.N.), India.

(Received On: 22-09-15; Revised & Accepted On: 26-10-15)

ABSTRACT

A dominating set D of a fuzzy graph $G = (V, E)$ is an independent dominating set if the induced subgraph $\langle D \rangle$ has no edges. An independent dominating set D of a fuzzy graph G is an accurate independent dominating set if $V - D$ has no independent dominating set of cardinality $|D|$. The fuzzy accurate independent domination number $i_{fa}(G)$ of G is the minimum cardinality of an accurate dominating set of G . In this paper we study a accurate independent domination in fuzzy graphs and investigate the relationship of $i_{fa}(G)$ with other known parameters.

Index terms: Fuzzy Graph, Fuzzy Independent Dominating set, Fuzzy Accurate Independent Dominating set, Fuzzy Accurate Independent Domination Number.

INTRODUCTION

A fuzzy subset of a non empty set V is a mapping $\sigma: V \rightarrow [0,1]$. A fuzzy relation on V is a fuzzy subset of $V \times V$. A fuzzy graph $G = (\sigma, \mu)$ is a pair of function $\sigma: V \rightarrow [0,1]$ and $\mu: V \times V \rightarrow [0, 1]$, where $\mu(u, v) \leq \sigma(u) \wedge \sigma(v)$ for all $u, v \in V$. The order p and size q of the fuzzy graph $G = (\sigma, \mu)$ are define by $p = \sum_{v \in V} \sigma(v)$ and $q = \sum_{u,v \in E} \mu(u, v)$. The complement of a fuzzy graph $G = (\sigma, \mu)$ is a fuzzy graph $\bar{G} = (\bar{\sigma}, \bar{\mu})$ where $\bar{\sigma} = \sigma$ and $\bar{\mu}(u, v) = \sigma(u) \wedge \sigma(v) - \mu(u, v)$ for all u, v in V . The fuzzy cardinality of a fuzzy subset D of V is $|D|_f = \sum_{v \in D} \sigma(v)$. An edge $e = \{u, v\}$ of a fuzzy graph is called an effective edge if $\mu(u, v) = \sigma(u) \wedge \sigma(v)$. The effective degree of a vertex u is defined to be the sum of the weights of the effective edges incident at u and is denoted by $d_E(u)$. The Minimum effective degree $\delta_E(G) = \min\{d_E(u) / u \in V(G)\}$ and the maximum effective degree $\Delta_E(G) = \max\{d_E(u) / u \in V(G)\}$. A set of fuzzy vertex which cover all the fuzzy edges is called a fuzzy vertex cover of G and the minimum cardinality of a fuzzy vertex cover is called a vertex covering number of G and is denoted by $\alpha_0(G)$. A set of fuzzy edge which cover all the fuzzy vertices is called a fuzzy edge cover of G and the minimum cardinality of a fuzzy edge cover is called a edge covering number of G and is denoted by $\alpha_1(G)$. The vertex independence number $\beta_0(G)$ of G is the maximum cardinality among the independent sets of vertices. The edge independence number $\beta_1(G)$ of G is the maximum cardinality among the independent sets of edges. For any graph H is a complete subgraph of G is called a Clique of G . The number of vertices in a largest Clique of G is called the Clique number $\omega(G)$ of G . If $\mu(u, v) = 0$ for every $v \in V$ then u is called isolated node. A set $S \subseteq V$ in a fuzzy graph G is said to be independent if $\mu(u, v) < \sigma(u) \wedge \sigma(v)$ for all $u, v \in S$. A dominating set is called an independent dominating set if D is independent. An independent dominating set S of a fuzzy graph G is said to be a maximal independent dominating set if there is no independent dominating set S^1 of G such that $S^1 \subset S$. An independent dominating set S of a fuzzy graph G is said to be a maximum independent dominating set if there is no independent dominating set S^1 of G such that $|S^1| > |S|$. The minimum scalar cardinality of an maximum independent dominating set of G is called the independent domination number of G and is denoted by $i(G)$. Let $x, y \in V$. We say that x dominates y in G if $\mu(u, v) = \sigma(u) \wedge \sigma(v)$. A subset S of V is called a dominating set in G if for every $v \notin S$, there exists $u \in S$ such that u dominates v . The minimum cardinality of a dominating set in G is called the domination number of G and is denoted by $\gamma(G)$. Let $G = (\sigma, \mu)$ be a fuzzy graph. A subset D of V is said to be fuzzy dominating set of G if for every $v \in V - D$ there exists $u \in D$ such that (u, v) is a strong arc. A dominating set D of a graph G is called minimal dominating set of G if for every node $v \in D$, $D - \{v\}$ is not a dominating set of the domination number $\gamma(G)$ is the minimum cardinalities taken over all minimal dominating sets of G . A dominating set D of a graph G is an accurate dominating set, if $V - D$ has no dominating set of cardinality $|D|$. The accurate domination number $\gamma_a(G)$ of G is the

Corresponding Author: *P. Gladys

minimum cardinality of an accurate dominating set. A dominating set D of a fuzzy graph G is an accurate dominating set, if $V-D$ has no dominating set of cardinality $|D|$. The fuzzy accurate domination number $\gamma_{fa}(G)$ of G is the minimum cardinality of an accurate dominating set. An independent dominating set D of a graph G is an accurate independent dominating set if $V-D$ has no independent dominating set of cardinality $|D|$. The accurate independent domination number $i_a(G)$ of G is the minimum cardinality of an accurate dominating set of G . An independent dominating set D of a fuzzy graph G is an accurate independent dominating set if $V-D$ has no independent dominating set of cardinality $|D|$. The fuzzy accurate independent domination number $i_{fa}(G)$ of G is the minimum cardinality of an accurate independent dominating set of G .

1. ACCURATE DOMINATION IN FUZZY GRAPHS

Definition: 1.1 A dominating set D of G is an accurate dominating set if $V-D$ has no dominating set of cardinality $|D|$. The accurate domination number $\gamma_{fa}(G)$ of G is the minimum cardinality of an accurate dominating set of G .


Theorem: 1.1 For any fuzzy graph $p - q \leq \gamma_{fa} \leq p - \delta_E$ where p, q and δ_E are the order, size and minimum effective incident degree of G respectively.

Proof: Let D be a accurate dominating set and γ_{fa} be the minimum fuzzy domination number in G . Then the scalar cardinality of $V-D$ is less than or equal to the scalar cardinality of $V \times V$. Hence $p-q \leq \gamma_{fa}$. Now, let u be the node with minimum effective incident degree δ_E , clearly $V-\{u\}$ is a accurate dominating set and hence $\gamma_{fa} \leq p - \delta_E$. Hence $p - q \leq \gamma_{fa} \leq p - \delta_E$ is true for any fuzzy graph.

Theorem: 1.2 $\lceil \frac{p}{1+\Delta(G)} \rceil \leq \gamma_{fa} \leq p - \Delta(G)$.

Theorem: 1.3 If G is a fuzzy graph without isolated nodes then $\gamma_{fa}(G) \leq \min\{\alpha_0(G), \alpha_1(G), \beta_0(G), \beta_1(G)\}$.

Example: 1.1



Here $D = \{3, 4\}, \gamma_{fa}(G) = 1.3$
 $\alpha_0(G) = 1.3, \alpha_1(G) = 1.3$
 $\beta_0(G) = 1.7, \beta_1(G) = 1.7$

Theorem: 1.4 For any fuzzy graph G and \bar{G} are both connected then $\gamma_{fa}(G) + \gamma_{fa}(\bar{G}) \leq p + 1$.

Proof: We know that $\gamma_{fa}(G) \leq p - \Delta(G)$ and $\gamma_{fa}(\bar{G}) \leq p - \Delta(\bar{G})$.

$$\begin{aligned} \text{Therefore } \gamma_{fa}(G) + \gamma_{fa}(\bar{G}) &\leq p - \Delta(G) + p - \Delta(\bar{G}) \\ &= 2p - (\Delta(G) + \Delta(\bar{G})) \\ &= 2p - (\Delta(G) + p - 1 - \delta(G)) \\ &= p + 1 + \delta(G) - \Delta(G) \text{ Since } \delta(G) - \Delta(G) \leq 0 \\ &\leq p + 1. \end{aligned}$$

Theorem: 1.5 For any fuzzy graph G and \bar{G} are both connected then $\gamma_{fa}(G) + \gamma_{fa}(\bar{G}) \leq p(p - 3)$.


Theorem: 1.6 For any fuzzy graph G and \bar{G} are both connected then

- (i) $\gamma_{fa}(G) + \gamma_{fa}(\bar{G}) \leq 2(p - 2)$
- (ii) $\gamma_{fa}(G) + \gamma_{fa}(\bar{G}) \leq (p - 2)^2$

2. ACCURATE INDEPENDENT DOMINATION IN FUZZY GRAPHS

Definition: An independent dominating set D of G is an accurate independent dominating set if $V-D$ has no independent dominating set of cardinality $|D|$. The accurate independent domination number $i_{fa}(G)$ of G is the minimum cardinality of an accurate dominating set of G

Example: 2.1



Here $D = \{v_2, v_5\}$ and $i_{fa}(G) = 1.4$
 $P = 3.7, q = 3.4, \Delta(G) = 1.5$
 $\delta(G) = 0.4, \alpha_0(G) = 1.4, \beta_0(G) = 1.9$

Theorem: 2.1 For any fuzzy graph G , $i_{fa}(G) \leq \beta_0(G)$.

Proof: Let S be an Independent set of nodes in G such that $|S| = \beta_0(G)$. Then G contains no larger independent set. Then $V - S$ has no independent dominating set of cardinality $|S|$. Therefore S is an accurate independent dominating set. Thus $i_{fa}(G) \leq |S|, \therefore i_{fa}(G) \leq \beta_0(G)$.

Theorem: 2.2 For any fuzzy graph G , $i_{fa}(G) \leq p - \gamma_f(G) + 1$.

Theorem: 2.3 For any fuzzy graph G , $\frac{p}{\Delta+1} \leq i_{fa}(G) \leq \frac{p\Delta}{\Delta+1} + 1$.

Proof: We know that $\frac{p}{\Delta+1} \leq \gamma_f(G) \rightarrow$ (a) and since $\gamma_f(G) \leq i_{fa}(G) \rightarrow$ (b)

From equation (a) and (b) we get $\frac{p}{\Delta+1} \leq i_{fa}(G)$. So lower bound is attained.

Using the previous theorem, $i_{fa}(G) \leq p - \gamma_f(G) + 1$
 $\leq p - \frac{p}{\Delta+1} + 1$
 $\leq \frac{p\Delta}{\Delta+1} + 1 \rightarrow$ (c)

From equation (a), (b) & (c) we get
 $\frac{p}{\Delta+1} \leq i_{fa}(G) \leq \frac{p\Delta}{\Delta+1} + 1$.

Theorem: 2.4 For any fuzzy graph G , $\lfloor \frac{p}{1+\Delta(G)} \rfloor \leq i_{fa}(G)$.

Theorem: 2.5 For any fuzzy graph G with $p \geq 2$ nodes, an independent dominating set with $\lfloor \frac{p}{2} \rfloor + 1$ nodes is an accurate independent dominating set.

Proof: Let D be an independent dominating set with $\lfloor \frac{p}{2} \rfloor + 1$ nodes. Then $|V-D| < \frac{p}{2}$. Hence D is an accurate Independent dominating set of G .

Theorem: 2.6 For any connected non trivial fuzzy graph G , $i_{fa}(G) + i_{fa}[L(G)] \leq p$. Where $L(G)$ is a line graph.

Proof: Let G be a connected graph. For any Fuzzy graph $i_{fa}[L(G)] \leq \beta_1(G)$

Also $i_{fa}(G) \leq \alpha_1(G)$

Hence $i_{fa}(G) + i_{fa}[L(G)] \leq \alpha_1(G) + \beta_1(G)$
 $= V(G)$
 $= p$

Therefore $i_{fa}(G) + i_{fa}[L(G)] \leq p$.


Theorem: 2.7 If G is a fuzzy graph without isolated nodes, then $i_{fa}(G) \leq \alpha_0(G) + 1$.

Theorem: 2.8 Let G be a fuzzy graph such that both G and \bar{G} have no isolated nodes then

$$i_{fa}(G) + i_{fa}(\bar{G}) \leq 2 \lceil \frac{p}{2} \rceil$$

$$i_{fa}(G) \cdot i_{fa}(\bar{G}) \leq \lceil \frac{p}{2} \rceil^2$$

Example: 2.2



Here $D = \{2,5\}$ $D = \{4,5\}$

$i_{fa}(G) = 1.1, p=2.8$ $i_{fa}(\bar{G}) = 1.3$

Theorem: 2.9 For any fuzzy graph G and \bar{G} have no isolated nodes then

$$i_{fa}(G) + i_{fa}(\bar{G}) \leq p + \alpha_0(G) - \omega(G) + 2.$$

Proof: From Theorem 2.7, $i_{fa}(G) \leq \alpha_0(G) + 1$

$$\begin{aligned} \text{Also } i_{fa}(\bar{G}) &\leq \alpha_0(\bar{G}) + 1 \\ &\leq p - \beta_0(\bar{G}) + 1 \\ &\leq p - \omega(G) + 1 \end{aligned}$$

Thus $i_{fa}(G) + i_{fa}(\bar{G}) \leq p + \alpha_0(G) - \omega(G) + 2.$

3. RELATION BETWEEN ACCURATE DOMINATION AND ACCURATE INDEPENDENT DOMINATION IN FUZZY GRAPHS

Theorem: 3.1 For any fuzzy graph G , $\gamma_{fa}(G) \leq i_{fa}(G) \rightarrow (1)$

Proof: Every independent accurate dominating set is an accurate dominating set. Thus (1) holds.

Theorem: 3.2 For any fuzzy graph G , $\gamma_f(G) \leq \gamma_{fa}(G) \leq i_{fa}(G)$

Theorem: 3.3 For any fuzzy graph G , $i_{fa}(G) \leq p - \gamma_{fa}(G) + 1.$

Proof: Let D be a minimum independent accurate dominating set G . Then for any node $v \in D$, $(V - D) \cup \{v\}$ is an accurate independent dominating set of G .

$$\begin{aligned} \text{Thus } i_{fa}(G) &\leq |(V - D) \cup \{v\}| \\ &= p - \gamma_{fa}(G) + 1. \end{aligned}$$

Theorem: 3.4 For any non-trivial connected fuzzy graph G , $\gamma_{fa}(G) + i_{fa}(G) \leq p + q.$

Proof: Since $i_{fa}(G) \leq \beta_0(G)$ [From Theorem 2.1]


Also $\gamma_{fa}(G) \leq \alpha_0(G)$

$$\begin{aligned} \text{Further } \gamma_{fa}(G) + i_{fa}(G) &\leq \alpha_0(G) + \beta_0(G) \\ &= \gamma(G) \cup E(G) \\ &= p + q \end{aligned}$$

Hence $\gamma_{fa}(G) + i_{fa}(G) \leq p + q$.

Theorem: 3.5 For any fuzzy graph G , $i_{fa}(G) \leq \gamma_{fa}(G) + \delta(G)$.

Example: 3.1



$D = \{V_4, V_5\}$
 $D = \{V_2, V_5\}$

$\gamma_{fa}(G) = 1.2$
 $i_{fa}(G) = 1.4$

REFERENCES

1. S.Ismail Mohideen and A.Mohamed Ismayil, Domination in Fuzzy Graph: A New Approach, International Journal of Computational Science and Mathematics (2010).
2. V.R.Kulli and M.B.Kattimani, Accurate domination in graphs, In V.R.Kulli, Ed., Advances in Domination Theory I, Vishwa International Publications, Gulbarga, India (2012) 1-8.
3. V.R.Kulli and M.B.Kattimani, The Accurate domination number of a graphs, Technical Report 2000:01, Dept. of. Math. Gulbarga University, Gulbarga. India (2000).
4. V.R.Kulli, Theory of Domination in Graphs, Vishwa International Publications, Gulbarga, India (2010).

Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2015. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]