

FIXED POINT, COINCIDENCE POINT AND COMMON FIXED POINT THEOREMS
UNDER VARIOUS EXPANSIVE CONDITIONS IN TRIANGULAR
INTUITIONISTIC FUZZY METRIC SPACE

RASHMI JAIN¹, R. D. DAHERIYA¹, MANOJ UGHAD^{*2}

¹Department of Mathematics,
Government J.H. Post Graduate College, Betul, P.O. Box 460001, (M.P.), India.

²Department of Mathematics,
Sarvepalli Radhakrishnan University, Bhopal, P.O. Box 462026, (M.P.), India.

(Received On: 20-09-15; Revised & Accepted On: 27-10-15)

ABSTRACT

In this article, we establish some fixed point, common fixed point and coincidence point theorems for expansive type mappings in the triangular intuitionistic fuzzy metric spaces. The presented theorems extend, generalize and improve many existing results in the literature.

2010 Mathematics Subject Classifications: 47H10, 54H25.

Keywords: intuitionistic fuzzy metric space; fixed point, common fixed point, coincidence point; expansive mappings.

1. INTRODUCTION AND PRELIMINARIES

The concept of fuzzy set was introduced by Zadeh [1] in 1965. In 1975, Kramosil and Michalek [2] introduced the notion of fuzzy metric space, which can be regarded as a generalization of the statistical (probabilistic) metric space. This work has provided an important basis for the construction of fixed point theory in fuzzy metric spaces.

In 2004, Park introduced the notion of intuitionistic fuzzy metric space [12]. He showed that, for each intuitionistic fuzzy metric space $(X, M, N, *, \diamond)$, the topology generated by the intuitionistic fuzzy metric (M, N) coincides with the topology generated by the fuzzy metric M . For more details on intuitionistic fuzzy metric space and related results we refer the reader to [12–20]. The study of expansive mappings is a very interesting research area in fixed point theory. In 1984, Wang et.al [31] introduced the concept of expanding mappings and proved some fixed point theorems in complete metric spaces. In 1992, Daffer and Kaneko [30] defined an expanding condition for a pair of mappings and proved some common fixed point theorems for two mappings in complete metric spaces. Chintaman and Jagannath [32] introduced several meaningful fixed point theorems for one expanding mapping.

In this paper, we present some new fixed point, coincidence point and common fixed point theorems under various expansive conditions in triangular intuitionistic fuzzy metric space. These results improve and generalize some important known results in [23-33]. Some related results to highlight the realized improvements is also furnished.

Throughout this paper \mathbb{R} and \mathbb{R}_+ will represents the set of real numbers and nonnegative real numbers, respectively. The following two definitions are required in the sequel which can be found in [12].

Definition 1.1: A binary operation $*$: $[0, 1] \times [0, 1] \rightarrow [0, 1]$ is continuous t-norm if $*$ satisfying the following conditions:

- (1). $*$ is commutative and associative;
- (2). $*$ is continuous;
- (3). $a * 1 = a, \forall a \in [0, 1]$;
- (4). $a * b \leq c * d$ whenever $a \leq c$ and $b \leq d, \forall a \in [0, 1]$.

**Corresponding Author: Manoj Ughade*², ²Department of Mathematics,
Sarvepalli Radhakrishnan University, Bhopal, P.O. Box 462026, (M.P.), India.**

Definition 1.2: A binary operation $\diamond : [0, 1] \times [0, 1] \rightarrow [0, 1]$ is continuous t-conorm if \diamond satisfying the following conditions:

- (1). \diamond is commutative and associative;
- (2). \diamond is continuous;
- (3). $a \diamond 0 = a, \forall a \in [0, 1]$;
- (4). $a \diamond b \leq c \diamond d$ whenever $a \leq c$ and $b \leq d, \forall a \in [0, 1]$.

In 2004, Park [12] introduced the concept of intuitionistic fuzzy metric space as follows.

Definition 1.3: A 5-tuple $(X, M, N, *, \diamond)$ is said to be an intuitionistic fuzzy metric space if X is an arbitrary set, $*$ is a continuous t-norm, \diamond is a continuous t-conorm, and M, N are two fuzzy sets on $X^2 \times (0, \infty)$ satisfying the following conditions, for all $x, y, z \in X$ and $s, t > 0$:

- (IFMS1). $M(x, y, t) + N(x, y, t) \leq 1$;
- (IFMS2). $M(x, y, t) > 0$;
- (IFMS3). $M(x, y, t) = 1$ for all $t > 0 \Leftrightarrow x = y$;
- (IFMS4). $M(x, y, t) = M(y, x, t)$;
- (IFMS5). $M(x, y, t) * M(y, z, s) \leq M(x, z, t + s)$;
- (IFMS6). $M(x, y, \cdot) : (0, \infty) \rightarrow [0, 1]$ is left continuous;
- (IFMS7). $\lim_{t \rightarrow \infty} M(x, y, t) = 1$;
- (IFMS8). $N(x, y, t) > 0$;
- (IFMS9). $N(x, y, t) = 0$ for all $t > 0 \Leftrightarrow x = y$;
- (IFMS10). $N(x, y, t) = N(y, x, t)$;
- (IFMS11). $N(x, y, t) \diamond N(y, z, s) \geq N(x, z, t + s)$;
- (IFMS12). $N(x, y, \cdot) : (0, \infty) \rightarrow [0, 1]$ is right continuous;
- (IFMS13). $\lim_{t \rightarrow \infty} N(x, y, t) = 0$;

Then (M, N) is called an intuitionistic fuzzy metric space on X . The functions $M(x, y, t)$ and $N(x, y, t)$ denote the degree of nearness and the degree on nonnearness between x and y with respect to t , respectively.

Definition 1.4: (see [12]) Let $(X, M, N, *, \diamond)$ be an intuitionistic fuzzy metric space. Then

- 1) a sequence $\{x_n\}$ is said to be Cauchy sequence whenever

$$\lim_{m, n \rightarrow \infty} M(x_n, x_m, t) = 1$$
 and

$$\lim_{m, n \rightarrow \infty} N(x_n, x_m, t) = 0$$
 for all $t > 0$. That is, for each $\varepsilon > 0$ and $t > 0$, there exists a natural number n_0 such that $M(x_n, x_m, t) > 1 - \varepsilon$ and $N(x_n, x_m, t) < \varepsilon$ for all $n, m \geq n_0$.
- 2) $(X, M, N, *, \diamond)$ is called complete whenever every Cauchy sequence is convergent with respect to the topology $\tau_{(M, N)}$.

Remark 1.5: Note that, if (M, N) is called an intuitionistic fuzzy metric space on X and $\{x_n\}$ is a sequence in X such that $\lim_{m, n \rightarrow \infty} M(x_n, x_m, t) = 1$ and $\lim_{m, n \rightarrow \infty} N(x_n, x_m, t) = 0$ for all $t > 0$ as from (IFMS1) of Definition 1.3, we know that $M(x, y, t) + N(x, y, t) \leq 1$ for all $x, y \in X$ and $t > 0$.

Let $(X, M, N, *, \diamond)$ be an intuitionistic fuzzy metric space. According to [8, 10], the fuzzy metric (M, N) is called triangular whenever $\frac{1}{M(x, y, t)} - 1 \leq \frac{1}{M(x, z, t)} - 1 + \frac{1}{M(z, y, t)} - 1$ and $N(x, y, t) \leq N(x, z, t) + N(z, y, t)$ for all $x, y, z \in X$ and $t > 0$.

Example 1.6: Let $X = \{(0,0), (0,4), (4,0), (4,5), (5,4)\}$ endowed with the metric $d: X \times X \rightarrow [0, +\infty)$ given by

$$d((x_1, x_2), (y_1, y_2)) = |x_1 - y_1| + |x_2 - y_2|$$

for all $(x_1, x_2), (y_1, y_2) \in X$. Define intuitionistic fuzzy metric by

$$M((x_1, x_2), (y_1, y_2), t) = \frac{t}{t + d((x_1, x_2), (y_1, y_2))} \quad \text{and} \quad N((x_1, x_2), (y_1, y_2), t) = \frac{d((x_1, x_2), (y_1, y_2))}{t + d((x_1, x_2), (y_1, y_2))}$$

for all $(x_1, x_2), (y_1, y_2) \in X$ and $t > 0$, where

$$a * b = \min\{a, b\} \quad \text{and} \quad a \diamond b = \max\{a, b\}.$$

Then X is a complete triangular intuitionistic fuzzy metric space.

Example 1.4: Let $X = [0, 2 - \sqrt{3})$ endowed with the usual distance $d(x, y) = |x - y|$. Consider

$$M(x, y, t) = \frac{t}{t+d(x,y)} \text{ and } N(x, y, t) = \frac{d(x,y)}{t+d(x,y)}$$

for all $x, y \in X$ and $t > 0$. Then X is a complete triangular intuitionistic fuzzy metric space.

2. MAIN RESULT

Now, we are ready to state and prove our main results.

Theorem 2.1: Let (X, M, N, \star, ϕ) be a complete triangular intuitionistic fuzzy metric space. Let $T: X \rightarrow X$ be a continuous self-mapping satisfying the condition

$$\frac{1}{M(Tx, Ty, t)} - 1 \geq a \left(\frac{1}{M(x, y, t)} - 1 \right) + b \left(\frac{1}{M(x, Tx, t)} - 1 \right) + c \left(\frac{1}{M(y, Ty, t)} - 1 \right) \tag{2.1}$$

for all $x, y \in X$, all $t > 0$ and $a > 1, b \in \mathbb{R}$ and $c \leq 1$ with $a + b + c > 1$. Then T has a unique fixed point.

Proof: Let x_0 be arbitrary in X , we define a sequence $\{x_n\}$ in X by the rule

$$x_0 = Tx_1, x_1 = Tx_2, \dots, x_n = Tx_{n+1} \tag{2.2}$$

If $x_n = x_{n+1}$ for some n then we have nothing to prove. Assume that $x_n \neq x_{n+1}$ for all n . Consider

$$\frac{1}{M(x_n, x_{n-1}, t)} - 1 = \frac{1}{M(Tx_{n-1}, Tx_n, t)} - 1 \tag{2.3}$$

Now by (2.1) and definition of sequence

$$\begin{aligned} \frac{1}{M(x_n, x_{n-1}, t)} - 1 &= \frac{1}{M(Tx_{n-1}, Tx_n, t)} - 1 \leq a \left(\frac{1}{M(x_{n-1}, x_n, t)} - 1 \right) + b \left(\frac{1}{M(x_{n+1}, Tx_{n+1}, t)} - 1 \right) + c \left(\frac{1}{M(x_n, Tx_n, t)} - 1 \right) \\ \frac{1}{M(x_n, x_{n-1}, t)} - 1 &\geq a \left(\frac{1}{M(x_{n+1}, x_n, t)} - 1 \right) + b \left(\frac{1}{M(x_{n+1}, x_n, t)} - 1 \right) + c \left(\frac{1}{M(x_n, x_{n-1}, t)} - 1 \right) \end{aligned} \tag{2.4}$$

By use of symmetric property, we have

$$\frac{1}{M(x_{n-1}, x_n, t)} - 1 \geq a \left(\frac{1}{M(x_n, x_{n+1}, t)} - 1 \right) + b \left(\frac{1}{M(x_n, x_{n+1}, t)} - 1 \right) + c \left(\frac{1}{M(x_{n-1}, x_n, t)} - 1 \right)$$

Thus $(1 - c) \left(\frac{1}{M(x_{n-1}, x_n, t)} - 1 \right) \geq (a + b) \left(\frac{1}{M(x_n, x_{n+1}, t)} - 1 \right)$

$$\frac{1}{M(x_n, x_{n+1}, t)} - 1 \leq \frac{1-c}{a+b} \left(\frac{1}{M(x_{n-1}, x_n, t)} - 1 \right) \tag{2.5}$$

Let $k = \frac{1-c}{a+b} < 1$. So the above inequality, become

$$\frac{1}{M(x_n, x_{n+1}, t)} - 1 \leq k \left(\frac{1}{M(x_{n-1}, x_n, t)} - 1 \right) \tag{2.6}$$

Also, we can show that

$$\frac{1}{M(x_{n-1}, x_n, t)} - 1 \leq k \left(\frac{1}{M(x_{n-2}, x_{n-1}, t)} - 1 \right) \tag{2.7}$$

So that

$$\frac{1}{M(x_n, x_{n+1}, t)} - 1 \leq k^2 \left(\frac{1}{M(x_{n-2}, x_{n-1}, t)} - 1 \right) \tag{2.8}$$

Proceeding in similar way we can get,

$$\frac{1}{M(x_n, x_{n+1}, t)} - 1 \leq k^n \left(\frac{1}{M(x_0, x_1, t)} - 1 \right) \tag{2.9}$$

for all n .

By using the triangular inequality, for each $m \geq n$, we obtain

$$\begin{aligned} \frac{1}{M(x_n, x_m, t)} - 1 &\leq \frac{1}{M(x_n, x_{n+1}, t)} - 1 + \frac{1}{M(x_{n+1}, x_{n+2}, t)} - 1 + \dots + \frac{1}{M(x_{m-1}, x_m, t)} - 1 \\ &\leq (k^n + k^{n+1} + \dots + k^{m-1}) \left(\frac{1}{M(x_0, x_1, t)} - 1 \right) \\ &\leq \frac{k^n}{1-k} \left(\frac{1}{M(x_0, x_1, t)} - 1 \right) \end{aligned} \tag{2.10}$$

Assume that $M(x_0, x_1, t) < 1$ that is, $\frac{1}{M(x_0, x_1, t)} - 1 > 0$. Letting $n \rightarrow \infty$, $\{x_n\}$ is a Cauchy sequence. Also, if $M(x_0, x_1, t) = 1$, then $M(x_n, x_m, t) = 1$ for all $m > n$ and hence $\{x_n\}$ is a Cauchy sequence. Since X is complete. So there must exist $x^* \in X$ such that

$$\lim_{n \rightarrow \infty} x_n = x^* \tag{2.11}$$

Now to show that x^* is a fixed point of T . Since T is continuous so

$$\lim_{n \rightarrow \infty} Tx_n = Tx^* \Rightarrow \lim_{n \rightarrow \infty} x_{n-1} = Tx^* \Rightarrow Tx^* = x^* \tag{2.12}$$

Hence x^* is the fixed point of T . Now suppose that T has another fixed point $y^* \neq x^*$. then we have

$$\begin{aligned} \frac{1}{M(x^*, y^*, t)} - 1 &= \frac{1}{M(Tx^*, Ty^*, t)} - 1 \\ &\geq a \left(\frac{1}{M(x^*, y^*, t)} - 1 \right) + b \left(\frac{1}{M(x^*, Tx^*, t)} - 1 \right) + c \left(\frac{1}{M(y^*, Ty^*, t)} - 1 \right) \\ &= a \left(\frac{1}{M(x^*, y^*, t)} - 1 \right) \end{aligned} \tag{2.13}$$

Since $a > 1$, so the above inequality is possible if

$$\frac{1}{M(x^*, y^*, t)} - 1 = 0 \tag{2.14}$$

which implies that $x^* = y^*$. Hence fixed point of T is unique.

Corollary 2.2: Let $(X, M, N, *, \delta)$ be a complete triangular intuitionistic fuzzy metric space. Let $T: X \rightarrow X$ be a continuous self-mapping satisfying the condition

$$\frac{1}{M(Tx, Ty, t)} - 1 \geq a \left(\frac{1}{M(x, y, t)} - 1 \right) \tag{2.15}$$

for all $x, y \in X$ and $a > 1$. Then T has a unique fixed point.

Proof: Putting $b = c = 0$ in Theorem 2.1 one can get the required result without any difficulty.

Corollary 2.3: Let $(X, M, N, *, \delta)$ be a complete triangular intuitionistic fuzzy metric space. Let $T: X \rightarrow X$ be a continuous self-mapping satisfying the condition

$$\frac{1}{M(Tx, Ty, t)} - 1 \geq a \left(\frac{1}{M(x, y, t)} - 1 \right) + b \left(\frac{1}{M(x, Tx, t)} - 1 \right) \tag{2.16}$$

for all $x, y \in X$ and $a > 1, b \in \mathbb{R}$ with $a + b > 1$. Then T has a unique fixed point.

Proof: Putting $c = 0$ in Theorem 2.1 one can get the required result without any difficulty.

Theorem 2.4: Let $(X, M, N, *, \delta)$ be a complete triangular intuitionistic fuzzy metric space. Let $T: X \rightarrow X$ be a surjective self-mapping satisfying the condition (2.1). Then T has a unique fixed point.

Proof: Let x_0 be arbitrary in X , we define a sequence $\{x_n\}$ in X by the rule $x_0 = Tx_1, x_1 = Tx_2, \dots, x_n = Tx_{n+1}$. Proceeding like Theorem 2.1, we obtain that $\{x_n\}$ is a Cauchy sequence in complete triangular intuitionistic fuzzy metric space. So there must exist $x^* \in X$ such that $\lim_{n \rightarrow \infty} x_n = x^*$. Now to show that x^* is a fixed point of T . Since T is surjective continuous mapping, so for any $p \in X, Tp = x^*$. Consider

$$\begin{aligned} \frac{1}{M(x_n, x^*, t)} - 1 &= \frac{1}{M(Tx_{n+1}, Tp, t)} - 1 \\ &\geq a \left(\frac{1}{M(x_{n+1}, p, t)} - 1 \right) + b \left(\frac{1}{M(x_{n+1}, Tx_{n+1}, t)} - 1 \right) + c \left(\frac{1}{M(p, Tp, t)} - 1 \right) \\ \frac{1}{M(x_n, x^*, t)} - 1 &\geq a \left(\frac{1}{M(x_{n+1}, p, t)} - 1 \right) + b \left(\frac{1}{M(x_{n+1}, x_n, t)} - 1 \right) + c \left(\frac{1}{M(p, u, t)} - 1 \right) \end{aligned} \tag{2.17}$$

Taking limit $n \rightarrow \infty$, we get

$$0 \geq (a + c) \left(\frac{1}{M(p, x^*, t)} - 1 \right) \Rightarrow M(p, x^*, t) = 1 \Rightarrow p = x^* \tag{2.18}$$

So $Tp = x^*$ becomes $Tx^* = x^*$. Thus x^* is a fixed point of T .

Uniqueness Follows from Theorem 2.1.

Theorem 2.5: Let $T, S: X \rightarrow X$ be two surjective self mappings of a complete triangular intuitionistic fuzzy metric space. Suppose that T and S satisfying the following inequalities

$$\left(\frac{1}{M(T(Sx), Sx, t)} - 1\right) + k \left(\frac{1}{M(T(Sx), x, t)} - 1\right) \geq a \left(\frac{1}{M(Sx, x, t)} - 1\right) \quad (2.19)$$

$$\text{and } \left(\frac{1}{M(S(Tx), Tx, t)} - 1\right) + k \left(\frac{1}{M(S(Tx), x, t)} - 1\right) \geq b \left(\frac{1}{M(Tx, x, t)} - 1\right) \quad (2.20)$$

for all $x, y \in X$ and some nonnegative real numbers a, b and k with $a > 1 + 2k$ and $b > 1 + 2k$. If T or S is continuous, then T and S have a common fixed point.

Proof: Let x_0 be arbitrary in X . since T is surjective, there exists $x_1 \in X$ such that $x_0 = Tx_1$. Also, since S is surjective, there exists $x_2 \in X$ such that $x_2 = Sx_1$. Continuing this process, we construct a sequence $\{x_n\}$ in X such that

$$x_{2n} = Tx_{2n+1} \text{ and } x_{2n+1} = Sx_{2n+2} \quad (2.21)$$

for all $n \in \mathbb{N} \cup \{0\}$. Now for $n \in \mathbb{N} \cup \{0\}$, we have

$$\left(\frac{1}{M(T(Sx_{2n+2}), Sx_{2n+2}, t)} - 1\right) + k \left(\frac{1}{M(T(Sx_{2n+2}), x_{2n+2}, t)} - 1\right) \geq a \left(\frac{1}{M(Sx_{2n+2}, x_{2n+2}, t)} - 1\right)$$

Thus, we have

$$\left(\frac{1}{M(x_{2n}, x_{2n+1}, t)} - 1\right) + k \left(\frac{1}{M(x_{2n}, x_{2n+2}, t)} - 1\right) \geq a \left(\frac{1}{M(x_{2n+1}, x_{2n+2}, t)} - 1\right)$$

which implies that

$$\left(\frac{1}{M(x_{2n}, x_{2n+1}, t)} - 1\right) + k \left(\frac{1}{M(x_{2n}, x_{2n+1}, t)} - 1\right) + k \left(\frac{1}{M(x_{2n+1}, x_{2n+2}, t)} - 1\right) \geq a \left(\frac{1}{M(x_{2n+1}, x_{2n+2}, t)} - 1\right)$$

$$\text{Hence } \left(\frac{1}{M(x_{2n+1}, x_{2n+2}, t)} - 1\right) \leq \frac{1+k}{a-k} \left(\frac{1}{M(x_{2n}, x_{2n+1}, t)} - 1\right) \quad (2.22)$$

On other hand, we have

$$\left(\frac{1}{M(S(Tx_{2n+1}), Tx_{2n+1}, t)} - 1\right) + k \left(\frac{1}{M(S(Tx_{2n+1}), x_{2n+1}, t)} - 1\right) \geq b \left(\frac{1}{M(Tx_{2n+1}, x_{2n+1}, t)} - 1\right)$$

Thus we have,

$$\left(\frac{1}{M(x_{2n-1}, x_{2n}, t)} - 1\right) + k \left(\frac{1}{M(x_{2n-1}, x_{2n+1}, t)} - 1\right) \geq b \left(\frac{1}{M(x_{2n}, x_{2n+1}, t)} - 1\right) \quad (2.23)$$

Since

$$\left(\frac{1}{M(x_{2n-1}, x_{2n}, t)} - 1\right) + \left(\frac{1}{M(x_{2n}, x_{2n+1}, t)} - 1\right) \geq \left(\frac{1}{M(x_{2n-1}, x_{2n+1}, t)} - 1\right)$$

We have

$$\left(\frac{1}{M(x_{2n-1}, x_{2n}, t)} - 1\right) + k \left(\frac{1}{M(x_{2n-1}, x_{2n}, t)} - 1\right) + k \left(\frac{1}{M(x_{2n}, x_{2n+1}, t)} - 1\right) \geq b \left(\frac{1}{M(x_{2n}, x_{2n+1}, t)} - 1\right) \quad (2.24)$$

$$\text{Hence, } \left(\frac{1}{M(x_{2n}, x_{2n+1}, t)} - 1\right) \leq \frac{1+k}{b-k} \left(\frac{1}{M(x_{2n-1}, x_{2n}, t)} - 1\right)$$

$$\text{Let } k = \max \left\{ \frac{1+k}{a-k}, \frac{1+k}{b-k} \right\}$$

Then by combining (2.22) and (2.24), we have

$$\left(\frac{1}{M(x_n, x_{n+1}, t)} - 1\right) \leq \lambda \left(\frac{1}{M(x_{n-1}, x_n, t)} - 1\right) \quad (2.25)$$

Repeating (2.25) n -times, we get

$$\left(\frac{1}{M(x_n, x_{n+1}, t)} - 1\right) \leq \lambda^n \left(\frac{1}{M(x_0, x_1, t)} - 1\right) \quad (2.26)$$

Thus, for $m > n$, we have

$$\frac{1}{M(x_n, x_m, t)} - 1 \leq \frac{1}{M(x_n, x_{n+1}, t)} - 1 + \frac{1}{M(x_{n+1}, x_{n+2}, t)} - 1 + \dots + \frac{1}{M(x_{m-1}, x_m, t)} - 1 \quad (2.27)$$

$$\leq (\lambda^n + \lambda^{n+1} + \dots + \lambda^{m-1}) \left(\frac{1}{M(x_0, x_1, t)} - 1\right) \leq \frac{\lambda^n}{1-\lambda} \left(\frac{1}{M(x_0, x_1, t)} - 1\right) \quad (2.28)$$

Assume that $M(x_0, x_1, t) < 1$ that is, $\frac{1}{M(x_0, x_1, t)} - 1 > 0$. Letting $n \rightarrow \infty$, $\{x_n\}$ is a Cauchy sequence. Also, if $M(x_0, x_1, t) = 1$, then $M(x_n, x_m, t) = 1$ for all $m > n$ and hence $\{x_n\}$ is a Cauchy sequence in X . So there must exist $x^* \in X$ such that $x_n \rightarrow x^*$ as $n \rightarrow +\infty$. therefore $x_{2n+1} \rightarrow x^*$ and $x_{2n+2} \rightarrow x^*$ as $n \rightarrow +\infty$. Without loss of generality, we may assume that T is continuous, then $Tx_{2n+1} \rightarrow Tx^*$ as $n \rightarrow +\infty$. But $Tx_{n+1} = x_{2n} \rightarrow x^*$ as $n \rightarrow +\infty$. Thus we have $Tx^* = x^*$. Since S is surjective, there exists $y^* \in X$ such that $Sy^* = x^*$.

Now,

$$\left(\frac{1}{M(T(Sy^*), Sy^*, t)} - 1\right) + k \left(\frac{1}{M(T(Sy^*), y^*, t)} - 1\right) \geq a \left(\frac{1}{M(Sy^*, y^*, t)} - 1\right)$$

implies that

$$k \left(\frac{1}{M(x^*, y^*, t)} - 1\right) \geq a \left(\frac{1}{M(x^*, y^*, t)} - 1\right)$$

Thus,
$$\left(\frac{1}{M(x^*, y^*, t)} - 1\right) \leq \frac{k}{a} \left(\frac{1}{M(x^*, y^*, t)} - 1\right) \tag{2.29}$$

Since $a > k$, we conclude that $\frac{1}{M(x^*, y^*, t)} - 1 = 0$ and so $x^* = y^*$. Hence $Tx^* = Sx^* = x^*$. Therefore x^* is a common fixed point of T and S .

By taking $b = a$ in Theorem 2.5, we have the following result.

Corollary 2.6: Let $T, S: X \rightarrow X$ be two surjective self mappings of a complete triangular intuitionistic fuzzy metric space. Suppose that T and S satisfying the following inequalities

$$\left(\frac{1}{M(T(Sx), Sx, t)} - 1\right) + k \left(\frac{1}{M(T(Sx), x, t)} - 1\right) \geq a \left(\frac{1}{M(Sx, x, t)} - 1\right) \tag{2.30}$$

and
$$\left(\frac{1}{M(S(Tx), Tx, t)} - 1\right) + k \left(\frac{1}{M(S(Tx), x, t)} - 1\right) \geq a \left(\frac{1}{M(Tx, x, t)} - 1\right) \tag{2.31}$$

for all $x, y \in X$ and some nonnegative real numbers a, b and k with $a > 1 + 2k$. If T or S is continuous, then T and S have a common fixed point.

By taking $S = T$ in Corollary 2.6, we have the following corollary.

Corollary 2.7: Let $S: X \rightarrow X$ be a continuous surjective self mappings of a complete triangular intuitionistic fuzzy metric space. Suppose that S satisfying the following inequalities

$$\left(\frac{1}{M(S(Sx), Sx, t)} - 1\right) + k \left(\frac{1}{M(S(Sx), x, t)} - 1\right) \geq a \left(\frac{1}{M(Sx, x, t)} - 1\right) \tag{2.32}$$

for all $x, y \in X$ and some nonnegative real numbers a, b and k with $a > 1 + 2k$. Then S has a common fixed point.

Theorem 2.8: Let $(X, M, N, *, \diamond)$ be a complete triangular intuitionistic fuzzy metric space. Let $T, f: X \rightarrow X$ be a mapping satisfying

$$\frac{1}{M(Tx, Ty, t)} - 1 \geq a \left(\frac{1}{M(fx, fy, t)} - 1\right) + b \left(\frac{1}{M(fx, Tx, t)} - 1\right) + c \left(\frac{1}{M(fy, Ty, t)} - 1\right) \tag{2.33}$$

for all $x, y \in X$, where $a, b, c \geq 0$, with $a + b + c > 1$. Suppose the following hypotheses:

- (1) $b < 1$ or $c < 1$.
- (2) $fX \subseteq TX$.
- (3) TX is a complete subspace of X .

Then T and f have a coincidence point.

Proof: Let $x_0 \in X$. Since $fX \subseteq TX$, we choose $x_1 \in X$ such that $Tx_1 = fx_0$. Again, we can choose that $x_2 \in X$ such that $Tx_2 = fx_1$. continuing in this same way, we construct a sequence $\{x_n\}$ in X such that for all $n \in \mathbb{N} \cup \{0\}$. If $fx_{m-1} = fx_m$ for some $m \in \mathbb{N}$, then $Tx_m = fx_m$. Thus x_m is a coincidence point of T and f . Now, assume that $x_{n-1} \neq x_n$ for all $m \in \mathbb{N}$.

Case-1: Suppose $b < 1$, by (2.33), we have

$$\begin{aligned} \frac{1}{M(fx_{n-1}, fx_n, t)} - 1 &= \frac{1}{M(Tx_n, Tx_{n+1}, t)} - 1 \\ &\geq a \left(\frac{1}{M(fx_n, fx_{n-1}, t)} - 1\right) + b \left(\frac{1}{M(fx_n, Tx_n, t)} - 1\right) + c \left(\frac{1}{M(fx_{n+1}, Tx_{n+1}, t)} - 1\right) \\ &= a \left(\frac{1}{M(fx_n, fx_{n-1}, t)} - 1\right) + b \left(\frac{1}{M(fx_n, fx_{n-1}, t)} - 1\right) + c \left(\frac{1}{M(fx_{n+1}, fx_n, t)} - 1\right) \end{aligned} \tag{2.34}$$

Thus, we have

$$(1 - b) \left(\frac{1}{M(fx_{n-1}, fx_n, t)} - 1 \right) \geq (a + c) \left(\frac{1}{M(fx_{n+1}, fx_n, t)} - 1 \right) \quad (2.35)$$

Hence,

$$\left(\frac{1}{M(fx_{n+1}, fx_n, t)} - 1 \right) \leq \left(\frac{1-b}{a+c} \right) \left(\frac{1}{M(fx_{n-1}, fx_n, t)} - 1 \right) \quad (2.36)$$

Case-2: Suppose $c < 1$, by (2.33), we have

$$\begin{aligned} \frac{1}{M(fx_n, fx_{n-1}, t)} - 1 &= \frac{1}{M(Tx_{n+1}, Tx_n, t)} - 1 \\ &\geq a \left(\frac{1}{M(fx_{n+1}, fx_n, t)} - 1 \right) + b \left(\frac{1}{M(fx_{n+1}, Tx_{n+1}, t)} - 1 \right) + c \left(\frac{1}{M(fx_n, Tx_n, t)} - 1 \right) \\ &= a \left(\frac{1}{M(fx_{n+1}, fx_{n+1}, t)} - 1 \right) + b \left(\frac{1}{M(fx_{n+1}, fx_n, t)} - 1 \right) + c \left(\frac{1}{M(fx_n, fx_{n-1}, t)} - 1 \right) \end{aligned} \quad (2.37)$$

Thus we have,

$$(1 - c) \left(\frac{1}{M(fx_{n-1}, fx_n, t)} - 1 \right) \geq (a + b) \left(\frac{1}{M(fx_{n+1}, fx_n, t)} - 1 \right) \quad (2.38)$$

Hence,

$$\left(\frac{1}{M(fx_{n+1}, fx_n, t)} - 1 \right) \leq \left(\frac{1 - c}{a + b} \right) \left(\frac{1}{M(fx_{n-1}, fx_n, t)} - 1 \right)$$

In case (1), we get $\lambda = \frac{1-b}{a+c}$ and in case (2), we get $\lambda = \frac{1-c}{a+b}$. Thus in both cases, we have $\lambda < 1$ and

$$\frac{1}{M(fx_{n+1}, fx_n, t)} - 1 \leq \lambda \left(\frac{1}{M(fx_{n-1}, fx_n, t)} - 1 \right) \quad (2.39)$$

Repeating (2.39) n-times we get,

$$\frac{1}{M(fx_{n+1}, fx_n, t)} - 1 \leq \lambda^n \left(\frac{1}{M(fx_0, fx_1, t)} - 1 \right) \quad (2.40)$$

So for $m > n$, we have

$$\begin{aligned} \frac{1}{M(fx_n, fx_m, t)} - 1 &\leq \frac{1}{M(fx_n, fx_{n+1}, t)} - 1 + \frac{1}{M(fx_{n+1}, fx_{n+2}, t)} - 1 + \dots + \frac{1}{M(fx_{m-1}, fx_m, t)} - 1 \\ &\leq (\lambda^n + \lambda^{n+1} + \dots + \lambda^{m-1}) \left(\frac{1}{M(fx_0, fx_1, t)} - 1 \right) \\ &\leq \frac{\lambda^n}{1-\lambda} \left(\frac{1}{M(fx_0, fx_1, t)} - 1 \right) \end{aligned} \quad (2.41)$$

Assume that $M(fx_0, fx_1, t) < 1$ that is, $\frac{1}{M(fx_0, fx_1, t)} - 1 > 0$. Letting $n \rightarrow \infty$, $\{Tx_n\}$ is a Cauchy sequence. Also, if $M(fx_0, fx_1, t) = 1$, then $M(Tx_n, Tx_m, t) = 1$ for all $m > n$ and hence $\{Tx_n\}$ is a Cauchy sequence in X. Therefore $\{Tx_n\}$ is a Cauchy sequence in TX. Since TX is complete. There is $x^* \in X$ such that $\{Tx_n\}$ converges to Tx^* as $n \rightarrow +\infty$. Hence fx_n converges to Tx^* as $n \rightarrow +\infty$. Since $a + b + c > 1$, we have a, b and c are not all 0. So we have the following cases.

Case-1: If $a \neq 0$, then

$$\begin{aligned} \frac{1}{M(Tx_n, Tx^*, t)} - 1 &\geq a \left(\frac{1}{M(fx_n, fx^*, t)} - 1 \right) + b \left(\frac{1}{M(fx_n, Tx_n, t)} - 1 \right) + c \left(\frac{1}{M(fu, Tu, t)} - 1 \right) \\ &\geq a \left(\frac{1}{M(fx_n, fu, t)} - 1 \right) \end{aligned}$$

Hence, $\left(\frac{1}{M(fx_n, fx^*, t)} - 1 \right) \leq \frac{1}{a} \left(\frac{1}{M(Tx_n, Tx^*, t)} - 1 \right) \quad (2.42)$

Since $\frac{1}{a} \left(\frac{1}{M(Tx_n, Tx^*, t)} - 1 \right) \rightarrow 0$ as $n \rightarrow +\infty$. Thus $fx_n \rightarrow fx^*$ as $n \rightarrow +\infty$. By uniqueness of limit, we have $Tx^* = fx^*$. Therefore T and f have a coincidence point.

Case-2: If $b \neq 0$, then

$$\begin{aligned} \frac{1}{M(Tx^*, Tx_n, t)} - 1 &\geq a \left(\frac{1}{M(fx_n, fx^*, t)} - 1 \right) + b \left(\frac{1}{M(fx^*, Tx^*, t)} - 1 \right) + c \left(\frac{1}{M(fx_n, Tx_n, t)} - 1 \right) \\ &\geq b \left(\frac{1}{M(fx^*, Tx^*, t)} - 1 \right) \end{aligned}$$

Hence, $\frac{1}{b} \frac{1}{M(Tx^*, Tx_n, t)} - 1 \geq \frac{1}{M(fx^*, Tx^*, t)} - 1 \quad (2.43)$

As similar proof of case (1), we can show that $fx^* = Tx^*$. Thus f and T have a coincidence point.

Case-3: If $c \neq 0$, then

$$\begin{aligned} \frac{1}{M(Tx_n, Tu, t)} - 1 &\geq a \left(\frac{1}{M(fx_n, fx^*, t)} - 1 \right) + b \left(\frac{1}{M(fx_n, Tx_n, t)} - 1 \right) + c \left(\frac{1}{M(Tu, fu, t)} - 1 \right) \\ &\geq c \left(\frac{1}{M(fx^*, Tx^*, t)} - 1 \right) \end{aligned}$$

Hence,
$$\frac{1}{M(fx^*, Tx^*, t)} - 1 \leq \frac{1}{c} \left(\frac{1}{M(Tx_n, Tx^*, t)} - 1 \right) \tag{2.44}$$

As similar proof of case (1), we can show that $fx^* = Tx^*$. Thus f and T have a coincidence point.

By taking $c = 0$ in Theorem 2.8, we have the following.

Corollary 2.9: Let (X, M, N, \star, ϕ) be a complete triangular intuitionistic fuzzy metric space. Let $T, f: X \rightarrow X$ be a mapping satisfying;

$$\frac{1}{M(Tx, Ty, t)} - 1 \geq a \left(\frac{1}{M(fx, fy, t)} - 1 \right) + b \left(\frac{1}{M(fx, Tx, t)} - 1 \right) \tag{2.45}$$

for all $x, y \in X$, where $a, b \geq 0$, with $a + b > 1$. Suppose the following hypotheses:

- (1) $b < 1$.
- (2) $fX \subseteq TX$.
- (3) TX is a complete subspace of X .

Then T and f have a coincidence point.

By taking $b = c = 0$ in Theorem 2.8, we have the following.

Corollary 2.10: Let (X, M, N, \star, ϕ) be a complete triangular intuitionistic fuzzy metric space. Let $T, f: X \rightarrow X$ be a mapping satisfying

$$\frac{1}{M(Tx, Ty, t)} - 1 \geq a \left(\frac{1}{M(fx, fy, t)} - 1 \right) \tag{2.46}$$

for all $x, y \in X$, where $a > 1$. Suppose the following hypotheses:

- (1) $fX \subseteq TX$.
- (2) TX is a complete subspace of X .

Then T and f have a coincidence point.

By taking $T = I$ (Identity on X) in Theorem 2.8, we have the following.

Corollary 2.11: Let (X, M, N, \star, ϕ) be a complete triangular intuitionistic fuzzy metric space. Let $T: X \rightarrow X$ be a mapping satisfying

$$\frac{1}{M(Tx, Ty, t)} - 1 \geq a \left(\frac{1}{M(x, y, t)} - 1 \right) + b \left(\frac{1}{M(x, Tx, t)} - 1 \right) + c \left(\frac{1}{M(y, Ty, t)} - 1 \right) \tag{2.47}$$

for all $x, y \in X$, where $a, b, c \geq 0$, with $a + b + c > 1$. Suppose that $b < 1$ or $c < 1$. Then T has a fixed point.

Corollary 2.12: Let (X, M, N, \star, ϕ) be a complete triangular intuitionistic fuzzy metric space. Let $T: X \rightarrow X$ be a mapping satisfying

$$\frac{1}{M(Tx, Ty, t)} - 1 \geq a \left(\frac{1}{M(x, y, t)} - 1 \right) + b \left(\frac{1}{M(x, Tx, t)} - 1 \right) \tag{2.48}$$

for all $x, y \in X$, where $a, b \geq 0$, with $a + b > 1$. Suppose that $b < 1$. Then T has a fixed point.

Corollary 2.13: Let (X, M, N, \star, ϕ) be a complete triangular intuitionistic fuzzy metric space. Let $T: X \rightarrow X$ be a mapping satisfying

$$\frac{1}{M(Tx, Ty, t)} - 1 \geq a \left(\frac{1}{M(x, y, t)} - 1 \right) + b \left(\frac{1}{M(y, Ty, t)} - 1 \right) \tag{2.49}$$

for all $x, y \in X$, where $a, b \geq 0$, with $a + b > 1$. Suppose that $b < 1$. Then T has a fixed point.

Corollary 2.14: Let (X, M, N, \star, ϕ) be a complete triangular intuitionistic fuzzy metric space. Let $T: X \rightarrow X$ be a mapping satisfying

$$\frac{1}{M(Tx, Ty, t)} - 1 \geq a \left(\frac{1}{M(x, y, t)} - 1 \right) \tag{2.50}$$

for all $x, y \in X$, where $a > 1$. Then T has a fixed point.

COMPETING INTERESTS

No conflict of interest was declared by the authors.

AUTHOR'S CONTRIBUTIONS

All authors contributed equally and significantly to writing this paper. All authors read and approved the final manuscript.

REFERENCES

1. L. A. Zadeh, "Fuzzy sets," *Information and Computation*, vol. 8, pp. 338–353, 1965.
2. I. Kramosil and J. Michalek, "Fuzzy metrics and statistical metric spaces," *Kybernetika*, vol. 11, no. 5, pp. 336–344, 1975.
3. M. Grabiec, "Fixed points in fuzzy metric spaces," *Fuzzy Sets and Systems*, vol. 27, no. 3, pp. 385–389, 1988.
4. A. George and P. Veeramani, "On some results in fuzzy metric spaces," *Fuzzy Sets and Systems*, vol. 64, no. 3, pp. 395–399, 1994.
5. C. Di Bari and C. Vetro, "Fixed points, attractors and weak fuzzy contractive mappings in a fuzzy metric space," *Journal of Fuzzy Mathematics*, vol. 13, no. 4, pp. 973–982, 2005.
6. D. Gopal, M. Imdad, C. Vetro, and M. Hasan, "Fixed point theory for cyclic weak ϕ -contraction in fuzzy metric spaces," *Journal of Nonlinear Analysis and Application*, vol. 2012, Article ID jnaa-00110, 11 pages, 2012.
7. P. Salimi, C. Vetro, and P. Vetro, "Some new fixed point results in non-Archimedean fuzzy metric spaces," *Nonlinear Analysis: Modelling and Control*, vol. 18, no. 3, pp. 344–358, 2013.
8. Y. Shen, D. Qiu, and W. Chen, "Fixed point theorems in fuzzy metric spaces," *Applied Mathematics Letters*, vol. 25, no. 2, pp. 138–141, 2012.
9. C. Vetro, "Fixed points in weak non-Archimedean fuzzy metric spaces," *Fuzzy Sets and Systems*, vol. 162, pp. 84–90, 2011.
10. C. Vetro, D. Gopal, and M. Imdad, "Common fixed point theorems for (ϕ, ψ) -weak contractions in fuzzy metric spaces," *Indian Journal of Mathematics*, vol. 52, no. 3, pp. 573–590, 2010.
11. C. Vetro and P. Vetro, "Common fixed points for discontinuous mappings in fuzzy metric spaces," *Rendiconti del Circolo Matematico di Palermo*, vol. 57, no. 2, pp. 295–303, 2008.
12. J. H. Park, "Intuitionistic fuzzy metric spaces," *Chaos, Solitons and Fractals*, vol. 22, no. 5, pp. 1039–1046, 2004.
13. C. Alaca, D. Turkoglu, and C. Yildiz, "Fixed points in intuitionistic fuzzy metric spaces," *Chaos, Solitons & Fractals*, vol. 29, no. 5, pp. 1073–1078, 2006.
14. D. Coker, "An introduction to intuitionistic fuzzy topological spaces," *Fuzzy Sets and Systems*, vol. 88, no. 1, pp. 81–89, 1997.
15. J. S. Park, Y. C. Kwun, and J. H. Park, "A fixed point theorem in the intuitionistic fuzzy metric spaces," *Far East Journal of Mathematical Sciences*, vol. 16, no. 2, pp. 137–149, 2005.
16. M. Rafi and M. S. M. Noorani, "Fixed point theorem on intuitionistic fuzzy metric spaces," *Iranian Journal of Fuzzy Systems*, vol. 3, no. 1, pp. 23–29, 2006.
17. B. Schweizer and A. Sklar, "Statistical metric spaces," *Pacific Journal of Mathematics*, vol. 10, pp. 313–334, 1960.
18. S. Manro, S. Kumar and S.S. Bhatia, Common fixed point theorems for weakly compatible maps satisfying common (E.A) like property in intuitionistic fuzzy metric spaces using implicit relation, *Journal of the Indian Math. Soc.*, 81(1-2) (2014), 123-133.
19. C. Ionescu, S. Rezapour, and M. E. Samei, "Fixed points of some new contractions on intuitionistic fuzzy metric spaces," *Fixed Point Theory and Applications*, vol. 2013, article 168, 2013.
20. N. Hussain, S. Khaleghizadeh, P. Salimi, and Afrah A. N. Abdou, "A New Approach to Fixed Point Results in Triangular Intuitionistic Fuzzy Metric Spaces", *Abstract and Applied Analysis*, Volume 2014, Article ID 690139, 16 pages.
21. N. Hussain, P. Salimi, V. Parvaneh, Fixed point results for various contractions in parametric and fuzzy b-metric spaces, *J. Nonlinear Sci. Appl.* 8 (2015), 719-739.
22. G. Jungck, "Fixed points for non-continuous non-self mappings on non-metric space," *Far East Journal of Mathematical Sciences*, vol. 4, pp. 199–212, 1996.
23. A. S. Saluja, A. K. Dhakda and D. Magarde, Some fixed point theorems for expansive type mapping in dislocated metric space, *Mathematical Theory and Modeling*, 3(2013).
24. R. D. Daheriya, Rashmi Jain, and Manoj Ughade "Some Fixed Point Theorem for Expansive Type Mapping in Dislocated Metric Space", *ISRN Mathematical Analysis*, Volume 2012, Article ID 376832, 5 pages, doi:10.5402/2012/376832.

25. Yan Han and Shaoyuan Xu, Some new theorems of expanding mappings without continuity in cone metric spaces, *Fixed Point Theory and Applications*, 2013, 2013:3.
26. Wasfi Shatanawi and Fadi Awawdeh, Some fixed and coincidence point theorems for expansive maps in cone metric spaces, *Fixed Point Theory and Applications* 2012, 2012:19.
27. Xianjiu Huang, Chuanxi Zhu, Xi Wen, Fixed point theorems for expanding mappings in partial metric spaces, *An. St. Univ. Ovidius Constant_a* Vol. 20(1), 2012, 213-224.
28. Z. Mustafa, F. Awawdeh and W. Shatanawi, Fixed Point Theorem for Expansive Mappings in G -Metric Spaces, *Int. J. Contemp. Math. Sciences*, Vol. 5, 2010, no. 50, 2463-2472.
29. A. Muraliraj and R. Jahir Hussain, Coincidence and Fixed Point Theorems for Expansive Maps in d -Metric Spaces, *Int. Journal of Math. Analysis*, Vol. 7, 2013, no. 44, 2171 – 2179.
30. P. Z. Daffer, H. Kaneko, *On expansive mappings*, *Math. Japonica*. 37 (1992), 733-735.
31. S. Z. Wang, B. Y. Li, Z. M. Gao, K. Iseki, *Some fixed point theorems for expansion mappings*, *Math. Japonica*. 29 (1984), 631-636.
32. Aage, CT, Salunke, JN: Some fixed point theorems for expansion onto mappings on cone metric spaces. *Acta Math. Sin. Engl. Ser.* 27(6), 1101-1106 (2011).
33. R. D. Daheriya, Rashmi Jain, and Manoj Ughade, Unified fixed point theorems for non-surjective expansion mappings by using class Ψ on 2- metric spaces, *Bessel J. Math.*, 3(3), (2013), 271-291.
34. G. Jungck, "Commuting mappings and fixed points," *The American Mathematical Monthly*, vol. 83, no.4, pp. 261-263, 1976.

Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2015. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]