

$I - g\alpha$ - CLOSED SETS IN IDEAL BITOPOLOGICAL SPACES

Stella Irene Mary. J*¹, Vinodhini. R.²

¹Associate Professor, ²M. phil. Scholar, Department of Mathematics,
PSG college of Arts and Science, Coimbatore - 641014, India.

(Received On: 29-10-15; Revised & Accepted On: 20-11-15)

ABSTRACT

In this paper a new class of closed sets namely $I - g\alpha$ - closed sets in ideal bitopological spaces (X, τ_1, τ_2, I) is introduced. Several properties and characterizations of this new class are investigated.

Keywords: Bitopological spaces, Ideal bitopological spaces, $(i, j) - g\alpha$ - closed sets, $(i, j) - I - g\alpha$ - closed sets, $(i, j) - I - g\alpha$ - open sets.

AMS subject classification: 54A05, 54E55, 54H05.

1. INTRODUCTION

The concept of Bitopological Spaces was initiated by Kelly [7] in 1963. He defined a set equipped with two topologies which is called a bitopological space and denoted it by (X, τ_1, τ_2) where (X, τ_1) and (X, τ_2) are the topological spaces. In 1970, Levine [9] introduced g - closed sets in Topological space (X, τ) and studied several properties. In 1986, Fukutake [2] extended the concept of generalized closed sets to bitopological spaces. In 1965, Njasted [12] and in 1990, Jelic [6] developed the concepts of alpha open sets and α - continuous functions in bitopological spaces. As an extension of α - open sets Maki *et.al* [10] defined $g\alpha$ - closed sets in topological spaces (X, τ) . In 2005, El - Tantawy and Abu.Donia [1] extended ag - closed sets induced by open sets which contains the class of $g\alpha$ - closed sets to bitopological spaces and studied several properties.

The concept of Ideal topological spaces was first introduced by Kuratowski [8] and thus opened the door for a large area of research. If I is an Ideal on X , then (X, τ_1, τ_2, I) is called an Ideal bitopological space. Vaidynathasamy [16] constructed local function in ideal topological space in 1945. Recently Tripathy and Hazarika [14], Tripathy and Mahanta [15] studied about the Ideal convergence in sequence spaces. In this article a new class of closed sets namely $I - g\alpha$ - closed sets in bitopological space is introduced. It is denoted by $(i, j) - I - g\alpha$ - closed sets where (i, j) means the topologies (τ_i, τ_j) , $i \neq j$; $i, j = 1, 2$ this class satisfies the inclusion relation given below.
 $\{(i, j) - g\alpha$ - closed sets $\} \subseteq \{(i, j) - I - g\alpha$ - closed sets $\}$

2. PRELIMINARIES:

Definition 2.1.1: [5] A **topology** on a set X is a collection of subsets of X having the following properties:

- 1) ϕ and X are in τ .
- 2) The union of the elements of any subcollection of τ is in τ .
- 3) The intersection of the elements of any finite subcollection of τ is in τ .

Definition 2.1.2: [7] A set X with topologies τ_1 and τ_2 is said to be **Bitopological space** and is denoted by (X, τ_1, τ_2) .

Definition 2.1.3: [13]

- (i) An **ideal I** on a non-empty set X is a collection of subsets of X which satisfies
- (a) $A \in I$ and $B \subseteq A$ implies $B \in I$ and
 - (b) $A \in I$ and $B \in I$ implies $A \cup B \in I$.

If I is an ideal on X , then (X, τ_1, τ_2, I) is called an **ideal bitopological space**.

- (ii) A subset A of X is said to be **$(i, j) - \text{clopen set}$** if the set is both open and closed.
-

Corresponding Author: Stella Irene Mary. J*¹

**¹Associate Professor, Department of Mathematics,
PSG college of Arts and Science, Coimbatore - 641014, India.**

Definition 2.1.4: [4] Let (X, τ_1, τ_2) be a bitopological space. We denote the bitopologies (τ_i, τ_j) by (i, j) .

- (i) A subset A of X is said to be (τ_i, τ_j) - α - open set or simply (i, j) - α - open set if $A \subseteq \tau_i - \text{int}[\tau_j - \text{cl}(\tau_i - \text{int}(A))]$, where $\text{int}(A)$ denote the interior of (A) and $\text{Cl}(A)$ denote the closure of (A) .
- (ii) A subset A of X is said to be (τ_i, τ_j) - α - closed set or simply (i, j) - α - closed set if $\tau_i - \text{cl}[\tau_j - \text{int}(\tau_i - \text{cl}(A))] \subseteq A$.
- (iii) The (i, j) - α - interior of A is defined as the union of all (i, j) - α - open sets contained in A . We denote (i, j) - α - interior of A by (i, j) - α - $\text{int}(A)$.
- (iv) The (i, j) - α - closure of A is defined as the intersection of all (i, j) - α - closed sets containing A . We denote (i, j) - α - closure of A by (i, j) - α - $\text{cl}(A)$.

3.1 Generalized α - closed sets in ideal bitopological spaces

Let (X, τ_1, τ_2) be a bitopological space. Throughout this chapter we denote the bitopologies (τ_i, τ_j) by (i, j) .

Definition 3.1.1: [1] A subset A of a bitopological space (X, τ_1, τ_2) is said to be (i, j) -generalized α – closed (In short, (i, j) - $g\alpha$ - closed) set if (j, i) - $\alpha\text{cl}(A) \subset U$ whenever $A \subset U$ and U is τ_i - α open in X .

In this section we prove several properties regarding (i, j) - α - interior and (i, j) - α - closure of sets in (X, τ_1, τ_2) .

Lemma 3.1.1: Let (X, τ_1, τ_2) be a bitopological space and A be a subset of X . Then

- (i) (i, j) - $\alpha\text{int}(A)$ is (i, j) - α - open.
- (ii) (i, j) - $\alpha\text{cl}(A)$ is (i, j) - α - closed.
- (iii) A is (i, j) - α - open if and only if $A = (i, j)$ - $\alpha\text{int}(A)$.
- (iv) A is (i, j) - α - closed if and only if $A = (i, j)$ - $\alpha\text{cl}(A)$.

Proof:

- (i) (i, j) - $\alpha\text{int}(A)$ is the union of all (i, j) - α - open sets contained in A . First we prove that (i, j) - α - open set is (i, j) - α - open. [Let $\{A_k\}$ where $k \in J$ be a family of (i, j) - α - open sets contained in A . Then for each $k \in J$,

$$\begin{aligned} A_k &\subseteq \tau_i - \text{int}[\tau_j - \text{cl}(\tau_i - \text{int} A_k)] \\ \text{Now, } \bigcup_k A_k &\subseteq \bigcup_k \{\tau_i - \text{int}[\tau_j - \text{cl}(\tau_i - \text{int} A_k)]\} \\ &\subseteq \tau_i - \text{int}[\bigcup_k \tau_j - \text{cl}(\tau_i - \text{int} A_k)] \\ &= \tau_i - \text{int}[\tau_j - \text{cl}(\bigcup_k \tau_i - \text{int} A_k)] \\ &\subseteq \tau_i - \text{int}[\tau_j - \text{cl}(\tau_i - \text{int} (\bigcup_k A_k))] \end{aligned}$$

Hence $\bigcup_k A_k$ is a (i, j) - α - open set

Therefore, (i, j) - $\alpha\text{int}(A)$ is (i, j) - α - open.

- (ii) (i, j) - $\alpha\text{cl}(A)$ is defined by intersection of all (i, j) - α - closed sets containing A . [Let $\{A_k\}$ where $k \in J$ is a family of (i, j) - α - closed sets containing A . This implies $\{A_k^c\}$ is a family of (i, j) - α - open sets contained in A^c . $\bigcup_k A_k^c$ is (i, j) - α - open set [by (i)]. Consequently $(\bigcap_k A_k)^c$ is (i, j) - α - open set and $\bigcap_k A_k$ (i, j) - α - closed set]. Therefore, (i, j) - $\alpha\text{cl}(A)$ is (i, j) - α - closed.
- (iii) Since (i, j) - $\alpha\text{int}(A)$ is the largest (i, j) - α - open set contained in A and A is (i, j) - α - open, we have $A = (i, j)$ - $\alpha\text{int}(A)$. Conversely, suppose $A = (i, j)$ - $\alpha\text{int}(A)$ then by (i), A is (i, j) - α - open.
- (iv) We know that, (i, j) - $\alpha\text{cl}(A)$ is the smallest (i, j) - α - closed set containing A . Since A itself is (i, j) - α - closed set, $A = (i, j)$ - $\alpha\text{cl}(A)$. Conversely, suppose $A = (i, j)$ - $\alpha\text{cl}(A)$ then by (ii), A is (i, j) - α - closed.

Lemma 3.1.2: Let (X, τ_1, τ_2) be a bitopological space and A be a subset of X . Then the following assertion hold.

- (i) $x \in (i, j)$ - $\alpha\text{cl}(A)$ if and only if for every (i, j) - α - open set U containing x , $U \cap A \neq \phi$.
- (ii) $x \in (i, j)$ - $\alpha\text{int}(A)$ if and only if there exists an (i, j) - α - open set U such that $x \in U \subseteq A$.
- (iii) If $A \subset B$, then (a) (i, j) - $\alpha\text{int}(A) \subset (i, j)$ - $\alpha\text{int}(B)$ and (b) (i, j) - $\alpha\text{cl}(A) \subset (i, j)$ - $\alpha\text{cl}(B)$

Proof:

- (i) Suppose $x \in (i, j)$ - $\alpha\text{cl}(A) = \bigcap_k B_k$, where B_k is a (i, j) - α - closed set containing A

Let U be an $(i, j) - \alpha$ - open set and $x \in U$ (3.1.1)

Suppose $U \cap A = \phi$ then $U^C \supset A$ and U^C is $(i, j) - \alpha$ - closed set containing A and hence $U^C = B_k$ for some k . Consequently, $x \in U^C$ which is a contradiction to (3.1.1). Hence $U \cap A \neq \phi$.

On the other hand, suppose for every $(i, j) - \alpha$ - open set U containing $x, U \cap A \neq \phi$. Suppose let $x \notin (i, j) - \alpha cl(A)$ then $x \notin \bigcap_k B_k$, where B_k is a $(i, j) - \alpha$ - closed set containing A . That means there exists $B_k \supset A$ such that $x \notin B_k$ and $x \in B_k^C$ where B_k^C is a $(i, j) - \alpha$ - open set contained in A and $B_k^C \cap A = \phi$ which is a contradiction. Therefore $x \in (i, j) - \alpha cl(A)$.

(ii) It is obvious by the definition of $(i, j) - \alpha int(A)$.

(iii) (a) $(i, j) - \alpha int(A) = \cup A_k$, where A_k is a $(i, j) - \alpha$ - open set contained in A (3.1.2)
 Since $A_k \subset A \subset B$ for all $k, \cup A_k \subset B$, by (3.1.2), $(i, j) - \alpha int(A) \subseteq B$. The fact that $(i, j) - \alpha int(A)$ is an $(i, j) - \alpha$ - open set and $(i, j) - \alpha int(B)$ is the largest $(i, j) - \alpha$ - open set contained in B , implies $(i, j) - \alpha int(A) \subset (i, j) - \alpha int(B)$.

(b) $A \subset B \Rightarrow B^C \subset A^C$. By (a), $(i, j) - \alpha int(B^C) \subset (i, j) - \alpha int(A^C)$ (3.1.3)
 Since $(i, j) - \alpha int(B^C) = \cup_k B_k^C$ and $(i, j) - \alpha int(A^C) = \cup_s A_s^C$ where B_k^C and A_s^C are $(i, j) - \alpha$ - open sets contained in B^C and A^C respectively.

By (3.1.3), $\cup_k B_k^C \subset \cup_s A_s^C \Rightarrow (\cap_k B_k)^C \subset (\cap_s A_s)^C \Rightarrow \cap_s A_s \subseteq \cap_k B_k$

where A_s and B_k are $(i, j) - \alpha$ - closed sets. Therefore, $(i, j) - \alpha cl(A) \subset (i, j) - \alpha cl(B)$.

Lemma 3.1.3: Let (X, τ_1, τ_2) be a bitopological space and A be a subset of X . Then

- (i) $X \setminus (i, j) - \alpha int(A) = (i, j) - \alpha cl(X \setminus A)$
- (ii) $X \setminus (i, j) - \alpha cl(A) = (i, j) - \alpha int(X \setminus A)$

Proof:

(i) $X \setminus (i, j) - \alpha int(A) = X \setminus \cup_k A_k$ where A_k is a $(i, j) - \alpha$ - open set contained in A
 $= \cap_k (X \setminus A_k)$
 $= \cap_k A_k^C$ where A_k^C is $(i, j) - \alpha$ - closed set containing A^C
 $= (i, j) - \alpha cl(A^C)$

(ii) $X \setminus (i, j) - \alpha cl(A) = X \setminus \cap_s A_s$ where A_s is a $(i, j) - \alpha$ - closed set containing A
 $= \cup (X \setminus A_s)$
 $= \cup A_s^C$ where A_s^C is $(i, j) - \alpha$ - open set contained in A^C
 $= (i, j) - \alpha int A^C$

3.2 $(i, j) - I$ - generalized α - closed sets

As an extension of $(i, j) -$ generalized α - closed sets in bitopological space to the ideal bitopological space we introduce the following definition.

Definition 3.2.1: Let (X, τ_1, τ_2, I) be an ideal bitopological space. A subset A of X is said to be **$(i, j) - I$ - generalized α - closed** (in short, $(i, j) - I - g\alpha$ - closed) set if $(j, i) - \alpha cl(A) \setminus B \in I$ whenever $A \subset B$ and B is $\tau_i - \alpha$ open in X , for $i, j = 1, 2$ and $i \neq j$. The family of all $(i, j) - I - g\alpha$ - closed sets of X is denoted by $(i, j) - I - g\alpha C(X)$. This new class $(i, j) - I - g\alpha$ - closed set contains the class of $(i, j) - g\alpha$ - closed set.

Definition 3.2.2: Let (X, τ_1, τ_2, I) be an ideal bitopological space and $A \subseteq X$, the intersection of all $(i, j) - I - g\alpha$ - closed sets containing A is called **$(i, j) - I - g\alpha$ - closure** of A and is denoted by $(i, j) - I - g\alpha - Cl(A)$.

The inclusion relation $\{(i, j) - g\alpha$ - closed sets $\} \subseteq \{(i, j) - I - g\alpha$ - closed sets $\}$ is proved in the following theorem.

Theorem 3.2.1: Every $(i, j) - g\alpha$ - closed is $(i, j) - I - g\alpha$ - closed.

Proof: Let A be $(i, j) - g\alpha$ - closed.

That is, $(j, i) - \alpha cl(A) \subset U$ whenever $A \subset U$ and U is $\tau_i - \alpha$ open in X (3.2.1)

We have to show that A is $(i, j) - I - g\alpha$ - closed. That is, $(j, i) - \alpha cl(A) \setminus B \in I$ whenever $A \subset B$ and B is $\tau_i - \alpha$ open in X , for $i, j = 1, 2$ and $i \neq j$. So, let $A \subset B$ and B is $\tau_i - \alpha$ open in X . By (3.2.1), $(j, i) - \alpha cl(A) \subset B$, and hence $(j, i) - \alpha cl(A) \setminus B = \phi \in I$. Thus A is $(i, j) - I - g\alpha$ - closed.

Remark 3.2.1: The converse of the above theorem is not necessarily true. This is proved in the following example.

Example 3.2.1: Let $X = \{a, b, c\}$. Consider the topologies $\tau_1 = \{\phi, \{a\}, X\}$, $\tau_2 = \{\phi, \{a\}, \{a, b\}, X\}$ and $I = \{\phi, \{b\}, \{c\}, \{b, c\}\}$. Here $\{a\}$ is $(1, 2) - I - g\alpha$ - closed set but not $(1, 2) - g\alpha$ - closed. Since $(2, 1) - \alpha cl(\{a\}) = X$ not a subset of $\{a\}$.

Theorem 3.2.2: Let (X, τ_1, τ_2, I) be an ideal bitopological space. If A is $(i, j) - I - g\alpha$ - closed and $A \subset B \subset (j, i) - \alpha cl(A)$ in X , then B is $(i, j) - I - g\alpha$ - closed in X , where $i, j = 1, 2$ and $i \neq j$.

Proof: Let $B \subset V$ and V is $\tau_i - \alpha$ open in X . Since $A \subset B \subset (j, i) - \alpha cl(A)$ and $B \subset V$, we have $A \subset V$. By hypothesis, $(j, i) - \alpha cl(A) \setminus V \in I$. Further $B \subset (j, i) - \alpha cl(A)$ implies that $(j, i) - \alpha cl(B) \subset (j, i) - \alpha cl(A) \Rightarrow (j, i) - \alpha cl(B) \setminus V \subset (j, i) - \alpha cl(A) \setminus V \in I$. Consequently B is $(i, j) - I - g\alpha$ - closed.

Set operations : The following Theorem and example prove that the class of $(i, j) - I - g\alpha$ - closed sets is closed for the set operation union but not for intersection.

Theorem 3.2.3: Union of two $(i, j) - I - g\alpha$ - closed sets in an ideal bitopological space (X, τ_1, τ_2, I) is also $(i, j) - I - g\alpha$ - closed.

Proof: Let A and B be two $(i, j) - I - g\alpha$ - closed sets with $A \cup B \subset V$, where V is any $\tau_i - \alpha$ open set. Clearly, $A \subset V$ and $B \subset V$ as $A \cup B \subset V$. Since A and B are $(i, j) - I - g\alpha$ - closed sets, we have $(j, i) - \alpha cl(A) \setminus V \in I$ and $(j, i) - \alpha cl(B) \setminus V \in I$. Now, $(j, i) - \alpha cl(A \cup B) \setminus V = ((j, i) - \alpha cl(A) \cup (j, i) - \alpha cl(B)) \setminus V = ((j, i) - \alpha cl(A) \setminus V) \cup ((j, i) - \alpha cl(B) \setminus V) \in I$ [since $A \in I$ and $B \in I \Rightarrow A \cup B \in I$]. Thus $(j, i) - \alpha cl(A \cup B) \setminus V \in I$. Hence $A \cup B$ is $(i, j) - I - g\alpha$ - closed set.

Remark 3.2.2: The intersection of two $(i, j) - I - g\alpha$ - closed sets need not be $(i, j) - I - g\alpha$ - closed. This is proved in the following example.

Example 3.2.2: Let $X = \{a, b, c\}$. Consider the topologies $\tau_1 = \{\phi, \{c\}, \{a, b\}, X\}$, $\tau_2 = \{\phi, \{a\}, \{b, c\}, X\}$ and $I = \{\phi\}$. Here $\{b, c\}$ and $\{a, c\}$ are $(1, 2) - I - g\alpha$ - closed set but $\{b, c\} \cap \{a, c\} = \{c\}$ is not $(1, 2) - I - g\alpha$ - closed.

Hereditary property : The following Theorem proves that the class of $(i, j) - I - g\alpha$ - closed sets admits hereditary property.

Theorem 3.2.4: Let (X, τ_1, τ_2, I) be an ideal bitopological space. Suppose A is $(i, j) - I - g\alpha$ - closed in X and $A \subset Y \subset X$. Then A is $(i, j) - I - g\alpha$ - closed relative to the subspace Y of X and with respect to the ideal $I_Y = \{P \subset Y : P \in I\}$.

Proof: Let V be $\tau_i - \alpha$ open in X and $A \subset Y \cap V$. Therefore, we have $A \subset V$. Since A is $(i, j) - I - g\alpha$ - closed in X , we have $(j, i) - \alpha cl(A) \setminus V \in I$. Further, we see that $((j, i) - \alpha cl(A) \cap Y) \setminus (Y \cap V) = ((j, i) - \alpha cl(A) \setminus V) \cap Y \in I_Y$.

Thus for $A \subset Y \cap V$ and V is $\tau_i - \alpha$ open, we have $((j, i) - \alpha cl(A) \cap Y) \setminus (Y \cap V) \in I_Y$. Hence A is $(i, j) - I - g\alpha$ - closed relative to the subspace $(Y, \tau_1|_Y, \tau_2|_Y)$.

3.3 $(i, j) - I -$ generalized α - open sets

We introduce the following definition namely $(i, j) - I -$ generalized α - open set.

Definition 3.3.1: Let (X, τ_1, τ_2, I) be an ideal bitopological space. A subset A of X is said to be $(i, j) - I -$ generalized α - open (in short, $(i, j) - I - g\alpha$ - open) set if $X \setminus A$ is $(i, j) - I - g\alpha$ - closed, for $i, j = 1, 2$ and $i \neq j$. The family of all $(i, j) - I - g\alpha$ - open sets contained in A is denoted by $(i, j) - I - g\alpha - O(X)$.

Lemma 3.3.1: Let (X, τ_1, τ_2, I) be an ideal bitopological space. Then $(j, i) - \alpha Cl(X \setminus A) = X \setminus (j, i) - \alpha int(A)$.

Proof: Let $x \in (j, i) - \alpha Cl(X \setminus A)$. That is for any $(j, i) - \alpha$ - open set U with $x \in U$, we have $U \cap (X \setminus A) \neq \phi$. This implies that there is no $(j, i) - \alpha$ - open set U with $x \in U$ so that $U \subseteq A$. Therefore, x is not $(j, i) - \alpha$ - interior point of A .

Thus $x \in X \setminus (j, i) - \alpha int(A)$. On the other hand, let $x \in X \setminus (j, i) - \alpha int(A)$. This implies $x \in X$ and $x \notin (j, i) - \alpha int(A)$. Since $x \notin (j, i) - \alpha int(A)$, there is no $(j, i) - \alpha$ - open set U with $x \in U$ so and $U \subseteq A$. We have $U \not\subseteq A$. Thus for any $(j, i) - \alpha$ - open set U with $x \in U$ we see that $U \cap (X \setminus A) \neq \phi$ which means that $x \in (j, i) - \alpha Cl(X \setminus A)$.

The following Theorem gives a Characterization of $(i, j) - I - g\alpha$ - open sets in an Ideal bitopological space.

Theorem 3.3.2: Let (X, τ_1, τ_2, I) be an ideal bitopological space. A subset A of X is $(i, j) - I - g\alpha$ - open in X if and if only $(B \setminus P) \subset (j, i) - \alpha int(A)$ for some $P \in I$, whenever $B \subset A$ and B is $\tau_i - \alpha$ closed.

Proof: Suppose A is $(i, j) - I - g\alpha$ - open in X . Let $B \subset A$ and B is $\tau_i - \alpha$ closed. Clearly $X \setminus A \subset X \setminus B$ and $X \setminus B$ is $\tau_i - \alpha$ open. Since A is $(i, j) - I - g\alpha$ - open, we have $X \setminus A$ is $(i, j) - I - g\alpha$ - closed. By definition, $(j, i) - \alpha cl(X \setminus A) \setminus (X \setminus B) \in I$. This implies $(j, i) - \alpha cl(X \setminus A) \cap (X \setminus B)^c = P$ for some $P \in I$
 $[(j, i) - \alpha cl(X \setminus A) \cap (X \setminus B)^c] \cup (X \setminus B) = P \cup (X \setminus B)$
 $[(j, i) - \alpha cl(X \setminus A) \cup (X \setminus B)] \cap [(X \setminus B)^c \cup (X \setminus B)] = P \cup (X \setminus B)$
 $(j, i) - \alpha cl(X \setminus A) \cup (X \setminus B) \cap X = P \cup (X \setminus B)$
 $(j, i) - \alpha cl(X \setminus A) \cup (X \setminus B) = P \cup (X \setminus B)$

This implies $(j, i) - \alpha cl(X \setminus A) \subset (X \setminus B) \cup P$, for some $P \in I$. Consequently $X \setminus ((X \setminus B) \cup P) \subset X \setminus (j, i) - \alpha cl(X \setminus A)$. By Lemma 3.3.1, we have $B \setminus P \subset X \setminus (X \setminus (j, i) - \alpha int(A))$. Hence $B \setminus P \subset (j, i) - \alpha int(A)$.

Conversely, Let $X \setminus A \subset B$ where B is $\tau_i - \alpha$ open in X , then $X \setminus B \subset A$ where B is $\tau_i - \alpha$ closed. By hypothesis, $(X \setminus B) \setminus P \subset (j, i) - \alpha int(A)$, where $P \in I$ which implies $(j, i) - \alpha cl(X \setminus A) \subset X \setminus [(X \setminus B) \setminus P] \subset B \cup P$. Therefore, $(j, i) - \alpha cl(X \setminus A) \setminus B \subset P \in I$ which implies $(j, i) - \alpha cl(X \setminus A) \setminus B \in I$ whenever $X \setminus A \subset B$ and B is $\tau_i - \alpha$ open $\Rightarrow X \setminus A$ is $(i, j) - I - g\alpha$ - closed in $X \Rightarrow A$ is $(i, j) - I - g\alpha$ - open in X .

Theorem 3.3.3: Let (X, τ_1, τ_2, I) be an ideal bitopological space. If A is $(i, j) - I - g\alpha$ - open in X and $(j, i) - \alpha int(A) \subset B \subset A$, then B is $(i, j) - I - g\alpha$ - open in X .

Proof: Assume that A is $(i, j) - I - g\alpha$ - open. Then $X \setminus A$ is $(i, j) - I - g\alpha$ - closed

We have $X \setminus A \subset X \setminus B \subset X \setminus (j, i) - \alpha int(A)$ (3.2.2)

$X \setminus (j, i) - \alpha int(A) = (j, i) - \alpha cl(X \setminus A)$. By (3.3.1), $X \setminus A \subset X \setminus B \subset (j, i) - \alpha cl(X \setminus A)$. By Theorem 3.3.1, we have $X \setminus B$ is $(i, j) - I - g\alpha$ - closed. Hence B is $(i, j) - I - g\alpha$ - open.

Set operations:

Theorem 3.3.4: The intersection of two $(i, j) - I - g\alpha$ - open sets in an ideal bitopological space (X, τ_1, τ_2, I) is also $(i, j) - I - g\alpha$ - open.

Proof: Suppose A and B be two $(i, j) - I - g\alpha$ - open sets in X . Then $X \setminus A$ and $X \setminus B$ are $(i, j) - I - g\alpha$ - closed. Now, we have $X \setminus (A \cap B) = (X \setminus A) \cup (X \setminus B)$ is $(i, j) - I - g\alpha$ - closed, by Theorem 3.2.3. Hence $A \cap B$ is $(i, j) - I - g\alpha$ - open.

Remark 3.3.1: The union of $(i, j) - I - g\alpha$ - open sets need not be $(i, j) - I - g\alpha$ - open set. This is proved in the following example.

Example 3.3.1: Let $X = \{a, b, c\}$. Consider the topologies $\tau_1 = \{\phi, \{a\}, \{b, c\}, X\}$, $\tau_2 = \{\phi, \{b\}, \{a, c\}, X\}$ and $I = \{\phi\}$. Here $\{b\}$ and $\{c\}$ are $(1, 2) - I - g\alpha$ - open but $\{b, c\}$ is not $(1, 2) - I - g\alpha$ - open.

The following Theorem proves that the union of any two $(i, j) - I - g\alpha$ - open set is open with the condition.

Theorem 3.3.5: If A and B are two $(i, j) - I - g\alpha$ - open sets in X such that $(j, i) - \alpha cl(A) \cap B = \phi$ and $A \cap (j, i) - \alpha cl(B) = \phi$, then $A \cup B$ is $(i, j) - I - g\alpha$ - open.

Proof: Let A and B be two $(i, j) - I - g\alpha$ - open sets in X such that $(j, i) - \alpha cl(A) \cap B = \phi$ and $A \cap (j, i) - \alpha cl(B) = \phi$. Suppose V is $\tau_i - \alpha$ closed and $V \subset A \cup B$. Then $V \cap (j, i) - \alpha cl(A) \subset (A \cup B) \cap (j, i) - \alpha cl(A) = (A \cap (j, i) - \alpha cl(A)) \cup (B \cap (j, i) - \alpha cl(A))$. $V \cap (j, i) - \alpha cl(A) \subset A \cap (j, i) - \alpha cl(A) = A$. Similarly, $V \cap (j, i) - \alpha cl(B) \subset B \cap (j, i) - \alpha cl(B) = B$. Since A and B are two $(i, j) - I - g\alpha$ - open sets in X , we have

$(V \cap (j, i) - acl(A)) \setminus P \subset (j, i) - \alpha int(A)$ and $(V \cap (j, i) - acl(B)) \setminus Q \subset (j, i) - \alpha int(B)$ for some $P, Q \in I$.

This implies that $(V \cap (j, i) - acl(A)) \setminus (j, i) - \alpha int(A) \subset P \in I$

$$\Rightarrow (V \cap (j, i) - acl(A)) \setminus (j, i) - \alpha int(A) \in I \quad (3.2.3)$$

$$\text{Similarly, } (V \cap (j, i) - acl(B)) \setminus (j, i) - \alpha int(B) \in I \quad (3.3.3)$$

By (3.3.2) and (3.3.3), we have

$(V \cap (j, i) - acl(A)) \setminus (j, i) - \alpha int(A) \cup (V \cap (j, i) - acl(B)) \setminus (j, i) - \alpha int(B) \in I$. Since for any two sets $A, B \in I$

$\Rightarrow A \cup B \in I$. Since for any sets A, B, C, D we have, $(A \cup C) \setminus (B \cup D) \subset (A \setminus B) \cup (C \setminus D)$

where $A = (V \cap (j, i) - acl(A))$; $B = (j, i) - \alpha int(A)$. $C = (V \cap (j, i) - acl(B))$; $D = (j, i) - \alpha int(B)$

$[(V \cap (j, i) - acl(A)) \cup (V \cap (j, i) - acl(B))] \setminus [(j, i) - \alpha int(A) \cup (j, i) - \alpha int(B)] \in I$. Also since

$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$, we have $V \cap [(j, i) - acl(A) \cup (j, i) - acl(B)] \setminus (j, i) - \alpha int(A) \cup (j, i) - \alpha int(B) \in I$. Thus $[V \cap ((j, i) - acl(A \cup B))] \setminus (j, i) - \alpha int(A) \cup (j, i) - \alpha int(B) \in I$

[Since $(j, i) - acl(A) \cup acl(B) = (j, i) - acl(A \cup B)$]. Further, $V = V \cap (A \cup B)$,

$V \subset V \cap ((j, i) - acl(A \cup B))$. Therefore,

$V \setminus [(j, i) - \alpha int(A \cup B)] \subset [V \cap ((j, i) - acl(A \cup B))] \setminus [(j, i) - \alpha int(A \cup B)] \subset [V \cap ((j, i) - acl(A \cup B))] \setminus ((j, i) - \alpha int(A) \cup (j, i) - \alpha int(B)) \in I$. This shows that $V \setminus R \subset (j, i) - \alpha int(A \cup B)$ for some $R \in I$

$\Rightarrow A \cup B$ is $(i, j) - I - g\alpha$ - open.

Theorem 3.3.6: Let (X, τ_1, τ_2, I) be an ideal bitopological space. If A is $(i, j) - I - g\alpha$ - open set relative to B such that $A \subset B \subset X$ and B is $(i, j) - I - g\alpha$ - open relative to X , then A is $(i, j) - I - g\alpha$ - open relative to X .

Proof: Let $U \subset A$ and U is $\tau_i - \alpha$ closed. Suppose A is $(i, j) - I - g\alpha$ - open relative to B . Then, we have $U \setminus P \subset (j, i) - \alpha int_B(A)$ for some $P \in I_B$, where $P \in I_B$ denotes the ideal of the set B . This implies that there exists a $(j, i) - \alpha$ open set V_1 , such that $U \setminus P \subset V_1 \cap B \subset A$. Let $U \subset B$ and U is $\tau_i - \alpha$ closed. Suppose B is $(i, j) - I - g\alpha$ - open relative to X . Then, we have $U \setminus Q \subset (j, i) - \alpha int(B)$ for some $Q \in I$. This implies that there exists a $(j, i) - \alpha$ open set V_2 , such that $U \setminus Q \subset V_2 \subset B$. Further $U \setminus (P \cup Q) = (U \setminus P) \cap (U \setminus Q) \subset ((V_1 \cap B) \cap V_2) \subset (V_1 \cap B) \cap B = V_1 \cap B \subset A$. This shows that $U \setminus (P \cup Q) \subset (j, i) - \alpha int(A)$ for some $P \cup Q \in I$. Hence A is $(i, j) - I - g\alpha$ - open relative to X .

REFERENCES:

1. O.A. El-Tantawy and H.M. Abu-Donia, Generalized Separation Axioms in Bitopological spaces, The Arabian JI for science and Engg. 30, 1A, (2005), 117-129.
2. T.Fukutake, On generalized closed sets in bitopological spaces, Bull.Fukuoka Univ.Ed.Part III, 35, (1985), 19- 28.
3. N.Gowrisankar and P.Senthil Kumar, Ideal bitopological $b-R_0(-R_1)$ spaces, International Journal of Innovative Science, Engineering & Technology, 2, (2015).
4. Q.H.Imran, Generalized Alpha Star Star Closed Sets in Bitopological Spaces, Gen. Math.Notes, 22, 2, (2014), 93-102.
5. J.R. Munkres, Topology, Massachusetts Institute of Technology, Pearson Education Inc.
6. M.Jelic, *Feebly p - continuous mappings*, Suppl.Rend Circ.Mat.Palemro (2), 24, (1990), 387-395.
7. J.C.Kelly, Bitopological spaces, Proc.London Math.Soc., 3 (13), (1963), 71-89.
8. K.Kuratowski, Topology, Academic Press, NewYork, (1966).
9. N.Levine, Generalized closed sets in Topology, Rend. Circ.Mat.Palermo, 19, (1970), 89-96.
10. H.Maki, R.Devi and K.Balachandran, Generalized alpha closed sets in topology, Bull Fukouka.Univ.Ed.part III, 42, (1993), 13-21.
11. Mashhour AS, Khedr FH and El-Deeb SN, Five separation axioms in bitopological spaces. Bull.Fac.Sci.Aasiut Univ.11, (1982), 53-67.
12. O.Njastad, On some class of nearly open sets, Pacific.J.Math., 15, (1965), 961-970.
13. B.C.Tripathy and D.J. Sarna, Generalized b - closed sets in ideal Bitopological spaces, Proyecciones Journal of Mathematics Vol.33, No. 3, (2014), 315-324.
14. B.C.Tripathy and B.Hazarika, I - convergent sequence spaces associated with multiplier Sequence spaces, Mathematical Inequalities and Applications; 11(3), (2008), 543-548.
15. B.C.Tripathy and S.Mahanta, On I - acceleration convergence of sequences, Journal of the Franklin Institute, 347, (2010), 591-598.
16. R.Vaidyanathaswamy, The localization theory in set topology, Proc.Indian Acad.Sci., 20, (1945), 51-61.
17. J.W.T.Young, A note on separation axioms and their application in the theory of locally Connected topological spaces. Bull.Amer.Math.Soc.49, (1943), 383-385.

Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2015. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]

