

## VERY $\beta_e$ - EXCELLENCE OF A GRAPH

N. SARADHA\*<sup>1</sup>, V. SWAMINATHAN<sup>2</sup>

<sup>1</sup>Assistant Professor, Department of Mathematics,  
SCSVMV University, Enathur, Kanchipuram, Tamilnadu, India.

<sup>2</sup>Coordinator, Ramanujan Research Center in Mathematics,  
Saraswathi Narayanan College, Madurai, Tamilnadu, India.

(Received On: 04-10-16; Revised & Accepted On: 22-10-16)

---

### ABSTRACT

Let  $G = (V, E)$  be a simple finite undirected graph. A subset  $S$  of  $V$  is called an equivalence set if every component of the induced sub graph  $\langle S \rangle$  is complete. The equivalence number  $\beta_e(G)$  is the maximum cardinality of an equivalence set of  $G$  [3]. A vertex  $u$  in  $V(G)$  is said to be  $\beta_e$ -good if  $u$  belongs to a  $\beta_e$ -set of  $G$ .  $G$  is said to be  $\beta_e$ -excellent if every vertex of  $G$  is  $\beta_e$ -good. A graph  $G = (V, E)$  is said to be very  $\beta_e$ -excellent if there exists a  $\beta_e$ -set  $S$  of  $G$  such that for every  $u$  in  $V-S$ , there exists a vertex  $v$  in  $S$  such that  $(S - \{v\}) \cup \{u\}$  is  $\beta_e$ -set of  $G$ .  $S$  is called a very  $\beta_e$ -excellent set of  $G$  and  $G$  is called a very  $\beta_e$ -excellent graph. An equivalence graph is a vertex disjoint union of complete graphs. The concept of equivalence set, sub chromatic number, generalized coloring and equivalence covering number were studied in [1], [2], [4], [5], [6], [8], [10]. In this paper the concept of very  $\beta_e$ -excellence is studied.

**Keywords:** Equivalence set, Equivalence graph,  $\beta_e$ -excellence, Very  $\beta_e$ -excellence.

---

### 1. INTRODUCTION

Gred.H. Fricke et al [7] called a vertex  $u$  of a graph  $G = (V, E)$  to be  $\mu$ -good if  $u$  is contained in a  $\mu(G)$ -set of  $G$  (where  $\mu$  is a parameter).  $G$  is said to be  $\mu$ -excellent if every vertex in  $V$  is  $\mu$ -good. A number of results has been proved by taking  $\mu$  as the domination parameter. Sridharan and Yamuna [12], [13] introduced several types of excellence, one of them being rigid excellence. A graph  $G$  is said to be rigid  $\mu$ -excellent if every vertex of  $G$  belongs to a unique  $\mu$ -set of  $G$ . Rigid  $\gamma$ -excellence was studied in [13]. A similar study was made with respect to the parameter  $\beta_0$  in [11]. A sub set  $S$  of  $V(G)$  is said to be an equivalence set if every component of  $\langle S \rangle$  is complete. A graph  $G$  is said to be an equivalence graph if  $V(G)$  is an equivalence set. The maximum cardinality of an equivalence set is denoted by  $\beta_e(G)$  [3]. In this paper, very  $\beta_e$ -excellence is defined and several results are derived.

### 2. Very $\beta_e$ -Excellence of a Graph


**Definition 2.1:** A graph  $G = (V, E)$  is said to be very  $\beta_e$ -excellent if there exists a  $\beta_e$ -set  $S$  of  $G$  such that for every  $u$  in  $V-S$ , there exists a vertex  $v$  in  $S$  such that  $(S - \{v\}) \cup \{u\}$  is a  $\beta_e$ -set of  $G$ .  $S$  is called a very  $\beta_e$ -excellent set of  $G$  and  $G$  is called a very  $\beta_e$ -excellent graph.

---

**Corresponding Author: N. Saradha\*<sup>1</sup>**

<sup>1</sup>Assistant Professor, Department of Mathematics,  
SCSVMV University, Enathur, Kanchipuram, Tamilnadu, India.

**Example 2.2:** Consider  $P_4$  with  $V(P_4) = \{u_1, u_2, u_3, u_4\}$ .



A graph which is very  $\beta_e$  -excellent

**Figure-2.1**

$S = \{u_1, u_2, u_4\}$  is a  $\beta_e$  -set of  $P_4$ . Also  $P_4$  is  $\beta_e$  -excellent.  $V - S = \{u_3\}$  and  $(S - \{u_2\}) \cup \{u_3\}$  is a  $\beta_e$  set of  $P_4$ . Therefore,  $S$  is a very  $\beta_e$  -excellent set of  $P_4$  and  $P_4$  is a very  $\beta_e$  -excellent graph.

**Remark 2.3:** Any very  $\beta_e$  -excellent graph is a  $\beta_e$  -excellent graph.

**Proof:** Let  $G$  be a very  $\beta_e$  -excellent graph and let  $S$  be a very  $\beta_e$  -excellent set of  $G$ . Let  $u \in V - S$ . Then there exist  $v \in S$  such that  $(S - \{v\}) \cup \{u\}$  is a  $\beta_e$  -set of  $G$ . Therefore, every vertex of  $V - S$  is an element of a  $\beta_e$  -set of  $G$ . Since  $S$  is a  $\beta_e$  -set of  $G$ , every element of  $V(G)$  is in a  $\beta_e$  -set of  $G$ . Therefore,  $G$  is  $\beta_e$  -excellent.

**Remark 2.4:** A very  $\beta_e$  -excellent graph need not be a rigid  $\beta_e$  -excellent graph. For example,  $P_4$  is a very  $\beta_e$  -excellent graph. But is not a rigid  $\beta_e$  -excellent graph.

**Very  $\beta_e$  -excellence for standard graphs**

1.  $K_n$  is very  $\beta_e$  -excellent for all  $n$ .
2.  $K_{1,n}$  is not a very  $\beta_e$  -excellent graph for any  $n \geq 2$ .
3.  $\overline{K_n}$  is a very  $\beta_e$  - excellent for all  $n$ .
4.  $W_n$  is not very  $\beta_e$  -excellent for  $n \geq 5$ .
5.  $K_{m,n}$  is not very  $\beta_e$  -excellent .
6. Petersen graph is not very  $\beta_e$  -excellent.
7. Any equivalence graph is very  $\beta_e$  -excellent.

**Proposition 2.5:**  $P_n$  is very  $\beta_e$  -excellent iff  $n = 2, 3, 4, 6, 7, 9, 12$ .

**Proof:** When  $n \equiv 2 \pmod{3}$ ,  $P_n$  is not  $\beta_e$  -excellent and hence not very  $\beta_e$  -excellent.

Therefore, the possible values of  $n$  are  $n = 15r, n = 15r + 1, n = 15r + 3, n = 15r + 4, n = 15r + 6, n = 15r + 7, n = 15r + 9, n = 15r + 10, n = 15r + 12, n = 15r + 13, n = 15r + 15 (r \geq 1)$ .

**Case I:**  $n = 15r$ . Let  $n = 3k$ . Then  $k = 5r$ ; If  $n = 3k$  then  $\beta_e = 2k = 10r$ .

Since the number of vertices is  $15r$ , there are  $3r$  consecutive five vertices set. For very  $\beta_e$  -excellence, from each set at most 3 vertices can be taken. Therefore, at most  $3(3r) = 9r$  vertices can be taken for constructing a very  $\beta_e$  -excellent set. But  $\beta_e(P_n) = 10r$  where  $n = 15r$ . Therefore,  $P_n$  where  $n = 15r$  is not very  $\beta_e$  excellent.

**Case II:**  $n = 15r + 1$ .

$n = 3k + 1$  implies  $k = 5r$ ;  $\beta_e = 2k + 1 = 10r + 1$ .

Since there are  $15r+1$  vertices, we have  $3r$  five consecutive element sets. From these sets as per the definition of very  $\beta_e$ -excellent set, at most 3 vertices can be taken from each set. The number of possible vertices chosen is  $3(3r)+1=9r+1$ . But  $\beta_e = 10r + 1$ . Therefore,  $P_n$  where  $n = 15r+1$  is not very  $\beta_e$ -excellent.

**Case III:**  $n = 15r + 3$

$n = 3k$  where  $k = 5r + 1$ ,  $\beta_e = 2k = 2(5r + 1)$ .

The number of possible vertices in a very  $\beta_e$ -excellent set chosen is  $3(3r) + 2 = 9r + 2$ .

Hence,  $P_n$  where  $n = 15r+3$  is not very  $\beta_e$ -excellent.

**Case IV:**  $n = 15r + 4$ .  $n = 3k + 1$  where  $k = 5r + 1$ ,  $\beta_e = 2k + 1 = 2(5r + 1) + 1 = 10r + 3$ .

The number of possible vertices chosen with respect to the definition of very  $\beta_e$ -excellent set is  $3(3r) + 3 = 9r + 3$ . But  $\beta_e = 10r + 3$ .

Therefore,  $P_n$  where  $n = 15r+4$  is not very  $\beta_e$ -excellent.

**Case V:**  $n = 15r + 6$ ,  $n = 3k$  where  $k = 5r + 2$ ,  $\beta_e = 2k = 2(5r + 2) = 10r + 4$ . There are  $3r+1$  five consecutive elements sets and from each set at most 3 vertices can be chosen is at most  $3(3r + 1) + 1 = 9r + 4$ . But  $\beta_e = 10r + 4$ . Therefore,  $P_n$  where  $n = 15r+6$  is not very  $\beta_e$ -excellent.

**Case VI:**  $n = 15r + 7$ ,  $n = 3k + 1$  where  $k = 5r + 2$ ,  $\beta_e = 2k + 1 = 2(5r + 2) + 1 = 10r + 5$ .

The number of maximum possible vertices chosen for a very  $\beta_e$ -excellent set is  $3(3r + 1) + 2 = 9r + 5$ . But  $\beta_e = 10r + 5$ . Therefore,  $P_n$  where  $n = 15r+7$  is not very  $\beta_e$ -excellent.

**Case VII:**  $n = 15r + 9$ ;  $n = 3k + 3$  where  $k = 5r + 3$ ,  $\beta_e = 2k = 2(5r + 3) = 10r + 6$ .

The number of possible vertices chosen for constructing a very  $\beta_e$ -excellent set is  $3(3r + 1) + 3 = 9r + 6$ . But  $\beta_e = 10r + 6$ . Therefore,  $P_n$  where  $n = 15r+9$  is not very  $\beta_e$ -excellent.

**Case VIII:**  $n = 15r + 10$ .  $n = 3k + 1$  where  $k = 5r + 3$ ;  $\beta_e = 2k + 1 = 2(5r + 3) + 1 = 10r + 7$ . The number of possible vertices chosen for constructing a very  $\beta_e$ -excellent set is  $3(3r + 2) = 9r + 6$ . But  $\beta_e = 10r + 7$ . Therefore,  $P_n$  where  $n = 15r+10$  is not very  $\beta_e$ -excellent.

**Case IX:**  $n = 15r + 12$ ,  $n = 3k$  where  $k = 5r + 4$ ;  $\beta_e = 2k = 2(5r + 4) = 10r + 8$ .

The number of possible vertices chosen for constructing a very  $\beta_e$ -excellent set is  $3(3r+2)+2=9r+8$ . But  $\beta_e = 10r + 8$ . Therefore,  $P_n$  where  $n = 15r + 12$  is not very  $\beta_e$ -excellent set.

**Case X:**  $n = 15r + 13$ .  $n = 3k + 1$  where  $k = 5r + 4$ ;  $\beta_e = 2k + 1 = 2(5r + 4) + 1 = 10r + 9$ .

The number of possible vertices chosen for constructing a very  $\beta_e$ -excellent set is  $3(3r+2)+2$ . But  $\beta_e = 10r + 9$ . Therefore  $P_n$  where  $n = 15r + 13$  is not very  $\beta_e$ -excellent set.

**Case XI:**  $n = 15r + 15$ .  $n = 3k$  where  $k = 5r + 5$ ;  $\beta_e = 2k = 2(5r + 5) = 10r + 10$ .

The number of possible vertices chosen for constructing a very  $\beta_e$ -excellent set is  $3(3r+5) = 9r+15$ . But  $\beta_e = 10r + 10$ . Therefore  $P_n$  where  $n = 15r + 15$  is not very  $\beta_e$ -excellent set.

When  $n = 1, 2, 3, 4$   $P_n$  is clearly very  $\beta_e$ -excellent.

When  $n = 6$ ,  $\{u_1, u_2, u_5, u_6\}$  is a very  $\beta_e$ -excellent set where  $V(P_6) = \{u_1, u_2, u_3, u_4, u_5, u_6\}$ .

When  $n = 7$ ,  $\{u_1, u_2, u_4, u_6, u_7\}$  is a very  $\beta_e$ -excellent set.

When  $n = 9$ ,  $\{u_1, u_2, u_4, u_6, u_7, u_9\}$  is a very  $\beta_e$ -excellent.

When  $n = 10$ ;  $n = 3k + 1$  where  $k = 3$ . There are two five consecutive elements set in  $P_{10}$  and at most 6 element are possible for a very  $\beta_e$ -excellent. Hence  $P_n$  is not very  $\beta_e$ -excellent.

When  $n = 12$ ,  $n = 3k$  where  $k = 4$ .  $\beta_e(P_n) = 8$ .

The set  $\{u_1, u_2, u_4, u_6, u_7, u_9, u_{11}, u_{12}\}$  is a very  $\beta_e$ -excellent and hence  $P_{12}$  is a very  $\beta_e$ -excellent graph.

When  $n = 13$ ,  $n = 3k+1$  where  $k=4$ .  $\beta_e(P_{13}) = 9$ . There are two five consecutive element sets with 3 elements remaining in the last. Hence at most 6 elements can be taken from the two consecutive elements sets and all the three remaining elements are to be taken for having 9 elements. This might will not give a  $\beta_e$ -set, since 3 consecutive elements cannot be taken in a  $\beta_e$ -set. Hence  $P_{13}$  is not very  $\beta_e$ -excellent.

**Proposition 2.6:**  $C_n$  is very  $\beta_e$ -excellent only if  $n = 3, 4, 5, 7, 10, 13$ .

**Proof:** Arguing as in the previous proposition 2.5 the above result is obtained.

**Remark 2.7:** If a graph  $G$  has a unique  $\beta_e$ -set which is not  $V(G)$  then  $G$  is not very  $\beta_e$ -excellent.

**Proposition 2.8:**  $C_n \circ K_1$  is not very  $\beta_e$ -excellent.

**Proof:**

**Case I:** Let  $n$  be even.

Let  $V(C_n \circ K_1) = \{u_1, u_2, \dots, u_n, v_1, v_2, \dots, v_n\}$ . Any  $\beta_e$ -set  $S$  of  $C_n \circ K_1$  consists of all  $v_i$ 's and alternate  $u_i$ 's. Any vertex outside  $S$  cannot come inside by replacing a vertex of  $S$  without affecting the equivalence nature of  $S$ . Therefore,  $C_n \circ K_1$  is not very  $\beta_e$ -excellent.

**Case II:** Let  $n$  be odd.

A similar argument as before shows that there exist no  $\beta_e$  excellent set which is very  $\beta_e$ -excellent.

**Observation 2.9:** A very  $\beta_e$ -excellent graph may have isolates. Also, there are non-equivalence graphs which have isolates and which are very  $\beta_e$ -excellent.

For example,  $K_m \cup \overline{K_n}$  is a very  $\beta_e$ -excellent graph which have isolates, but this is an equivalence graph.  $C_4 \cup K_1$  is a non equivalence graph which is very  $\beta_e$ -excellent and which has an isolate.

**Remark 2.10:** If G is a very  $\beta_e$ -excellent graph then  $G \cup \overline{K_m}$  is also very  $\beta_e$ -excellent.

**Proposition 2.11:** Let G be very  $\beta_e$ -excellent graph without isolates. Let S be a very  $\beta_e$ -excellent set of G. Then for any  $u \in S$ ,  $|pn[u, S]| \geq 1$ .

**Proof:** Let G be a very  $\beta_e$ -excellent graph and let S be a very  $\beta_e$ -excellent set of G. Let  $u \in S$ . Suppose u is an isolate of S and any neighbor of u in G is adjacent with some vertex of S other than u. Then  $pn[u, S] = 1$ . Also, if all the neighbors of u form a complete sub graph with u, then  $pn[u, S] = 1$ .

**Corollary 2.12:**  $P_6$  is very  $\beta_e$ -excellent.  $S = \{u_1, u_2, u_5, u_6\}$  is a very  $\beta_e$  excellent set of G and  $pn[u_5, S] = 2 > 1$

**Remark 2.13:** Let G be a graph without isolates. Let S be a very  $\beta_e$ -excellent set of G. Let  $x \in V - S$ . Then there exist  $u \in S$  such that  $(S - \{u\}) \cup \{x\}$  is a  $\beta_e$ -set of G. x need not be a private neighbour of u. For example,  $P_7$  is very  $\beta_e$ -excellent. Let  $V(P_7) = \{u_1, u_2, u_3, u_4, u_5, u_6, u_7\}$ . Let  $S = \{u_1, u_2, u_4, u_6, u_7\}$ . Then S is a very  $\beta_e$ -excellent subset of  $V(G)$ .  $(S - \{u_2\}) \cup \{u_3\}$  is a  $\beta_e$ -set of G. But  $u_3$  is not a private neighbour of  $u_2$ .

**Theorem 2.14:** Let G be a graph without isolates. Suppose there exist a  $\beta_e$ -set S of G such that for every  $x \in V - S$ , there exist  $u \in S$  such that  $x \in pn(u, S)$ . Then G is very  $\beta_e$ -excellent.

**Proof:** By hypothesis, there exist a  $\beta_e$ -set S of G such that for every  $x \in V - S$ , there exist  $u \in S$  such that  $x \in pn(u, S)$ . Then  $(S - \{u\}) \cup \{x\}$  is a  $\beta_e$ -set of G. Therefore S is a very  $\beta_e$ -excellent set of G. Hence G is a very  $\beta_e$ -excellent graph.

**Illustration 2.15:** Let  $V(C_4) = \{u_1, u_2, u_3, u_4\}$ . Let  $S = \{u_1, u_2\}$ . Then  $u_3$  and  $u_4$  are private neighbours of S.

$(S - \{u_2\}) \cup \{u_3\}$  is a  $\beta_e$ -set.

$(S - \{u_1\}) \cup \{u_4\}$  is a  $\beta_e$ -set.

**Theorem 2.16:** Let G be a graph such that G is an equivalence graph. Let  $V_1, V_2, \dots, V_k$  be the components of G which are complete. Add vertices  $u_1, u_2, \dots, u_k$ . Join  $u_i$  only with every vertex of  $V_i, 1 \leq i \leq k$ . Let H be the resulting graph. Then H is very  $\beta_e$ -excellent.

**Proof:** Clearly H is an equivalence graph and H is very  $\beta_e$ -excellent.

## REFERENCES

1. M.O. Albertson, R.E. Jamison, S.T. Hedetniemi and S.C.Locke, The subchromatic number of a graph, Discrete Math., 74(1989), 33-49.
2. N. Alon, Covering graphs by the minimum number of equivalence relations, Combinatorica 6 (3) (1986), 201-206.
3. S. Arumugam, M.Sundarakannan, Equivalence Dominating Sets in Graphs, Utilitas Mathematica 91(2013) 231-242.
4. A. Blokhuis and T. Kloks, On the equivalence covering number of split graphs, Information Processing Letters, 54(1995), 301-304.

5. P.Duchet, Representations, noyaux en theorie des graaphes et hyper graphes, These de Doctoral d'EtatI., Universite Paris VI, 1979.
6. R.D. Dutton and R.C. Brigham, Domination in Claw-Free Graphs, Congr. Numer., 132(1998), 69-75.
7. G.H. Fricke, Teresa W.Haynes, S.T. Hedetniemi, S.M. Hedetniemi and R.C. Laskar, Excellent trees.preprint.
8. J.Gimbel and C. Hartman, Subcolorings and the subchromatic number of a graph, Discrete Math., 272 (2003), 139-154.
9. Harary, Graph Theory,Addison-Wesley/Narosa,1988.
10. C.Mynhardt and I. Broere, Generalized colorings of graphs, In Y.Alavi, G.Chartrand, L.Lesniak, D.R. Lick and C.E. Wall, editors, Graph Theory with Applications to Algorithms and Computer Science, Wiley, (1985),583-594.
11. A.P.Pushpalatha, Study of graphs with respect to the independence parameter and forcing domination, Ph.D. Thesis, Madurai Kamaraj University, 2010.
12. N.Sridharan and M.Yamuna, A note of Excellent graphs, Ars Combinatoria 78 (2006), pp.267-276.
13. M.Yamuna, Excellent-Just Excellent-Very Excellent Graphs, Ph.D. Thesis, Alagappa University, Karaikudi, 2003.

**Source of support: Nil, Conflict of interest: None Declared**

***[Copy right © 2016. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]***