

ON FIXED POINT THEOREM IN A SPACE WITH TWO METRICS

GANESH KUMAR SONI*

Dept. of Mathematics, Govt. P. G. College, Narsinghpur - (M.P.), India.

(Received On: 14-12-16; Revised & Accepted On: 25-01-17)

ABSTRACT

In the present paper, we shall prove a fixed point theorem for mapping in two metric spaces.

Keywords: Two metric space, Continuous mapping, Fixed point.

INTRODUCTION

In 1968 Maia (1) generalized the result of well known Banach, S. (2) contraction Principle by taking two metrics on a set X . During the past few years Maia's theorem was generalized and fixed point theorems proved in several directions by Iseki, K. (3), Iyer, S. (4), Mishra (5) Ray, B.K. (6), Rus, I.A. (7), Singh, S.P. and Pant, S.P. (8) Bhore, S.K. (9) Phatak, H.K. (10) and others

OUR MAIN RESULTS

Theorem 1: Let (X, d_1, d_2) be metric space such that the following conditions hold,

- (i) $d_1(x, y) \leq d_2(x, y)$ for all $x, y \in X$.
- (ii) (X, d_1) is complete space.
- (iii) Two mappings $S, T : X \rightarrow X$ be a mapping are continuous with respect to d_1 and satisfies inequality

$$[d_2(Sx, Ty)]^2 \leq C \frac{d_2(x, y) d_2(Sx, Ty) [1 + 2\{\sqrt{d_2(x, y)} + d_2(y, Ty)\}^2]}{[1 + \{\sqrt{d_2(x, Ty)} + d_2(y, Sx)\}^2 + \{\sqrt{d_2(x, Sx)} + d_2(y, Ty)\}^2]}$$

Then S and T have a common fixed point, further if $0 < C < 1$ then each of S and T has a unique fixed point and these two fixed points coincide.

Proof: Let $x_0 \in X$ be an arbitrary and define a sequence $\{x_n\}$ be a sequence of X then we defined as below $x_{2m-1} = Sx_{2m-2}$, $x_{2m} = Tx_{2m-1}$, $m = 1, 2, 3, \dots$ then we show that the sequence $\{x_n\}$ of iterates at x_0 is a Cauchy sequence. If for some k be the positive integers such that $x_{k-1} = x_k$ then $\{x_n\}$ is a Cauchy sequence. Suppose that the positive $x_{m-1} \neq x_m$ for each $m = 1, 2, 3, \dots$ then for $x = x_{2m-2}$ and $y = x_{2m-1}$ using inequality (iv), we have

$$\begin{aligned} [d_2(x_{2m-1}, x_{2m})]^2 &\leq C \frac{d_2(x_{2m-2}, x_{2m-1}) d_2(x_{2m-1}, x_{2m}) [1 + 2\{\sqrt{d_2(x_{2m-2}, x_{2m-1})} + d_2(x_{2m-1}, x_{2m})\}^2]}{[1 + \{\sqrt{d_2(x_{2m-2}, x_{2m})} + d_2(x_{2m-1}, x_{2m-1})\}^2 + \{\sqrt{d_2(x_{2m-2}, x_{2m-1})} + d_2(x_{2m-1}, x_{2m})\}^2]} \\ &\leq C \frac{d_2(x_{2m-2}, x_{2m-1}) d_2(x_{2m-1}, x_{2m}) [1 + 2\{\sqrt{d_2(x_{2m-2}, x_{2m})}\}^2]}{[1 + \{\sqrt{d_2(x_{2m-2}, x_{2m})}\}^2 + \{\sqrt{d_2(x_{2m-2}, x_{2m})}\}^2]} \\ &\leq C \frac{d_2(x_{2m-2}, x_{2m-1}) d_2(x_{2m-1}, x_{2m}) [1 + 2d_2(x_{2m-2}, x_{2m})]}{[1 + 2d_2(x_{2m-2}, x_{2m})]} \end{aligned}$$

$$[d_2(x_{2m-1}, x_{2m})]^2 \leq C d_2(x_{2m-2}, x_{2m-1}) d_2(x_{2m-1}, x_{2m})$$

$$d_2(x_{2m-1}, x_{2m}) \leq C d_2(x_{2m-2}, x_{2m-1})$$

Where $C < 1$. Proceeding in this way we get $d_2(x_{2m-1}, x_{2m}) \leq C^{2m-1} d_2(x_{2m-2}, x_{2m-1})$

By calculations we can easily see that the following inequality holds for sequence $\{x_n\}$.

$$\begin{aligned} d_2(x_i, x_j) &\leq \sum_{k=1}^{j-1} d_2(x_k, x_{k+1}) \\ &\leq \left(\frac{C^i}{1-C}\right) d_2(x_0, x_1) \quad j < i \end{aligned}$$

Tends to zero as $i \rightarrow \infty$.

Corresponding Author: Ganesh Kumar Soni*

Dept. of Mathematics, Govt. P. G. College, Narsinghpur - (M.P.), India.

It follows that $\{x_n\}$ is a Cauchy sequence with respect to d_2 . Further by condition (i) sequence is a Cauchy under metric d_1 . Since X is complete space there is some u in X such that $\lim_{n \rightarrow \infty} x_n = p$ in the metric d_1 . Since S is continuous on X it is continuous at p also. $Sp = S \lim_{m \rightarrow \infty} x_n = S \lim_{m \rightarrow \infty} x_{2m} = \lim_{m \rightarrow \infty} Sx_{2m} = \lim_{m \rightarrow \infty} x_{2m+1} = p$

Similarly, by the continuity of T we have $Tp = p$. Hence p is the a common fixed point of S and T .

Let p and q be two distinct common fixed point of S and T then

$$\begin{aligned} [d_2(p, q)]^2 &= [d_2(Sp, Tq)]^2 \\ &\leq C \frac{d_2(p, q)d_2(Sp, Tq)[1+2\{\sqrt{d_2(p, q)+d_2(q, Tq)}\}^2]}{[1+\{\sqrt{d_2(p, Tq)+d_2(q, Sp)}\}^2+\{\sqrt{d_2(p, Sp)+d_2(q, Tq)}\}^2]} \\ &\leq C \frac{d_2(p, q)d_2(p, q)[1+2\{\sqrt{d_2(p, q)+d_2(q, q)}\}^2]}{[1+\{\sqrt{d_2(p, q)+d_2(q, p)}\}^2+\{\sqrt{d_2(p, p)+d_2(q, q)}\}^2]} \\ &= C[d_2(p, q)]^2 \end{aligned}$$

Since $0 < C < 1$ we arrive at a contradiction then $p = q$. Therefore S and T have unique common fixed point.

REFERENCES

1. Maia M.G., Un' osservazione sulle contrazioni metriche, Rend. Sem. Math. Univ. Padova 40, 139-143 (1968).
2. Banach S., Sur les operations dans les ensembles abstraits et leur application aux equations integrals Fund.Math 3, 133-181(1922).
3. Iseki K., A common fixed point theorem, Rend. Sem. Math. Padova 53, 13-14 (1975).
4. Iyer S., Fixed point theorems in Bimetric spaces, Journal of M.A.C.T., 15 (1982).
5. Mishra S.N., Remarks on some fixed point theorems in Bimetric spaces, Indian J.pure appl. Math 9(12), 1271-1274 (1979).
6. Ray B.K., On a fixed point theorem in a space with two metrics, Mathematics Education Vol. IX No. 3 (1975).
7. Rus I.A., On a fixed point theorem of Maia, Studia Univ. Babes - Bolyai 22, 40-42(1977).
8. Singh S.P. and Pant R.P., A common fixed point theorem in a metric space with two metrics, Pure appl. Math. Sci. 14 (1-2), 35-37(1981).
9. Bhore S.K., Fixed points of orbitally continuous mappings, The Mathematics Education Vol. XXI No. 4 Dec. 1987.
10. Pathak H.K., On some non-unique fixed point theorems for the maps of Dhage type, Pure and Appl. Math. Science Vol. XXVII No. 1-2 March 1988.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2016. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]