

SOME NEW CONCEPTS OF CONTINUITY IN TOPOLOGICAL SPACES

C. SANTHINI¹, S. LAKSHMI PRIYA*²

¹Associate Professor,
Department of Mathematics,
V. V. Vanniaperumal College for Women, Virudhunagar - 626 001, India.

²M.Phil Scholar,
Department of Mathematics,
V. V. Vanniaperumal College for Women, Virudhunagar - 626 001, India.

(Received On: 15-03-17; Revised & Accepted On: 10-04-17)

ABSTRACT

The purpose of this paper is to introduce and investigate several continuous functions namely gs_a^{**} -continuous functions and contra gs_a^{**} -continuous functions along with their several characterizations. Further we introduce new types of graphs called gs_a^{**} -closed graphs, contra gs_a^{**} -closed graphs and investigated several characterizations of such notions.

MSC: 54C08, 54C05.

Keywords: gs_a^{**} -continuous functions, contra gs_a^{**} -continuous functions, gs_a^{**} -closed graph, contra gs_a^{**} -closed graph, locally gs_a^{**} -indiscrete space.

1. INTRODUCTION

In recent literature, we find many topologists have focused their research in the direction of investigating types of generalized continuity. The notion of contra-continuity was first investigated by Dontchev[7]. A good number of researchers have initiated different types of contra-continuous functions which are found in the papers [4],[5],[6]. In 1970, Levine [10] discussed the notion of generalized closed sets in topological spaces. Extensive research on generalizing closedness was done in recent years. In 1963, Levine [11] introduced the concepts of semi-open sets in topological spaces. W. Dunham [9] introduced the concept of generalized closure and defined a new topology τ^* and investigated some of their properties. Quite recently the authors Robert.A and Pious Missier.S introduced and studied semi-open [15] sets and semi α -open [15] sets using the generalized closure operator. Recently Santhini *et.al* [16] introduced gs_a^{**} -closed sets in topological spaces. In 1969, Long [12] introduced closed graphs in topological spaces. In this paper, by means of gs_a^{**} -closed sets, we introduce namely, gs_a^{**} -continuous functions and contra gs_a^{**} -continuous functions along with their several properties, characterizations and mutual relationships. Further we introduce new types of graphs, called gs_a^{**} -closed graphs, contra gs_a^{**} -closed graphs via gs_a^{**} -open sets. Several characterizations and properties of such notions are investigated.

2. PRELIMINARIES

In this section, we recall some basic definitions and properties used in our paper.

Definition 2.1: A subset A of a space (X, τ) is said to be

- (i) semi-open [11] if $A \subseteq cl(intA)$.
- (ii) semi-open if [15] $A \subseteq cl_{\alpha}(intA)$.
- (iii) semi α -open [15] if $A \subseteq cl_{\alpha}(intA)$.
- (iv) a g-closed set [2] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is open in X.

*Corresponding Author: S. Lakshmi Priya*²*

²M.Phil Scholar, Department of Mathematics,

V. V. Vanniaperumal College for Women, Virudhunagar - 626 001, India.

- (v) a ω -closed set [17] if $\text{cl}(A) \subseteq U$ whenever A and U is semi-open in X .
- (vi) a generalized-semi closed set(briefly gs-closed) [5] if $\text{scl}(A) \subseteq U$ whenever $A \subseteq U$ and U is open in X .
- (vii) a g^*s -closed set[14] if $\text{scl}(A) \subseteq U$ whenever $A \subseteq U$ and U is gs-open in X .
- (viii) a generalized semi pre-closed set(briefly gsp-closed)[8] if $\text{spcl}(A) \subseteq U$ whenever $A \subseteq U$ and U is open in X .

Definition 2.2: A subset A of a space (X, τ) is called generalized gs_α^{**} -closed set (briefly gs_α^{**} -closed) [16] if $\text{scl}(A) \subseteq U$ whenever $A \subseteq U$ and U is semi $^*\alpha$ -open in (X, τ) .

The class of all gs_α^{**} -open subsets of X is denoted by $gs_\alpha^{**}O(X, \tau)$ and the class of all gs_α^{**} -open subsets of X containing x is denoted by $gs_\alpha^{**}O(X, x)$.

Definition 2.3: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called a

- (1) semi-continuous [11] if $f^{-1}(V)$ is semi-closed set in (X, τ) for every closed set V in (Y, σ) .
- (2) semi * -continuous [13] if $f^{-1}(V)$ is semi * -closed set in (X, τ) for every closed set V in (Y, σ) .
- (3) semi $^*\alpha$ -continuous [15] if $f^{-1}(V)$ is semi $^*\alpha$ -closed set in (X, τ) for every closed set V in (Y, σ) .
- (4) g -continuous [2] if $f^{-1}(V)$ is g -closed set in (X, τ) for every closed set V in (Y, σ) .
- (5) generalized semi-continuous(briefly gs-continuous) [5] if $f^{-1}(V)$ is gs-closed set in (X, τ) for every closed set V in (Y, σ) .
- (6) generalized semi-precontinuous (briefly gsp-continuous) [8] if $f^{-1}(V)$ is gsp-closed set in (X, τ) for every closed set V in (Y, σ) .
- (7) ω -continuous [17] if $f^{-1}(V)$ is ω -closed set in (X, τ) for every closed set V in (Y, σ) .
- (8) g^*s -continuous [14] if $f^{-1}(V)$ is g^*s -closed set in (X, τ) for every closed set V in (Y, σ) .

Definition 2.4: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is said to be

- (1) contra-continuous [7] if $f^{-1}(V)$ is closed in (X, τ) for every open set V in (Y, σ) .
- (2) contra semi-continuous [6] if $f^{-1}(V)$ is semi-closed in (X, τ) for every open set V in (Y, σ) .
- (3) contra semi * -continuous [13] if $f^{-1}(V)$ is semi * -closed in (X, τ) for every open set V in (Y, σ) .
- (4) contra semi $^*\alpha$ -continuous [15] if $f^{-1}(V)$ is semi $^*\alpha$ -closed in (X, τ) for every open set V in (Y, σ) .
- (5) contra gs-continuous [3] if $f^{-1}(V)$ is gs-closed in (X, τ) for every open set V in (Y, σ) .
- (6) contra gsp-continuous [1] if $f^{-1}(V)$ is gsp-closed in (X, τ) for every open set V in (Y, σ) .
- (7) contra g -continuous [4] if $f^{-1}(V)$ is g -closed in (X, τ) for every open set V in (Y, σ) .
- (8) contra g^*s -continuous [14] if $f^{-1}(V)$ is g^*s -closed in (X, τ) for every open set V in (Y, σ) .

Definition 2.5: A space X is locally indiscrete [18] if every open set in X is closed.

Definition 2.6:

- (i) A space (X, τ) is called a ${}_aT_{s^{**}}$ -space [16] if every gs_α^{**} -closed set in it is closed.
- (ii) A space (X, τ) is called a $T_{s^{**}}^\alpha$ -space [16] if every gs-closed set in it is gs_α^{**} -closed.

3. gs_α^{**} -Continuous and gs_α^{**} -Irresolute functions

In this section, the concepts of gs_α^{**} -continuity and gs_α^{**} -irresoluteness are introduced and studied.

Definition 3.1: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called gs_α^{**} -continuous if $f^{-1}(V)$ is gs_α^{**} -closed set in (X, τ) for every closed set V in (Y, σ) .

Example 3.2: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}, \{a\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = c, f(b) = a, f(c) = b$ is gs_α^{**} -continuous.

Theorem 3.3:

- (1) Every continuous function is gs_α^{**} -continuous.
- (2) Every ω -continuous function is gs_α^{**} -continuous.
- (3) Every g^*s -continuous function is gs_α^{**} -continuous.
- (4) Every semi-continuous function is gs_α^{**} -continuous.
- (5) Every semi $^*\alpha$ -continuous function is gs_α^{**} -continuous.
- (6) Every gs_α^{**} -continuous function is gs-continuous.
- (7) Every gs_α^{**} -continuous function is gsp-continuous.

Proof:

- (1) Let V be a closed set in Y . Since, f is continuous, $f^{-1}(V)$ is closed in X . By theorem 3.2 [16], $f^{-1}(V)$ is gs_α^{**} -closed in X and so f is gs_α^{**} -continuous.
- (2)-(7). Similar to the proof of (1).

Remark 3.4: The converses of the above theorems are not be true as seen from the following examples.

Example 3.5: Let $X = Y = \{a, b, c, d\}$, $\tau = \{\emptyset, X, \{a\}, \{d\}, \{a, d\}, \{c, d\}, \{a, c, d\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = b, f(b) = c, f(c) = d, f(d) = a$ is gs_α^{**} -continuous but not continuous.

Example 3.6: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, b\}, \{b\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = c, f(b) = a, f(c) = b$ is gs_α^{**} -continuous but not ω -continuous.

Example 3.7: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{b, c\}, \{a\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = b, f(b) = c, f(c) = a$ is gs_α^{**} -continuous but not g^*s continuous.

Example 3.8: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{b, c\}, \{a\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = c, f(b) = a, f(c) = b$ is gs_α^{**} -continuous but not semi-continuous.

Example 3.9: Let $X = Y = \{a, b, c, d\}$, $\tau = \{\emptyset, X, \{b, c, d\}, \{a, d\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = f(b) = a, f(c) = b, f(d) = c$ is gs_α^{**} -continuous but not semi*-continuous.

Example 3.10: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, b\}, \{a\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{b, c\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = c, f(b) = a, f(c) = b$ is gs -continuous but not gs_α^{**} -continuous.

Example 3.11: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = c, f(b) = a, f(c) = b$ is gsp -continuous but not gs_α^{**} -continuous.

Remark 3.12: gs_α^{**} -continuous and g -continuous functions are independent of each other.

Example 3.13: Let $X = \{a, b, c, d\}$, $Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{b\}, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = b, f(b) = f(c) = c, f(d) = a$ is g -continuous but not gs_α^{**} -continuous.

Example 3.14: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{b\}, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = b, f(b) = c, f(c) = d, f(d) = a$ is gs_α^{**} -continuous but not g -continuous.

Remark 3.15: gs_α^{**} -continuous and semi*- α -continuous functions are independent of each other.

Example 3.16: Let $X = \{a, b, c, d\}$, $Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$ and $\sigma = \{\emptyset, Y, \{b, c\}, \{a\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = f(d) = b, f(b) = a, f(c) = c$ is gs_α^{**} -continuous but not semi*- α -continuous.

Example 3.17: Let $X = \{a, b, c, d\}$, $Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{b\}, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(b) = f(c) = a, f(a) = c, f(d) = b$ is semi*- α -continuous but not gs_α^{**} -continuous.

4. Characterizations of gs_α^{**} -continuous functions

Theorem 4.1: The following are equivalent for a function $f: (X, \tau) \rightarrow (Y, \sigma)$. Assume that $gs_\alpha^{**}O(X, \tau)$ is closed under any union.

- (i) f is gs_α^{**} -continuous.
- (ii) For each $x \in X$ and each open set F in Y containing $f(x)$, there exists a gs_α^{**} -open set U in X containing x such that $f(U) \subseteq F$.

Proof:

(i) \Rightarrow (ii): Let $x \in X$ and F be an open set in Y containing $f(x)$. Since f is gs_α^{**} -continuous, $f^{-1}(F)$ is gs_α^{**} -open in X containing x . Take $U = f^{-1}(F)$ then U is a gs_α^{**} -open set in X containing x such that $f(U) \subseteq F$.

(ii) \Rightarrow (i): Let F be an open set in Y such that $x \in f^{-1}(F)$. Then F is an open set containing $f(x)$. By (ii), there exists a gs_α^{**} -open set U_x in X containing x such that $f(U_x) \subseteq F$ which implies $U_x \subseteq f^{-1}(F)$. Therefore $f^{-1}(F) = \cup \{U_x : x \in f^{-1}(F)\}$. Since U_x is gs_α^{**} -open and $gs_\alpha^{**}O(X, \tau)$ is closed under any union. Hence $f^{-1}(F)$ is open and so f is gs_α^{**} -continuous.

Theorem 4.2: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is gs_α^{**} -continuous if and only if $f^{-1}(V)$ is gs_α^{**} -open in X for every open set V in Y .

Proof: Since $f^{-1}(V^c) = (f^{-1}(V))^c$, proof follows.

Remark 4.3: The composition of two gs_{α}^{**} -continuous functions is not gs_{α}^{**} -continuous.

Example 4.4: Let $X = Y = Z = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{a, b\}\}$, $\sigma = \{\emptyset, Y, \{a, b\}\}$ and $\mu = \{\emptyset, Z, \{a\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = a, f(b) = c, f(c) = b$ and $g: (Y, \sigma) \rightarrow (Z, \mu)$ defined by $g(a) = b, g(b) = a, g(c) = c$. Then f and g are gs_{α}^{**} -continuous but $g \circ f: (X, \tau) \rightarrow (Z, \mu)$ is not gs_{α}^{**} -continuous.

Theorem 4.5: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ and $g: (Y, \sigma) \rightarrow (Z, \mu)$ be any functions. Then

- (i) $g \circ f: (X, \tau) \rightarrow (Z, \mu)$ is gs_{α}^{**} -continuous if g is continuous and f is gs_{α}^{**} -continuous.
- (ii) $g \circ f: (X, \tau) \rightarrow (Z, \mu)$ is gsp-continuous if g is continuous and f is gs_{α}^{**} -continuous.

Proof:

- (i) Let V be any closed set in Z . Since g is continuous, $g^{-1}(V)$ is closed in Y . Since f is gs_{α}^{**} -continuous, $f^{-1}(g^{-1}(V)) = (g \circ f)^{-1}(V)$ is gs_{α}^{**} -closed set in X . Hence $g \circ f$ is gs_{α}^{**} -continuous.
- (ii) Similar to the proof of (i).

Theorem 4.6: Let X and Z be any topological spaces and Y be a ${}_{\alpha}T_{s^{**}}$ -space then the following hold.

- (i) $g \circ f: (X, \tau) \rightarrow (Z, \mu)$ is gs_{α}^{**} -continuous if g is gs_{α}^{**} -continuous and f is gs_{α}^{**} -continuous.
- (ii) $g \circ f: (X, \tau) \rightarrow (Z, \mu)$ is semi-continuous if g is gs_{α}^{**} -continuous and f is semicontinuous.
- (iii) $g \circ f: (X, \tau) \rightarrow (Z, \mu)$ is g^*s -continuous if g is gs_{α}^{**} -continuous and f is g^*s -continuous.

Proof: (i) Let U be any closed set in Z . Since g is gs_{α}^{**} -continuous, $g^{-1}(U)$ is gs_{α}^{**} -closed in Y . But Y is a ${}_{\alpha}T_{s^{**}}$ -space implies $g^{-1}(U)$ is closed in Y . Since f is gs_{α}^{**} -continuous, $f^{-1}(g^{-1}(U)) = (g \circ f)^{-1}(U)$ is gs_{α}^{**} -closed in X and hence $g \circ f$ is gs_{α}^{**} -continuous.

(ii)-(iii) similar to the proof of (i).

Theorem 4.7: If a function $f: X \rightarrow Y$ is gs_{α}^{**} -continuous where X is a ${}_{\alpha}T_{s^{**}}$ -space then f is continuous.(resp.semi-continuous)

Proof: Let V be a closed set in Y . Since f is gs_{α}^{**} -continuous, $f^{-1}(V)$ is gs_{α}^{**} -closed in X . Since X is a ${}_{\alpha}T_{s^{**}}$ -space, $f^{-1}(V)$ is closed in X and so f is continuous.

Theorem 4.8: If a function $f: X \rightarrow Y$ is gs_{α}^{**} -continuous where X is a ${}_{\alpha}T_{s^{**}}$ -space then f is gs -continuous.

Proof: Let V be a closed set in Y . Since f is gs_{α}^{**} -continuous, $f^{-1}(V)$ is gs_{α}^{**} -closed in X . Since X is a ${}_{\alpha}T_{s^{**}}$ -space, $f^{-1}(V)$ is closed in X By theorem 3.2[16], $f^{-1}(V)$ is gs -closed in X and so f is gs -continuous.

Definition 4.9: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called a gs_{α}^{**} -irresolute if $f^{-1}(V)$ is gs_{α}^{**} -closed set in (X, τ) for every gs_{α}^{**} -closed set V in (Y, σ) .

Example 4.10: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{b, c\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = b, f(b) = a, f(c) = c$ is gs_{α}^{**} -irresolute.

Theorem 4.11:

- (1) Every gs_{α}^{**} -irresolute function is gs_{α}^{**} -continuous.
- (2) Every gs_{α}^{**} -irresolute function is gs -continuous.
- (3) Every gs_{α}^{**} -irresolute function is gsp-continuous.

Proof:

(1) Let V be a closed set in Y . By theorem 3.2[16], V is gs_{α}^{**} -closed in Y . Since f is gs_{α}^{**} -irresolute, $f^{-1}(V)$ is gs_{α}^{**} -closed set in X and so f is gs_{α}^{**} -continuous.

(2)-(3) similar to the proof of (1).

Remark 4.12: The converses of the above theorems are not true as seen from the following example.

Example 4.13: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{a, b\}\}$.

Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = b, f(b) = a, f(c) = c$, is gs_{α}^{**} -continuous but not gs_{α}^{**} -irresolute.

Example 4.14: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{b, c\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{b, c\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = a, f(b) = c, f(c) = b$, is gs -continuous but not gs_{α}^{**} -irresolute.

Example 4.15: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{b, c\}\}$.

Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = a, f(b) = c, f(c) = b$, is gsp-continuous but not gs_{α}^{**} -irresolute.

Theorem 4.16: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ and $g: (Y, \sigma) \rightarrow (Z, \mu)$ be any functions. Then the following holds.

- (i) $g \circ f: (X, \tau) \rightarrow (Z, \mu)$ is gs_α^{**} -irresolute if g is gs_α^{**} -irresolute and f is gs_α^{**} -irresolute.
- (ii) $g \circ f: (X, \tau) \rightarrow (Z, \mu)$ is gs_α^{**} -continuous if g is gs_α^{**} -continuous and f is gs_α^{**} -irresolute.

Proof:

- (i) Let V be gs_α^{**} -irresolute in Z . Then $g^{-1}(V)$ is gs_α^{**} -closed in Y . Also f is gs_α^{**} -irresolute, $f^{-1}(g^{-1}(V)) = (g \circ f)^{-1}(V)$ is gs_α^{**} -closed set in X . Hence $g \circ f$ is gs_α^{**} -irresolute.
- (ii) Similar to the proof of (i).

Theorem 4.17: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is gs_α^{**} -irresolute if and only if $f^{-1}(V)$ is gs_α^{**} -open in X for every gs_α^{**} -open set V in Y .

Proof: Since $f^{-1}(V^c) = (f^{-1}(V))^c$, the proof follows.

Theorem 4.18: If a function $f: X \rightarrow Y$ is gs_α^{**} -continuous where X is a ${}_aT_{s^{**}}$ -space then f is gs_α^{**} -irresolute.

Proof: Let U be a gs_α^{**} -closed set in Y . Since Y is a ${}_aT_{s^{**}}$ -space, then U is closed in Y . By theorem 3.2 [16], U is gs_α^{**} -closed set in Y . Since f is gs_α^{**} -irresolute, $f^{-1}(U)$ is gs_α^{**} -closed in X and so f is gs_α^{**} -irresolute.

Theorem 4.19: Let X and Z be any topological spaces and Y be a ${}_aT_{s^{**}}$ -space then $g \circ f: (X, \tau) \rightarrow (Z, \mu)$ is gs_α^{**} -continuous if g is gs_α^{**} -irresolute and f is gs_α^{**} -continuous.

Proof: Let U be any closed set in Z . Since g is gs_α^{**} -irresolute, $g^{-1}(U)$ is gs_α^{**} -closed in Y . But X is a ${}_aT_{s^{**}}$ -space which implies $g^{-1}(U)$ is closed in Y . Since f is gs_α^{**} -continuous, $f^{-1}(g^{-1}(U)) = (g \circ f)^{-1}(U)$ is gs_α^{**} -closed in X and hence $g \circ f$ is gs_α^{**} -continuous.

Theorem 4.20: Let X and Z be any topological spaces and Y be a ${}^aT_{s^{**}}$ -space then $g \circ f: (X, \tau) \rightarrow (Z, \mu)$ is gs_α^{**} -continuous if g is gs -continuous and f is gs_α^{**} -irresolute.

Proof: Let U be any closed set in Z . Since g is gs -continuous, $g^{-1}(U)$ is gs -closed in Y . But Y is a ${}^aT_{s^{**}}$ -space implies $g^{-1}(U)$ is gs_α^{**} -closed in Y . Since f is gs_α^{**} -irresolute, $f^{-1}(g^{-1}(U)) = (g \circ f)^{-1}(U)$ is gs_α^{**} -closed in X . Consequently $g \circ f$ is gs_α^{**} -continuous.

5. Contra gs_α^{**} -continuous functions

In this section, we define contra gs_α^{**} -continuous functions and derives some of their properties.

Definition 5.1: A function $f: X \rightarrow Y$ is said to be contra gs_α^{**} -continuous if $f^{-1}(V)$ is gs_α^{**} -closed in X for every open set V in Y .

Example 5.2: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b, c\}\}$ and $\sigma = \{\emptyset, Y, \{a\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = c, f(b) = a, f(c) = b$ is a contra gs_α^{**} -continuous.

Theorem 5.3: The following are equivalent for a function $f: (X, \tau) \rightarrow (Y, \sigma)$.

Assume that $gs_\alpha^{**}O(X, \tau)$ is closed under any union.

- (1) f is contra gs_α^{**} -continuous.
- (2) For every closed set F of Y , $f^{-1}(F)$ is gs_α^{**} -open in X .
- (3) For each $x \in X$ and each closed set F of Y containing $f(x)$, there exists gs_α^{**} -open set U containing x in X such that $f(U) \subset F$.

Proof:

(1) \Rightarrow (2): Let F be a closed set in Y . Then $Y - F$ is an open set in Y . By (1), $f^{-1}(Y - F) = X - f^{-1}(F)$ is gs_α^{**} -closed in X . which implies $f^{-1}(F)$ is gs_α^{**} -open in X .

(2) \Rightarrow (1): Similar to the proof of (1).

(2) \Rightarrow (3): Let F be a closed set in Y containing $f(x)$. Then $x \in f^{-1}(F)$. By (2), $f^{-1}(F)$ is gs_α^{**} -open in X containing x .

Let $U = f^{-1}(F)$. Then U is gs_α^{**} -open in X containing x and $f(U) = f(f^{-1}(F)) \subset F$.

(3) \Rightarrow (2): Let F be a closed set in Y containing $f(x)$ which implies $x \in f^{-1}(F)$. From (3), there exists gs_α^{**} -open set U_x in X containing x such that $f(U_x) \subset F$ which implies $U_x \subset f^{-1}(F)$. Therefore $f^{-1}(F) = \cup \{U_x : x \in f^{-1}(F)\}$, Since U_x is gs_α^{**} -open and $gs_\alpha^{**}O(X, \tau)$ is closed under any union, $f^{-1}(F)$ is gs_α^{**} -open in X .

Remark 5.4: Composition of two contra gs_α^{**} -continuous function is not contra gs_α^{**} -continuous.

Example 5.5: $X = \{a, b, c, d\}$, $Y = Z = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, b\}, \{a, b, c\}, \{a, b, d\}\}$, and $\sigma = \{\emptyset, Y, \{a\}, \{a, b\}\}$ and $\mu = \{\emptyset, Z, \{a\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = f(b) = c$, $f(c) = b$, $f(d) = a$ and $g: (Y, \sigma) \rightarrow (Z, \mu)$ defined by $g(a) = b$, $g(b) = c$, $g(c) = a$ are gs_α^{**} -continuous but $g \circ f: (X, \tau) \rightarrow (Z, \mu)$ is not gs_α^{**} -continuous.

Theorem 5.6:

- (i) Every contra-continuous function is contra gs_α^{**} -continuous.
- (ii) Every contra semi-continuous function is contra gs_α^{**} -continuous.
- (iii) Every contra semi*-continuous function is contra gs_α^{**} -continuous.
- (iv) Every contra gs_α^{**} -continuous function is contra gs -continuous.
- (v) Every contra gs_α^{**} -continuous function is contra gsp -continuous.

Proof:

(i) Let V be any open set in Y . Since f is contra-continuous, $f^{-1}(V)$ is closed in X . By theorem 3.2[16], $f^{-1}(V)$ is gs_α^{**} -closed in X . Hence f is contra gs_α^{**} -irresolute.

(ii) - (v). Similar to the proof of (i).

Remark 5.7: The converses of the above theorems are not true as seen from the following examples.

Example 5.8: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{a\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = c$, $f(b) = a$, $f(c) = b$ is contra gs_α^{**} -continuous but not contra-continuous.


Example 5.9: Let $X = \{a, b, c, d\}$, $Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, b\}, \{a, b, c\}, \{a, b, d\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = f(d) = b$, $f(c) = a$, $f(b) = c$ is contra gs_α^{**} -continuous but not contra semi-continuous.

Example 5.10: Let $X = \{a, b, c, d\}$, $Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{c\}, \{a, d\}, \{a, c, d\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = b$, $f(b) = a$, $f(c) = d$, $f(d) = c$ is contra gs_α^{**} -continuous but not contra semi*-continuous.

Example 5.11: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = b$, $f(b) = c$, $f(c) = a$ is contra gs -continuous but not contra gs_α^{**} -continuous.

Example 5.12: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = c$, $f(b) = a$, $f(c) = b$ is contra gsp -continuous but not contra gs_α^{**} -continuous.

Remark 5.13: From the above results we have the following diagram.



In the above diagram $A \rightarrow B$ denotes A implies B but not conversely.

Remark 5.14: Contra g -continuous function and contra gs_α^{**} -continuous functions are independent of each other.

Example 5.15: Let $X = \{a, b, c, d\}$, $Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{d\}, \{a, d\}, \{c, d\}, \{a, c, d\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = f(b) = a$, $f(c) = d$, $f(d) = b$ is contra g -continuous but not contra gs_α^{**} -continuous.

Example 5.16: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{b, c\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = c$, $f(b) = a$, $f(c) = b$ is contra gs_α^{**} -continuous but not contra g -continuous.

Remark 5.17: Contra gs_α^{**} -continuous function and contra semi α -continuous functions are independent of each other.

Example 5.18: Let $X = \{a, b, c, d\}$, $Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{b, c\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = f(d) = b$, $f(b) = a$, $f(c) = c$ is contra gs_α^{**} -continuous but not contra semi α -continuous.

Example 5.19: Let $X = \{a, b, c, d\}$, $Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{b\}, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = f(c) = a$, $f(b) = c$, $f(d) = b$ is contra semi α -continuous but not contra gs_α^{**} -continuous.

Theorem 5.20:

- (i) If $f: X \rightarrow Y$ is gs_α^{**} -continuous and $h: Y \rightarrow Z$ is contra-continuous then $h \circ f: X \rightarrow Z$ is contra gs_α^{**} -continuous.
- (ii) If $f: X \rightarrow Y$ is contra gs_α^{**} -continuous and $h: Y \rightarrow Z$ is continuous then $h \circ f: X \rightarrow Z$ is contra gs_α^{**} -continuous.
- (iii) If $f: X \rightarrow Y$ is contra gs_α^{**} -continuous and $h: Y \rightarrow Z$ is contra-continuous then $h \circ f: X \rightarrow Z$ is gs_α^{**} -continuous.

Proof:

(i) Let V be an open set in Z . Since h is contra-continuous, $h^{-1}(V)$ is closed in Y . Since f is gs_α^{**} -continuous, $f^{-1}(h^{-1}(V)) = (h \circ f)^{-1}(V)$ is gs_α^{**} -closed in X and hence $h \circ f$ is gs_α^{**} -continuous.

(ii) - (iii) Similar to the proof of (i).

Remark 5.21: The concept of gs_α^{**} -continuity and contra gs_α^{**} -continuity are independent.

Example 5.22: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, b\}, \{a\}\}$ and $\sigma = \{\emptyset, Y, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = c$, $f(b) = a$, $f(c) = b$ is contra gs_α^{**} -continuous but not gs_α^{**} -continuous.

Example 5.23: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, b\}\}$ and $\sigma = \{\emptyset, Y, \{a\}, \{a, b\}\}$. Then $f: (X, \tau) \rightarrow (Y, \sigma)$ defined by $f(a) = b$, $f(b) = a$, $f(c) = c$ is gs_α^{**} -continuous but not contra gs_α^{**} -continuous.

Theorem 5.24: If $f: (X, \tau) \rightarrow (Y, \sigma)$ is gs_α^{**} -irresolute and $g: (Y, \sigma) \rightarrow (Z, \mu)$ is a contra gs_α^{**} -continuous function then $g \circ f: X \rightarrow Z$ is contra gs_α^{**} -continuous.

Proof: Let V be an open set in Z . Since g is contra gs_α^{**} -continuous, $g^{-1}(V)$ is gs_α -closed in Y . Since f is contra gs_α^{**} -irresolute, $f^{-1}(g^{-1}(V)) = (g \circ f)^{-1}(V)$ is gs_α^{**} -closed in X and hence $g \circ f$ is contra gs_α^{**} -continuous.

Theorem 5.25: If a function $f: X \rightarrow Y$ is contra gs_α^{**} -continuous and Y is regular, then f is gs_α^{**} -continuous.

Proof: Let $x \in X$ and V be an open set in Y containing $f(x)$. Since Y is regular there exists an open set W in Y containing $f(x)$ such that $cl(W) \subset V$. Since f is contra gs_α^{**} -continuous. By theorem 4.1, there exists gs_α^{**} -open set U in X containing x such that $f(U) \subset cl(W)$. Then $f(U) \subset cl(W) \subset V$. Therefore f is gs_α^{**} -continuous.

Theorem 5.26: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a function and X is a ${}_aT_{s^{**}}$ -space. Then the following are equivalent.

- (i) f is contra semi-continuous.
- (ii) f is contra gs_α^{**} -continuous.

Proof:

(i) \Rightarrow (ii): By theorem 5.6, proof follows.

(ii) \Rightarrow (i): Let V be any open set in Y . Since f is contra gs_α^{**} -continuous, $f^{-1}(V)$ is gs_α^{**} -closed in X . Since X is ${}_aT_{s^{**}}$ -space, $f^{-1}(V)$ is closed in X and hence $f^{-1}(V)$ is semi-closed in X f is contra semi-continuous.

Theorem 5.27: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a function and X is a ${}^aT_{s^{**}}$ -space. Then the following are equivalent.

- (i) f is contra gs_α^{**} -continuous.
- (ii) f is contra gs -continuous.

Proof: Similar to the proof of theorem 5.26.

Theorem 5.28: If f is gs_α^{**} -continuous and if Y is locally indiscrete then f is contra gs_α^{**} -continuous.

Proof: Let V be an open set in Y . Since Y is locally indiscrete, V is closed in X . Since f is gs_α^{**} -continuous, $f^{-1}(V)$ is gs_α^{**} -closed in X hence f is contra gs_α^{**} -continuous.

Theorem 5.29: If a function $f: (X, \tau) \rightarrow (Y, \sigma)$ is continuous and X is locally indiscrete then f is contra gs_{α}^{**} -continuous.

Proof: Let V be an open set in (Y, σ) . Since f is continuous, $f^{-1}(V)$ is open in X . Since X is locally indiscrete, $f^{-1}(V)$ is closed set in X . By theorem 3.2[16], $f^{-1}(V)$ is gs_{α}^{**} -closed in X and hence f is contra gs_{α}^{**} -continuous.

Theorem 5.30: If a function $f: (X, \tau) \rightarrow (Y, \sigma)$ is contra gs_{α}^{**} -continuous and X is a ${}_aT_{s^{**}}$ - space then $f: (X, \tau) \rightarrow (Y, \sigma)$ is contra gs -continuous.

Proof: Let V be an open set in Y . Since f is contra gs_{α}^{**} -continuous, $f^{-1}(V)$ is gs_{α}^{**} -closed in X . Since X is ${}_aT_{s^{**}}$ -space, $f^{-1}(V)$ is closed in X and so gs -closed in X and hence f is contra gs_{α}^{**} -continuous.

Definition 5.31: A space X is called locally gs_{α}^{**} -indiscrete if every gs_{α}^{**} -open set is closed in X .

Theorem 5.32: If a function $f: (X, \tau) \rightarrow (Y, \sigma)$ is gs_{α}^{**} -continuous and the space X is locally gs_{α}^{**} -indiscrete then f is contra continuous.

Proof: Let V be an open set in Y . Since f is gs_{α}^{**} -continuous, $f^{-1}(V)$ is gs_{α}^{**} -open in X . Since X is locally gs_{α}^{**} -indiscrete, $f^{-1}(V)$ is closed in X and by theorem 3.2[16], $f^{-1}(V)$ is gs_{α}^{**} -closed in X . Consequently f is contra gs_{α}^{**} -continuous.

Theorem 5.33: If a function $f: (X, \tau) \rightarrow (Y, \sigma)$ is gs_{α}^{**} -irresolute where Y is a locally gs_{α}^{**} -indiscrete space and $g: (Y, \sigma) \rightarrow (Z, \mu)$ is contra gs_{α}^{**} -continuous function then $g \circ f$ is gs_{α}^{**} -continuous.

Proof: Let V be any closed set in Z . Since g is contra gs_{α}^{**} -continuous, $g^{-1}(V)$ is gs_{α}^{**} -open in Y . But Y is locally gs_{α}^{**} -indiscrete implies $g^{-1}(V)$ is closed in Y . By theorem 3.2[16], $g^{-1}(V)$ is gs_{α}^{**} -closed in Y . Since f is gs_{α}^{**} -irresolute, $f^{-1}(g^{-1}(V)) = (g \circ f)^{-1}(V)$ is gs_{α}^{**} -closed in X and hence $g \circ f$ is gs_{α}^{**} -continuous.

Theorem 5.34: If a function $f: (X, \tau) \rightarrow (Y, \sigma)$ is gs_{α}^{**} -continuous and the space (X, τ) is locally gs_{α}^{**} -indiscrete space then f is contra gs_{α}^{**} -continuous.

Proof: Let V be any open set in (Y, σ) . Since f is gs_{α}^{**} -continuous, $f^{-1}(V)$ is gs_{α}^{**} -open in X . Since X is locally gs_{α}^{**} -indiscrete, $f^{-1}(V)$ is closed in X . By theorem 3.2 [16], $f^{-1}(V)$ is gs_{α}^{**} -closed set in X and hence f is contra gs_{α}^{**} -continuous.

6. Contra gs_{α}^{**} -closed graph

Definition 6.1: The graph $G(f)$ of a function $f: X \rightarrow Y$ is said to be gs_{α}^{**} -closed (resp.contra gs_{α}^{**} -closed) if for each $(x, y) \in (X \times Y) - G(f)$, there exist an $U \in gs_{\alpha}^{**}O(X, x)$ and an open (resp.closed) set V in Y such that $(U \times V) \cap G(f) = \emptyset$.

Lemma 6.2: A function $f: X \rightarrow Y$ is gs_{α}^{**} -closed (resp.contra gs_{α}^{**} -closed) if for each $(x, y) \in (X \times Y) - G(f)$ there exists $U \in gs_{\alpha}^{**}O(X, x)$ and an open set (resp.closed set) V in Y containing y such that $f(U) \cap V = \emptyset$.

Proof: We shall prove that $f(U) \cap V = \emptyset$ iff $(U \times V) \cap G(f) = \emptyset$. Let $(U \times V) \cap G(f) \neq \emptyset$. Then there exists $(x, y) \in (U \times V)$ and $(x, y) \in G(f)$ which implies $x \in U, y \in V$ and $y = f(x) \in V$. Therefore $f(U) \cap V \neq \emptyset$.

Theorem 6.3: If a function $f: X \rightarrow Y$ is gs_{α}^{**} -continuous and Y is a T_1 -space then $G(f)$ is contra gs_{α}^{**} -closed in $X \times Y$.

Proof: Let $(x, y) \in (X \times Y) - G(f)$. Then $y \neq f(x)$. Since Y is T_1 , there exists an open set V of Y such that $f(x) \in V, y \notin V$. Since f is gs_{α}^{**} -continuous, by theorem 4.1 there exists a gs_{α}^{**} -open set U of X containing x such that $f(U) \subset V$. Therefore $f(U) \cap (Y - V) = \emptyset$ where $Y - V$ is closed in Y containing y . By lemma 6.2, $G(f)$ is a gs_{α}^{**} -closed graph in $X \times Y$.

Theorem 6.4: Let $f: X \rightarrow Y$ be a function and $g: X \times Y$ be the graph of f defined by $g(x) = (x, f(x))$ for every $x \in X$. If g is contra gs_{α}^{**} -continuous, then f is contra gs_{α}^{**} - Continuous.

Proof: Let U be an open set in Y , then $X \times U$ is an open set in $X \times Y$. Since g is contra gs_{α}^{**} -continuous, $f^{-1}(U) = g^{-1}(X \times U)$ is gs_{α}^{**} -closed in X . Thus f is contra gs_{α}^{**} -continuous.

Definition 6.5:

- (i) $gs_{\alpha}^{**} - T_0$ if for every pair of distinct points x, y in X there exists a gs_{α}^{**} -open set U containing one of the points but not the other.
- (ii) $gs_{\alpha}^{**} - T_1$ if for every pair of distinct points x, y in X there exists a gs_{α}^{**} -open set U containing x not y and a gs_{α}^{**} -open set V containing y but not x .
- (iii) $gs_{\alpha}^{**} - T_2$ if for every pair of distinct points x, y in X there exists disjoint gs_{α}^{**} -open sets U and V containing x and y respectively.

Theorem 6.6: If $f: (X, \tau) \rightarrow (Y, \sigma)$ is an injective function with the gs_{α}^{**} -closed graph $G(f)$ then X is $gs_{\alpha}^{**} - T_1$.

Proof: Let x and y be two distinct points of X , then $f(x) \neq f(y)$. Thus $(x, f(y)) \in X \times Y - G(f)$. Since $G(f)$ is gs_{α}^{**} -closed, there exists a gs_{α}^{**} -open set U containing x and an open set V containing $f(y)$ such that $f(U) \cap V = \emptyset$. By theorem 3.2 [16], U and V are gs_{α}^{**} -open sets containing x and $f(y)$ such that $f(U) \cap V = \emptyset$. Hence $y \notin U$. Similarly there exist gs_{α}^{**} -open sets M and N containing y and $f(x)$ such that $f(M) \cap N = \emptyset$. Hence $x \notin M$. It follows that X is $gs_{\alpha}^{**} - T_1$.

Theorem: 6.7: If $f: (X, \tau) \rightarrow (Y, \sigma)$ is a surjective function with the gs_{α}^{**} -closed graph $G(f)$ then Y is $gs_{\alpha}^{**} - T_1$.

Proof: Let y and z be two distinct points of Y . Since f is surjective there exist a point x in X such that $f(x) = z$. Therefore $(x, y) \notin G(f)$, by lemma 6.2, there exists a gs_{α}^{**} -open set U containing x and an open set V containing y such that $f(U) \cap V = \emptyset$. By theorem 3.2[16], U and V are gs_{α}^{**} -open sets containing x and y such that $f(U) \cap V = \emptyset$. It follows that $z \notin V$. Similarly there exist $w \in X$ such that $f(w) = y$. Hence $(w, z) \notin G(f)$. Similarly there exist gs_{α}^{**} -open sets M and N containing w and z respectively such that $f(M) \cap N = \emptyset$. Thus $y \notin N$. Hence the space Y is $gs_{\alpha}^{**} - T_1$.

REFERENCES

1. Alli K. *Contra $g^{\#}p$ -continuous functions*, International Journal of Mathematics Trends and Technology- volume 4 Issue 11-Dec 2013.
2. Balachandran K. Sundaram P. and Maki H. *On Generalised Continuous Maps in topological spaces*, Memoirs of the Faculty of Science Kochi University Series A, vol. 12, pp. 513, 1991.
3. Bhattacharya P. and Lahiri B.K. *Semi-generalized closed sets in Topology*, Indian J. Math., 29(1987), 375-382.
4. Caldas M. Jafari S. Noiri T. Simeos, M. *A new generalization of contra-continuity via Levines g -closed sets*, Chaos Solitons Fractals 42 (2007), 1595-1603.
5. Devi R. Maki H. and Balachandran K. *semi generalized closed maps and generalized semi closed maps*, Memoirs of the Faculty of Science Kochi University Series A, vol. 14, pp. 4154, 1993.
6. Dontchev J. Noiri T. *Contra-semi continuous functions*, Math. Pannon, 10 (1999), 159- 168.
7. Dontchev J. *Contra-continuous functions and strongly S -closed spaces*, Internat. J.Math. Math. sci. 19 (1996), 303-310.
8. Dontchev J. *On generalizing semi-preopen sets*, Mem. Fac. Sci. Kochi Univ. Ser.A. Math., 16(1995), 35-48.
9. Dunham W., *A New closure operator for Non- T_1 Topologies*, Kyungpook Math. J. 22 (1982), 55-60.
10. Levine, N., *Generalized closed sets in topology*, Rend. Circ. Mat. Palermo., 19(2), (1970), 89-96.
11. Levine N., *Semi-open sets and Semi-continuity in topological spaces*, Amer. Math. Monthly, 70(1), (1963), 36-41.
12. Long P.E. *Functions with closed graphs*, Amer. Math. Monthly, 76(1969), 930-932.
13. Othman H.A., *New types of α -continuous mapping Master Degree Thesis*, Al- Muatansiriya University, IRAQ (2004).
14. Pushpalatha A. *On g^*s closed sets in Topological spaces*, Int.J.Contemp Math. Sciences, vol. 6 19, pp 917-929, 2011.
15. Robert A. and Pious Missier, S. *Semi-star-Alpha-Open Sets and Associated Functions*, International Journal of Computer Applications, vol.104, 16, 2014.
16. Santhini C., and Lakshmi Priya S., *New class of generalized closed sets in topological spaces*, Proceedings of International Conference on Recent trends in Applied Mathematics, S.R.N.M college, Sattur, 2017.
17. Sundaram P., and Sherik John M., *On ω -closed sets in topology*, Acta ciencia Indica, 4(2000), 43-65.
18. Willard S., *General Topology*, Addition Wesley (1970).

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2017. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]