

## On $rgw\alpha$ - Open Sets in Topological Spaces

Dr. R. S. WALI<sup>1</sup>, VIJAYLAXMI R. PATIL\*<sup>2</sup>

<sup>1</sup>Department of Mathematics,  
Bhandari and Rathi College, Guledagudd, Karnataka, INDIA.

<sup>2</sup>Department of Mathematics,  
Rani Channamma University Belagavi, Karnataka, INDIA.

(Received On: 16-03-17; Revised & Accepted On: 17-04-17)

---

### ABSTRACT

In this paper, we introduced and studied  $rgw\alpha$ -open sets in topological space and obtain some of their properties. Also we introduce  $rgw\alpha$ -interior,  $rgw\alpha$ -closure,  $rgw\alpha$ -neighbourhood and  $rgw\alpha$ -limit points in topological spaces.

**Keywords:**  $rgw\alpha$ -open sets,  $rgw\alpha$ -interior,  $rgw\alpha$ -closure,  $rgw\alpha$ -neighbourhood,  $rgw\alpha$ -limit points.

**Mathematics Subject Classification (2010):** 54A05.

---

### 1] INTRODUCTION

Regular open sets have been introduced and investigated by Stone [6]. P.Sundaram and M.Sheik John [8] defined and studied  $w$ -closed sets in topological spaces. S.S Benchalli and R.S.Wali [12] introduced and studied  $rw$ -closed sets. N.Jasted [7] introduced and studied  $\alpha$ -sets. S.S.Benchalli *et al.* [11] studied  $w\alpha$ -closed sets in topological spaces. S.S.Benchalli *et al.* [10] introduced  $g\alpha$ -closed sets. and P.G.Patil *et al.* [9] introduced  $g^*w\alpha$ -closed set. A. Vadivel and Vairamanickam [2] introduced  $rg\alpha$ -closed sets and  $rg\alpha$ -open sets in topological spaces. In this paper we define  $rgw\alpha$ -open sets, its properties and  $rgw\alpha$ -interior,  $rgw\alpha$ -closure,  $rgw\alpha$ -neighbourhood and  $rgw\alpha$ -limit points and obtain some of its basic properties.

### 2] PRELIMINARIES

Throughout the paper  $X$  and  $Y$  denote the topological space  $(X, \tau)$  and  $(Y, \tau)$  respectively. And on which no separation axioms are assumed unless otherwise explicitly stated. For a subset  $A$  of space  $X$ ,  $cl(A)$ ,  $int(A)$ ,  $A^c$ , and  $rcl(A)$  denote the closure of  $A$ , Interior of  $A$ , complement of  $A$  and regular closure of  $A$  in  $X$  respectively.

**Definition 2.1:** A subset  $A$  of a space  $X$  is called

- 1) a regular open set [6] if  $A = int(cl(A))$  and a regular closed set if  $A = cl(int(A))$ .
- 2) a  $\alpha$ -open set [7] if  $A \subseteq int(cl(int(A)))$  and  $\alpha$ -closed set if  $cl(int(cl(A))) \subseteq A$ .
- 3) a weakly closed set (briefly,  $w$ -closed) [1] if  $cl(A) \subseteq U$  whenever  $A \subseteq U$  &  $U$  is semi open in  $X$ .
- 4) a weakly  $\alpha$ -closed set (briefly,  $w\alpha$ -closed) [11] if  $\alpha cl(A) \subseteq U$  whenever  $A \subseteq U$  &  $U$  is  $w$ -open in  $X$ .
- 5) a regular  $\alpha$ -open set (2) if there is a regular open set  $U \ni U \subseteq A \subseteq \alpha cl(U)$

The intersection of all regular closed (resp.  $\alpha$ -closed,  $w\alpha$ -closed and regular  $\alpha$ -closed) subsets of space  $X$  containing  $A$  is called regular closure (Resp.  $\alpha$ -closure,  $w\alpha$ -closure and regular  $\alpha$ -closure) of  $A$  and denoted by  $rcl(A)$  (resp.  $\alpha cl(A)$ ,  $w\alpha cl(A)$  and  $racl(A)$ ).

---

**Corresponding Author: Vijaylaxmi R. Patil\*<sup>2</sup>**

**<sup>2</sup>Department of Mathematics, Rani Channamma University Belagavi, Karnataka, INDIA.**

**Definition 2.2:** A subset  $A$  of a space  $X$  is called

- 1) generalized  $\alpha$ -closed set (briefly  $g\alpha$ -closed) [4], if  $\text{acl}(A) \subseteq U$  whenever  $A \subseteq U$  and  $U$  is  $\alpha$ -open in  $X$ .
- 2) generalized semi-pre closed set (briefly  $gsp$ -closed) [5] if  $\text{spcl}(A) \subseteq U$  whenever  $A \subseteq U$  and  $U$  is open in  $X$ .
- 3) generalized weak  $\alpha$ -closed (briefly  $gwa$ -closed) set [10] if  $\text{acl}(A) \subseteq U$  whenever  $A \subseteq U$  &  $U$  is  $w\alpha$ -open in  $X$ .
- 4) generalized star weakly  $\alpha$ -closed set (briefly  $g^*w\alpha$ -closed) [9] if  $\text{cl}(A) \subseteq U$  whenever  $A \subseteq U$  &  $U$  is  $w\alpha$ -open in  $X$ .
- 5) regular generalized  $\alpha$ -closed set (briefly  $rg\alpha$ -closed) [2] if  $\text{acl}(A) \subseteq U$  whenever  $A \subseteq U$  &  $U$  is regular  $\alpha$ -open in  $X$ .

The complements of the above mentioned closed sets are respective open sets.

### 3. $rgw\alpha$ -closed sets in topological spaces.

**Definition 3.1 [13]:** A subset  $A$  of a space  $X$  is called regular generalized weakly  $\alpha$ -closed set (briefly  $rgw\alpha$ -closed) if  $\text{r}\alpha\text{cl}(A) \subseteq U$  whenever  $A \subseteq U$  &  $U$  is weak  $\alpha$ -open set in  $X$ .

**Results 3.2** from [13]:

- 1) Every closed set is  $rgw\alpha$ -closed set in  $X$ .
- 2) Every regular closed set is  $rgw\alpha$ -closed set in  $X$ .
- 3) Every weak- closed set is  $rgw\alpha$ -closed set in  $X$ .
- 4) Every  $\alpha$ -closed,  $g\alpha$ -closed,  $rg\alpha$ -closed,  $gwa$ -closed and  $g^*w\alpha$ -closed sets are  $rgw\alpha$ -closed sets in  $X$ .
- 5) Every  $rw$ -closed,  $r\alpha$ -closed,  $rs$ -closed and  $w\alpha$ -closed sets are  $rgw\alpha$ -closed sets in  $X$ .
- 6) Every  $rgw\alpha$ -closed set is  $g\beta$ -closed set in  $X$ .
- 7) The union of two  $rgw\alpha$ -closed sets of  $X$  is  $rgw\alpha$ -closed set in  $X$ .
- 8) The intersection of two  $rgw\alpha$ -closed sets of  $X$  is need not be  $rgw\alpha$ -closed set.

### 4. $rgw\alpha$ -open sets and their basic properties

In this section we introduce and study  $rgw\alpha$ -open sets in topological spaces and obtain some of their properties.

**Definition 4.1:** A subset  $A$  of  $X$  is called regular generalized weakly- $\alpha$  open set ( $rgw\alpha$ -open set) in  $X$  if  $A^c$  is  $rgw\alpha$ -closed in  $X$ . We denote the family of all  $rgw\alpha$ -open sets in  $X$  by  $RGW\alpha O(X)$ .

**Theorem 4.2:** If a subset  $A$  of a space  $X$  is  $w$ -open then it is  $rgw\alpha$ -open set, but not conversely.

**Proof:** Let  $A$  be a  $w$ -open set in a space  $X$ . Then  $A^c$  is  $w$ -closed set. By result 3.2(3)  $A^c$  is  $rgw\alpha$ -closed. Therefore  $A$  is  $rgw\alpha$ -open set in  $X$ . The converse of this theorem need not be true as seen from the following example.

**Example 4.3:** Let  $X = \{a, b, c, d, e\}$  with topology  $\tau = \{X, \phi, \{a\}, \{d\}, \{e\}, \{a, d\}, \{a, e\}, \{d, e\}, \{a, d, e\}\}$ . Then the set  $A = \{c\}$  is  $rgw\alpha$ -open set but not  $w$ -open set in  $X$ .

**Corollary 4.4:** Every open set is  $rgw\alpha$ -open set but not conversely.

**Proof:** Follows from definition and theorem 4.2.

**Corollary 4.5:** Every regular open set is  $rgw\alpha$ -open set but not conversely.

**Proof:** Follows from definition and theorem 4.2.

**Theorem 4.6:** If a subset  $A$  of a space  $X$  is  $rgw\alpha$ -open, then it is  $g\beta$ -open set in  $X$ .

**Proof:** Let  $A$  be  $rgw\alpha$ -open set in  $X$ . Then  $A^c$  is  $rgw\alpha$ -closed set in  $X$ . By result 3.2(6)  $A^c$  is  $g\beta$ -closed set in  $X$ . Therefore  $A$  is  $g\beta$ -open set in space  $X$ . The converse of this theorem need not be true as seen from the following example.

**Example 4.7:** In example 4.3 the subset  $\{b, c\}$  of  $X$  is  $g\beta$ -open set but not  $rgw\alpha$ -open set.

**Theorem 4.8:** If a subset  $A$  of  $X$  is  $g\alpha$ -open set then it is  $rgw\alpha$ -open set in  $X$ , but not conversely.

**Proof:** Let  $A$  be a  $g\alpha$ -open set in a space  $X$ . Then  $A^c$  is  $g\alpha$ -closed set. By result 3.2(4)  $A^c$  is  $rgw\alpha$ -closed. Therefore  $A$  is  $rgw\alpha$ -open set in  $X$ . The converse of this theorem need not be true as seen from the following example.

**Example 4.9:** In example 4.3 the subset  $A=\{a, b\}$  of  $X$  is  $rgw\alpha$ -open set but not  $g\alpha$ -open set in  $X$ .

**Theorem 4.10:** If a subset  $A$  of  $X$  is  $gwa$ -open set then it is  $rgw\alpha$ -open set in  $X$ , but not conversely.

**Proof:** Let  $A$  be a  $gwa$ -open set in a space  $X$ . Then  $A^c$  is  $gwa$ -closed set. By result 3.2 (4)  $A^c$  is  $rgw\alpha$ -closed. Therefore  $A$  is  $rgw\alpha$ -open set in  $X$ . The converse of this theorem need not be true as seen from the following example.

**Example 4.11:** In example 4.3 the sub set  $A= \{b\}$  of  $X$  is  $rgw\alpha$ -open set but not  $gwa$ -open set in  $X$ .

**Corollary 4.12:** If a subset  $A$  of  $X$  is  $g^*wa$ -open set then it is  $rgw\alpha$ -open set in  $X$ , but not conversely.

**Proof:** it follows from the theorem 4.10 and the implication  $gwa \Rightarrow g^*wa$  set.

**Theorem 4.13:** If  $A$  and  $B$  are  $rgw\alpha$ -open sets in a space  $X$ . Then  $A \cap B$  is also  $rgw\alpha$ -open set in  $X$ .

**Proof:** If  $A$  and  $B$  are  $rgw\alpha$ -open sets in a space  $X$ . Then  $A^c$  and  $B^c$  are  $rgw\alpha$ -closed sets in a space  $X$ . By result 3.2(7).  $A^c \cup B^c$  is also  $rgw\alpha$ -closed set in  $X$ . That is  $A^c \cup B^c = (A \cap B)^c$  is a  $rgw\alpha$ -closed set in  $X$ . Therefore  $A \cap B$  is  $rgw\alpha$ -open set in  $X$ .

**Remark 4.14:** The union of two  $rgw\alpha$ -open sets in  $X$  is generally not a  $rgw\alpha$ -open in  $X$ .

**Example 4.15:** In example 4.3 the sets  $A=\{a,b\}$  and  $B=\{c\}$  are  $rgw\alpha$ -open sets in  $X$ , But  $A \cup B=\{a,b,c\}$  is not  $rgw\alpha$ -open set in  $X$ .

**Theorem 4.16:** If a set  $A$  is  $rgw\alpha$ -open in a space  $X$ , then  $G=X$ , whenever  $G$  is  $wa$ -open and  $\text{int}(A) \cup A^c \subset G$ .

**Proof:** Suppose that  $A$  is  $rgw\alpha$ -open in  $X$ . Let  $G$  be weak  $\alpha$ -open and  $\text{int}(A) \cup A^c \subset G$ . This implies  $G^c \subset (\text{int}(A) \cup A^c)^c = (\text{int}(A))^c \cap A$ . That is  $G^c \subset (\text{int}(A))^c - A^c$ . Thus  $G^c \subset \text{cl}(A)^c - A^c$ , since  $(\text{int}(A))^c = \text{cl}(A^c)$ . Now  $G^c$  is also weak  $\alpha$ -open and  $A^c$  is  $rgw\alpha$ -closed then by theorem it follows that  $G^c = \emptyset$ . Hence  $G=X$ . The converse of this theorem need not be true as seen from the following example.

**Example 4.17:** In Example 4.3  $RGW\alpha O(X) = \{X, \phi, \{a\}, \{b\}, \{c\}, \{d\}, \{e\}, \{a, b\}, \{a, c\}, \{a, d\}, \{a, e\}, \{b, d\}, \{d, e\}, \{c, d\}, \{c, e\}, \{a, b, d\}, \{a, b, e\}, \{a, c, d\}, \{a, c, e\}, \{a, d, e\}, \{b, c, d\}, \{b, d, e\}, \{c, d, e\}, \{a, b, c, d\}, \{a, b, d, e\}, \{a, b, c, e\}, \{a, c, d, e\}, \{b, c, d, e\}\}$ . And  $WaO(X) = \{X, \phi, \{a\}, \{b\}, \{d\}, \{e\}, \{b, d\}, \{a, d\}, \{a, b\}, \{a, c\}, \{a, e\}, \{b, e\}, \{d, e\}, \{a, b, c\}, \{a, b, d\}, \{a, b, e\}, \{b, d, e\}, \{a, d, e\}, \{a, b, d, e\}, \{a, c, d, e\}\}$ . Take  $A = \{a, c, d\}$ . then  $A$  is not  $rgw\alpha$ -open. However  $\text{int}(A) \cup A^c = \{a, d\} \cup \{b, e\} = \{a, b, d, e\}$ . so for some weak  $\alpha$ -open  $G$ , we have  $\text{int}(A) \cup A^c = \{a, b, d, e\} \subset G$  gives  $G=X$ , but  $A$  is not  $rgw\alpha$ -open.

**Theorem 4.18:** A subset  $A$  of  $(X, \tau)$  is  $rgw\alpha$ -open set if and only if  $U \subseteq r\alpha \text{int}(A)$  whenever  $U$  is  $wa$ -closed and  $U \subseteq A$ .

**Proof:** Assume that  $A$  is  $rgw\alpha$ -open in  $X$  and  $U$  is  $wa$ -closed set of  $(X, \tau)$  such that  $U \subseteq A$ . Then  $X-A$  is  $rgw\alpha$ -closed set in  $(X, \tau)$ . Also  $X-A \subseteq X-U$  and  $X-U$  is  $wa$ -open set of  $(X, \tau)$ . This implies that  $r\alpha \text{cl}(X-A) \subseteq X-U$ . But  $r\alpha \text{cl}(X-A) = X - r\alpha \text{int}(A)$ . Thus  $X - r\alpha \text{int}(A) \subseteq X-U$ . So  $U \subseteq r\alpha \text{int}(A)$ .

Conversely, Suppose  $U \subseteq r\text{aint}(A)$  whenever  $U$  is  $w\alpha$ -closed and  $U \subseteq A$ , To prove that  $A$  is  $rgw\alpha$ -open in  $X$ . Let  $G$  be  $w\alpha$ -open set of  $(X, \tau)$  s.t.  $X-A \subseteq G$ . Then  $X-G \subseteq A$ . Now  $X-G$  is  $w\alpha$ -closed set containing  $A$ . So  $X-G \subseteq r\text{aint}(A)$ ,  $X-r\text{aint}(A) \subseteq G$ , But  $r\text{acl}(X-A) = X - r\text{aint}(A)$ . Thus  $r\text{acl}(X-A) \subseteq G$ . i.e  $X-A$  is  $rgw\alpha$ -closed set. Hence  $A$  is  $rgw\alpha$ -open set.

**Theorem 4.19:** If  $A$  is  $w\alpha$ -open and  $rgw\alpha$ -closed set then  $A$  is  $r\alpha$ -closed.

**Proof:** Since  $A \subseteq A$  and  $A$  is  $w\alpha$ -open and  $rgw\alpha$ -closed we have  $r\text{acl}(A) \subseteq A$ . Thus  $r\text{acl}(A) = A$ . Hence  $A$  is  $r\alpha$ -closed set of  $(X, \tau)$ .

**Theorem 4.20:** If  $r\text{aint}(A) \subseteq B \subseteq A$  and  $A$  is  $rgw\alpha$ -open set in  $X$ , then  $B$  is  $rgw\alpha$ -open set in  $X$ .

**Proof:** If  $r\text{aint}(A) \subseteq B \subseteq A$ , then  $X-A \subseteq X-B \subseteq X-r\text{aint}(A) = r\text{acl}(X-A)$ . Since  $(X-A)$  is  $rgw\alpha$ -closed set, then by theorem 3.15 [13]  $X-B$  is also  $rgw\alpha$ -closed set in  $X$ . Therefore  $B$  is  $rgw\alpha$ -open set in  $X$ .

**Theorem 4.21:** If  $A$  is  $rgw\alpha$ -closed set in  $X$ , then  $r\text{acl}(A)-A$  is  $rgw\alpha$ -open set in  $X$ .

**Proof:** Let  $A$  be  $rgw\alpha$ -closed set in  $X$ , Let  $F$  be an  $w\alpha$ -open s.t.  $F \subseteq r\text{acl}(A)-A$ . Since  $A$  is  $rgw\alpha$ -closed, then by theorem 3.12[13]  $r\text{acl}(A)-A$  does not contain any non empty  $w\alpha$ -closed set in  $X$ . Thus  $F = \emptyset$ . Then  $F \subseteq r\text{aint}(r\text{acl}(A)-A)$ . Therefore by theorem 4.18  $r\text{acl}(A)-A$  is  $rgw\alpha$ -open set in  $X$ .

**Theorem 4.22:** If  $A$  and  $B$  be subsets of space  $(X, \tau)$ . If  $B$   $rgw\alpha$ -open and  $r\text{aint}(B) \subseteq A$ , then  $A\alpha B$  is  $rgw\alpha$ -open set in  $X$ .

**Proof:** Let  $B$  is  $rgw\alpha$ -open in  $X$ .  $r\text{aint}(B) \subseteq A$  and  $r\text{aint}(B) \subseteq B$  is always true, then  $r\text{aint}(B) \subseteq A\alpha B$ . also  $r\text{aint}(B) \subseteq A\alpha B \subseteq B$  and  $B$  is  $rgw\alpha$ -open set then by theorem 4.20  $A\alpha B$  is also  $rgw\alpha$ -open set in  $X$ .

## 5. $rgw\alpha$ -Closure and $rgw\alpha$ -Interior

In this section the notation of  $rgw\alpha$ -Closure and  $rgw\alpha$ -Interior is defined and some of its basic properties are studied.

**Definition 5.1:** For a subset  $A$  of  $X$ ,  $rgw\alpha$ -Closure of  $A$  is denoted by  $rgw\alpha\text{cl}(A)$  and defined as  $rgw\alpha\text{cl}(A) = \bigcap \{G: A \subseteq G, G \text{ is } rgw\alpha\text{-closed in } X\}$  or  $\bigcap \{G: A \subseteq G, G \in RGW\alpha C(X)\}$ .

**Theorem 5.2:** If  $A$  and  $B$  are subsets of a space  $X$  then

- i)  $rgw\alpha\text{cl}(X) = X$ ,  $rgw\alpha\text{cl}(\emptyset) = \emptyset$ .
- ii)  $A \subseteq rgw\alpha\text{cl}(A)$ .
- iii) If  $B$  is any  $rgw\alpha$ -closed set containing  $A$ , then  $rgw\alpha\text{cl}(A) \subseteq B$ .
- iv) If  $A \subseteq B$  then  $rgw\alpha\text{cl}(A) \subseteq rgw\alpha\text{cl}(B)$ .
- v)  $rgw\alpha\text{cl}(A) = rgw\alpha\text{cl}(rgw\alpha\text{cl}(A))$ .
- vi)  $rgw\alpha\text{cl}(A \cup B) = rgw\alpha\text{cl}(A) \cup rgw\alpha\text{cl}(B)$ .

**Proof:** i) By definition of  $rgw\alpha$ -Closure,  $X$  is Only  $rgw\alpha$ -closed set containing  $X$ , therefore  $rgw\alpha\text{cl}(X) =$  Intersection of all the  $rgw\alpha$ -closed set containing  $X = \bigcap \{X\} = X$ , therefore  $rgw\alpha\text{cl}(X) = X$ . Again By the Definition of  $rgw\alpha$ -Closure  $rgw\alpha\text{cl}(\emptyset) =$  Intersection of all  $rgw\alpha$ -closed set containing  $\emptyset = \emptyset \cap$  any  $rgw\alpha$ -closed set containing  $\emptyset = \emptyset$ . Therefore  $rgw\alpha\text{cl}(\emptyset) = \emptyset$ .

ii) By definition of  $rgw\alpha$ -Closure of  $A$  it is obvious that  $A \subseteq rgw\alpha\text{cl}(A)$ .

iii) Let B be any  $rgw\alpha$ -closed set containing A, Since  $rgw\alpha cl(A)$  is the intersection of all  $rgw\alpha$ -closed set containing A,  $rgw\alpha cl(A)$  is contained in every  $rgw\alpha$ -closed set containing A. Hence in particular  $rgw\alpha cl(A) \subseteq B$ .

iv) Let A and B be subsets of X, such that  $A \subseteq B$  by definition of  $rgw\alpha$ -Closure,  $rgw\alpha cl(B) = \bigcap \{F: B \subseteq F \in RGW\alpha C(X)\}$ . If  $B \subseteq F \in RGW\alpha C(X)$ , then  $rgw\alpha cl(B) \subseteq F$ . Since  $A \subseteq B$ ,  $A \subseteq B \subseteq F \in RGW\alpha C(X)$ , we have  $rgw\alpha cl(A) \subseteq F$ ,  $rgw\alpha cl(A) \subseteq \bigcap \{F: B \subseteq F \in RGW\alpha C(X)\} = rgw\alpha cl(B)$ . Therefore  $rgw\alpha cl(A) \subseteq rgw\alpha cl(B)$ .

v) Let A be any subset of X by definition of  $rgw\alpha$ -Closure,  $rgw\alpha cl(A) = \bigcap \{F: A \subseteq F \in RGW\alpha C(X)\}$ . Therefore  $A \subseteq F \in RGW\alpha C(X)$  then  $rgw\alpha cl(A) \subseteq F$ , Since F is  $rgw\alpha$ -closed set containing  $rgw\alpha cl(A)$  by (iii)  $rgw\alpha cl(rgw\alpha cl(A)) = \bigcap \{F: A \subseteq F \in RGW\alpha C(X)\} = rgw\alpha cl(A)$ . therefore  $rgw\alpha cl(rgw\alpha cl(A)) = rgw\alpha cl(A)$

vi) Let A and B be subsets of X, clearly  $A \subseteq A \cup B$ ,  $B \subseteq A \cup B$  from (iv)  $rgw\alpha cl(A) \subseteq rgw\alpha cl(A \cup B)$ ,  $rgw\alpha cl(B) \subseteq rgw\alpha cl(A \cup B)$ . Hence  $rgw\alpha cl(A) \cup rgw\alpha cl(B) \subseteq rgw\alpha cl(A \cup B)$ . Now we have to prove  $rgw\alpha cl(A \cup B) \subseteq rgw\alpha cl(A) \cup rgw\alpha cl(B)$ .

Suppose  $x \notin rgw\alpha cl(A) \cup rgw\alpha cl(B)$  then  $\exists$   $rgw\alpha$ -closed set  $A_1$  and  $B_1$  with  $A \subseteq A_1$ ,  $B \subseteq B_1$  &  $x \notin A_1 \cup B_1$ . We have  $A \cup B \subseteq A_1 \cup B_1$  and  $A_1 \cup B_1$  is the  $rgw\alpha$ -closed set such that  $x \notin A_1 \cup B_1$ . Thus  $x \notin rgw\alpha cl(A \cup B)$  hence  $rgw\alpha cl(A \cup B) \subseteq rgw\alpha cl(A) \cup rgw\alpha cl(B)$  (2). From (1) and (2) we have  $rgw\alpha cl(A \cup B) = rgw\alpha cl(A) \cup rgw\alpha cl(B)$ .

**Theorem 5.3:** If  $A \subseteq X$  is  $rgw\alpha$ -closed set then  $rgw\alpha cl(A) = A$ .

**Proof:** Let A be  $rgw\alpha$ -closed subset of X. We know that  $A \subseteq rgw\alpha cl(A)$  - (1). Also  $A \subseteq A$  and A is  $rgw\alpha$ -closed set by theorem 5.2 (iii)  $rgw\alpha cl(A) \subseteq A$  - (2). Hence  $rgw\alpha cl(A) = A$ .

The converse of the above need not be true as seen from the following example.

**Example 5.4:** Let  $X = \{a, b, c, d, e\}$  with topology  $\tau = \{X, \phi, \{a\}, \{d\}, \{e\}, \{a, d\}, \{a, e\}, \{d, e\}, \{a, d, e\}\}$  here  $A = \{a, d\}$  and  $rgw\alpha cl(A) = \{a, d\} = A$  but A is not  $rgw\alpha$ -closed set.

**Theorem 5.5:** If A and B are subsets of Space X then  $rgw\alpha cl(A \cap B) \subseteq rgw\alpha cl(A) \cap rgw\alpha cl(B)$ .

**Proof:** Let A and B be subsets of X, clearly  $A \cap B \subseteq A$ ,  $A \cap B \subseteq B$ , by theorem 5.2 (iv)  $rgw\alpha cl(A \cap B) \subseteq rgw\alpha cl(A)$ ,  $rgw\alpha cl(A \cap B) \subseteq rgw\alpha cl(B)$ , hence  $rgw\alpha cl(A \cap B) \subseteq rgw\alpha cl(A) \cap rgw\alpha cl(B)$ .

**Remark 5.6:** In general  $rgw\alpha cl(A) \cap rgw\alpha cl(B) \not\subseteq rgw\alpha cl(A \cap B)$ .

**Theorem 5.7:** For an  $x \in X$ ,  $x \in rgw\alpha cl(X)$  if and only if  $A \cap V \neq \phi$  for every  $rgw\alpha$ -open set V containing x.

**Proof:** Let  $x \in rgw\alpha cl(A)$ . To prove  $A \cap V \neq \phi$  for every  $rgw\alpha$ -open set V containing x by contradiction. Suppose  $\exists$   $rgw\alpha$ -open set V containing x s.t.  $A \cap V = \phi$  then  $A \subseteq X - V$ ,  $X - V$  is  $rgw\alpha$ -closed set,  $rgw\alpha cl(A) \subseteq X - V$ . This Shows that  $x \notin rgw\alpha cl(A)$  which is contradiction. Hence  $A \cap V \neq \phi$  for every  $rgw\alpha$ -open set V containing x.

Conversely: Let  $A \cap V \neq \phi$  for every  $rgw\alpha$ -open set V containing x. To prove  $x \in rgw\alpha cl(A)$ . We prove the result by contradiction. Suppose  $x \notin rgw\alpha cl(A)$  then there exist a  $rgw\alpha$ -closed subset F containing A s.t.  $x \notin F$ . Then  $x \in X - F$  is  $rgw\alpha$ -open. Also,  $(X - F) \cap A = \phi$  which is contradiction. Hence  $x \in rgw\alpha cl(A)$ .

**Theorem 5.8:** If A is subset of space X, then

- i)  $rgw\alpha cl(A) \subseteq cl(A)$
- ii)  $rgw\alpha cl(A) \subseteq r\alpha cl(A)$

**Proof:** Let  $A$  be subset of space  $X$  by definition of closure  $cl(A) = \bigcap \{F : A \subseteq F \in C(X)\}$  If  $A \subseteq F \in C(X)$  then  $A \subseteq F \in RGW\alpha C(X)$  because every closed set is  $rgw\alpha$ -closed that is  $rgw\alpha cl(A) \subseteq F$ , therefore  $rgw\alpha cl(A) \subseteq \bigcap \{F : A \subseteq F \in C(X)\}$  Hence  $rgw\alpha cl(A) \subseteq cl(A)$ .

ii) let  $A$  be subset of space  $X$  by definition of  $ra$ -closure  $racl(A) = \bigcap \{F : A \subseteq F \in raC(x)\}$ , If  $A \subseteq F \in raC(x)$  then  $A \subseteq F \in rgw\alpha C(x)$  because every  $ra$ -closed set is  $rgw\alpha$ -closed that is  $rgw\alpha cl(A) \subseteq F$  therefore  $rgw\alpha cl(A) \subseteq \bigcap \{F : A \subseteq F \in raC(x)\} = racl(A)$ . Hence  $rgw\alpha cl(A) \subseteq racl(A)$ .

**Remark 5.9:** Containment relation in the above theorem 5.8 may be proper as seen from following example.

**Example 5.10:** Let  $X = \{a, b, c, d, e\}$  with topology  $\tau = \{X, \phi, \{a\}, \{d\}, \{e\}, \{a, d\}, \{a, e\}, \{d, e\}, \{a, d, e\}\}$ ,  $A = \{a, b, d, e\}$ ,  $cl(A) = \{X\}$ ,  $rgw\alpha cl(A) = \{a, b, d, e\}$  &  $racl(A) = \{X\}$ . It follows that  $rgw\alpha cl(A) \subset cl(A)$  and  $rgw\alpha cl(A) \subset racl(A)$ .

**Theorem 5.11:** If  $A$  is subset of space  $X$  then  $gspcl(A) \subseteq rgw\alpha cl(A)$  where  $gspcl(A) = \bigcap \{F : A \subseteq F \in GSPC(X)\}$ .

**Proof:** Let  $A$  be subset of  $X$  by definition of  $rgw\alpha$ -closure  $rgw\alpha cl(A) = \bigcap \{F : A \subseteq F \in RGW\alpha C(X)\}$ . If  $A \subseteq F \in RGW\alpha C(X)$  then  $A \subseteq F \in GSPC(X)$ , because every  $rgw\alpha$ -closed is  $gsp$ -closed i.e.  $gspcl(A) \subseteq F$ . therefore  $gspcl(A) \subseteq \bigcap \{F : A \subseteq F \in RGW\alpha C(X)\} = rgw\alpha cl(A)$ .

Hence  $gspcl(A) \subseteq rgw\alpha cl(A)$ .

**Theorem 5.12:**  $rgw\alpha$ -Closure is a kuratowski-Closure operator on a space  $X$ .

**Proof:** Let  $A$  and  $B$  be the subsets of space  $X$ . i)  $rgw\alpha cl(x) = x$ ,  $rgw\alpha cl(\phi) = \phi$  ii)  $A \subseteq rgw\alpha cl(A)$  iii)  $rgw\alpha cl(A) = rgw\alpha cl(rgw\alpha cl(A))$  iv)  $rgw\alpha cl(A \cup B) = rgw\alpha cl(A) \cup rgw\alpha cl(B)$  by theorem 5.2 Hence,  $rgw\alpha$ -Closure is a Kuratowski-Closure operator on a space  $X$ .

**Definition 5.13:** For a subset  $A$  of  $X$ ,  $rgw\alpha$ -Interior of  $A$  is denoted by  $rgw\alpha int(A)$  and defined as  $rgw\alpha int(A) = \bigcup \{G : G \subseteq A \text{ and } G \text{ is } rgw\alpha\text{-open in } X\}$  or  $\bigcup \{G : G \subseteq A \text{ and } G \in RGW\alpha O(X)\}$ .

i.e.  $rgw\alpha\text{-int}(A)$  is the union of all  $rgw\alpha$ -open set contained in  $A$ .

**Theorem 5.14:** Let  $A$  and  $B$  be subset of space  $x$  then

- i)  $rgw\alpha int(X) = X$ ,  $rgw\alpha int(\phi) = \phi$
- ii)  $rgw\alpha int(A) \subseteq A$
- iii) If  $B$  is any  $rgw\alpha$ -open set contained in  $A$  then  $B \subseteq rgw\alpha int(A)$
- iv) If  $A \subseteq B$  then  $rgw\alpha int(A) \subseteq rgw\alpha int(B)$
- v)  $rgw\alpha int(A) = rgw\alpha int(rgw\alpha int(A))$ .
- vi)  $rgw\alpha int(A \cap B) = rgw\alpha int(A) \cap rgw\alpha int(B)$

**Proof:** i) and ii) by definition of  $rgw\alpha$ -Interior of  $A$ , it is obvious.

iii) Let  $B$  be any  $rgw\alpha$ -open set such that  $B \subseteq A$ . Let  $x \in B$ ,  $B$  is an  $rgw\alpha$ -open set contained in  $A$ ,  $x$  is an element of  $rgw\alpha$ -Interior of  $A$  i.e.  $x \in rgw\alpha int(A)$ . Hence  $B \subseteq rgw\alpha int(A)$ .

iv), v) vi) similar proof as theorem 5.2 and definition of  $rgw\alpha$ -Interior.

**Theorem 5.15:** If a subset  $A$  of  $X$  is  $rgw\alpha$ -open then  $rgw\alpha int(A) = A$ .

**Proof:** Let  $A$  be  $rgw\alpha$ -open subset of  $X$ . We know that  $rgw\alpha int(A) \subseteq A$  –(1) Also  $A$  is  $rgw\alpha$ -open set contained in  $A$  from theorem 5.13 iii)  $A \subseteq rgw\alpha int(A)$  –(2) hence from (1) and (2)  $rgw\alpha int(A) = A$ .

**Theorem 5.16:** If A and B are subsets of space X then  $rgwaint(A) \cup rgwaint(B) \subseteq rgwaint(A \cup B)$

**Proof:** We know that  $A \subseteq (A \cup B)$  and  $B \subseteq (A \cup B)$  we have theorem 5.13 iv)  $rgwaint(A) \subseteq rgwaint(A \cup B)$  and  $rgwaint(B) \subseteq rgwaint(A \cup B)$ . This implies that  $rgwaint(A) \cup rgwaint(B) \subseteq rgwaint(A \cup B)$ .

**Remarks 5.17:** The converse of the above theorem need not be true.

**Theorem 5.18:** If A is a subset of X then i)  $int(A) \subseteq rgwaint(A)$  ii)  $raint(A) \subseteq rgwaint(A)$ .

**Proof:** Let A be a subset of a space X. Let  $x \in int(A) \Rightarrow x \in \bigcup \{G : G \text{ is open, } G \subseteq A\}$

$\Rightarrow \exists$  an open set G s.t.  $x \in G \subseteq A \Rightarrow \exists$  an  $rgw\alpha$ -open set G s.t.  $x \in G \subseteq A$  as every open set is  $rgw\alpha$ -open set in X  $\Rightarrow x \in \bigcup \{G : G \text{ is } rgw\alpha\text{-open set in X}\} \Rightarrow x \in rgwaint(A)$ , thus  $x \in int(A) \Rightarrow x \in rgwaint(A)$ , Hence,  $int(A) \subseteq rgwaint(A)$ .

ii) Let A be a subset of space X. Let  $x \in raint(A)$ ,  $\Rightarrow x \in \bigcup \{G : G \text{ is } r\alpha\text{-open } G \subseteq A\}$

$\Rightarrow \exists$  an  $r\alpha$ -open set G s.t.  $x \in G \subseteq A$

$\Rightarrow \exists$  an  $rgw\alpha$ -open set G s.t.  $x \in G \subseteq A$ , as every  $r\alpha$ -open set is an  $rgw\alpha$ -open set in X  $\Rightarrow x \in \bigcup \{G : G \text{ is } rgw\alpha\text{-open set in X}\} \Rightarrow x \in rgwaint(A)$ .

Thus  $x \in raint(A) \Rightarrow x \in rgwaint(A)$ .

Hence  $raint(A) \subseteq rgwaint(A)$ .

**Remark 5.19:** Containment relation in the above theorem may be proper as seen from the following example.

**Example 5.20:** Let  $X = \{a, b, c, d, e\}$  with topology  $\tau = \{\emptyset, \{a\}, \{d\}, \{e\}, \{a, d\}, \{a, e\}, \{a, d, e\}\}$   $A = \{a, b\}$   
 $int(A) = \{a\}$ ,  $raint(A) = \{a\}$ ,  $rgwaint(A) = \{a, b\}$  therefore  $int(A) \subset rgwaint(A)$  and  $raint(A) \subset rgwaint(A)$

**Theorem 5.21:** If A is subset of X, then  $rgwaint(A) \subseteq gspint(A)$ , where  $gspint(A)$  is given by  $gspint(A) = \bigcup \{G \subseteq X : G \text{ is } gsp\text{-open, } G \subseteq A\}$ .

**Proof:** Let A be a subset of a space X. Let  $x \in rgwaint(A) \Rightarrow x \in \bigcup \{G : G \text{ is } rgw\alpha\text{-open } G \subseteq A\}$

$\Rightarrow \exists$  an  $rgw\alpha$ -open set G s.t.  $x \in G \subseteq A$ , as every  $rgw\alpha$ -open set is an  $gsp$ -open set in X  $\Rightarrow x \in \bigcup \{G : G \text{ is } gsp\text{-open } G \subseteq A\} \Rightarrow x \in gspint(A)$ .

Thus,  $x \in rgwaint(A) \Rightarrow x \in gspint(A)$  Hence,  $rgwaint(A) \subseteq gspint(A)$ .

**Theorem 5.22:** For any subset A of X

i)  $X\text{-}rgwaint(A) = rgw\alpha cl(X-A)$

ii)  $X\text{-}rgw\alpha cl(A) = rgwaint(X-A)$

**Proof:**  $x \in X\text{-}rgwaint(A)$ , then x is not in  $rgwaint(A)$  i.e. every  $rgw\alpha$ -open set G containing x such that  $G \subseteq A$ . This implies every  $rgw\alpha$ -open set G containing x intersects  $(X-A)$  i.e.  $G \cap (X-A) \neq \emptyset$ . Then by theorem 5.7  $x \in rgw\alpha cl(X-A)$   
 Therefore  $X\text{-}rgw\alpha cl(A) \subseteq rgw\alpha cl(X-A)$ ---(1)

and let  $x \in rgw\alpha cl(X-A)$ , then every  $rgw\alpha$ -open set G containing x intersects  $X-A$  i.e.  $G \cap (X-A) \neq \emptyset$ . i.e. every  $rgw\alpha$ -open set G containing x s.t.  $G \subseteq A$ . Then by definition 5.12. x is not in  $rgw\alpha cl(A)$ , i.e.  $x \in X\text{-}rgwaint(A)$  and so  $rgw\alpha cl(X-A) \subseteq X\text{-}rgwaint(A)$ ---(2)

Thus  $X\text{-}rgwaint(A) = rgw\alpha cl(X-A)$ . Similarly we can prove ii).

## 6. $rgw\alpha$ -Neighbourhood and $rgw\alpha$ -Limit points

In this section we define the notation of  $rgw\alpha$ -Neighbourhood,  $rgw\alpha$ -Limit points and  $rgw\alpha$ -Derived set and some of their basic properties and analogous to those for open sets.

**Definition 6.1:** Let  $(X, \tau)$  be a topological space and let  $x \in X$ , A subset  $N$  of  $X$  is said to be  $rgw\alpha$ -Neighbourhood of  $x$  if there exists an  $rgw\alpha$ -open set  $G$  s.t.  $x \in G \subseteq N$ .

**Definition 6.2:** i) Let  $(X, \tau)$  be a topological space and  $A$  be a subset of  $X$ , A subset  $N$  of  $X$  is said to be  $rgw\alpha$  Neighbourhood of  $A$ , if there exists an  $rgw\alpha$ -open set  $G$  s.t.  $A \subseteq G \subseteq N$

ii) The collection of all  $rgw\alpha$ -Neighbourhood of  $x \in X$  called  $rgw\alpha$ -Neighbourhood system at  $x$  and shall be denoted by  $rgw\alpha N(x)$

**Definition 6.3:** i) Let  $(X, \tau)$  be a topological space and  $A$  be a subset of  $X$ , then a point  $x \in X$  is called a  $rgw\alpha$ -Limit point of  $A$  if every  $rgw\alpha$ -Neighbourhood of  $x$  contains a point of  $A$  distinct from  $x$  i.e.  $(N - \{x\}) \cap A \neq \emptyset$  for each  $rgw\alpha$ -Neighbourhood  $N$  of  $x$ . Also equivalently iff, every  $rgw\alpha$ -open set  $G$  containing  $x$  contains a point of  $A$  other than  $x$ .

ii) The set of all  $rgw\alpha$ -Limit points of the set  $A$  is called Derived set of  $A$  and is denoted by  $rgwad(A)$ .

**Theorem 6.4:** Every neighbourhood  $N$  of  $x \in X$  is called is a  $rgw\alpha$ -Neighbourhood of  $x \in X$ .

**Proof:** Let  $N$  be neighbourhood of point  $x \in X$ . To prove that  $N$  is a  $rgw\alpha$ -Neighbourhood of  $x$  by definition of neighbourhood,  $\exists$  an open set  $G$  s.t.  $x \in G \subseteq N \Rightarrow \exists$  an  $rgw\alpha$ -open set  $G$  s.t.  $x \in G \subseteq N$ , as every open set is  $rgw\alpha$ -open set. Hence  $N$  is  $rgw\alpha$ -Neighbourhood of  $x$ ,

**Remark 6.5:** In general, a  $rgw\alpha$ -nbhd  $N$  of  $x \in X$ . need not be a nbhd of  $x$  in  $X$ , as seen from the following example.

**Example 6.6 :** Let  $X = \{a, b, c, d, e\}$  with topology  $\tau = \{x, \phi, \{a\}, \{d\}, \{e\}, \{a, d\}, \{a, e\}, \{d, e\}, \{a, d, e\}\}$ . The set  $\{a, b\}$  is  $rgw\alpha$ -Neighbourhood of the point  $b$ , since  $\exists$  the  $rgw\alpha$ -open set  $\{b\}$  s.t.  $b \in \{b\} \subseteq \{a, b\}$ , However the set  $\{a, b\}$  is not a nbhd of the point  $b$ . Since no open set  $G$  exists s.t.  $b \in G \subseteq \{a, b\}$

**Theorem 6.7:** If a subset  $N$  of a space  $X$  is  $rgw\alpha$ -open, then  $N$  is  $rgw\alpha$ -nbhd of each of its points.

**Proof:** Suppose  $N$  is  $rgw\alpha$ -open. Let  $x \in N$ . We claim that  $N$  is  $rgw\alpha$ -nbhd of  $x$ . For  $N$  is a  $rgw\alpha$ -open set such that  $x \in N \subseteq N$ . Since  $x$  is an arbitrary point of  $N$ , it follows that  $N$  is a  $rgw\alpha$ -nbhd of each of its points.

**Remark 6.8:** Let  $X = \{a, b, c, d, e\}$  with topology  $\tau = \{X, \phi, \{a\}, \{d\}, \{e\}, \{a, d\}, \{a, e\}, \{d, e\}, \{a, d, e\}\}$ . The set  $\{b, c\}$  is a  $rgw\alpha$ -nbhd of the point  $b$ , since the  $rgw\alpha$ -open set  $\{b\}$  is s.t.  $b \in \{b\} \subseteq \{b, c\}$ , Also the set  $\{b, c\}$  is  $rgw\alpha$ -nbhd of the point  $c$ , Since the  $rgw\alpha$ -open set  $\{c\}$  is s.t.  $c \in \{c\} \subseteq \{b, c\}$ . That is  $\{b, c\}$  is a  $rgw\alpha$ -nbhd of each of its points. However the set  $\{b, c\}$  is not  $rgw\alpha$ -open set in  $X$ .

**Theorem 6.10:** Let  $X$  be a topological space. If  $F$  is a  $rgw\alpha$ -closed subset of  $X$ , and  $x \in F^c$ . Prove that there exists  $rgw\alpha$ -nbhd  $N$  of  $x$  such that  $N \cap F = \emptyset$ . **Proof:** let  $F$  be  $rgw\alpha$ -closed subset of  $X$  and  $x \in F^c$ . Then  $F^c$  is  $rgw\alpha$ -open set of  $X$ . So by theorem 6.7  $F^c$  contains a  $rgw\alpha$ -nbhd of each of its points. Hence there exists a  $rgw\alpha$ -nbhd of  $N$  of  $x$  such that  $N \subseteq F^c$ . That is  $N \cap F = \emptyset$ .

**Theorem 6.11:** Let  $X$  be a topological space and for each  $x \in X$ , Let  $rgw\alpha$ - $N(x)$  be the collection of all  $rgw\alpha$ -nbhd of  $x$ . Then we have following results.

- i)  $\forall x \in X, rgw\alpha$ - $N(x) \neq \emptyset$ .

- ii)  $N \in rgw\alpha\text{-}N(x) \Rightarrow x \in N$ .
- iii)  $N \in rgw\alpha\text{-}N(x), M \supset N \Rightarrow M \in rgw\alpha\text{-}N(x)$
- iv)  $N \in rgw\alpha\text{-}N(x), M \in rgw\alpha\text{-}N(x) \Rightarrow N \cap M \in rgw\alpha\text{-}N(x)$
- v)  $N \in rgw\alpha\text{-}N(x) \Rightarrow$  There exists  $M \in rgw\alpha\text{-}N(x)$  such that  $M \in N$  &  $M \in rgw\alpha\text{-}N(y)$ , for every  $y \in M$ .

**Proof:** i) Since  $X$  is a  $rgw\alpha$ -open set, it is  $rgw\alpha$ -nbhd of every  $x \in X$ . Hence there exists at least one  $rgw\alpha$ -nbhd (namely  $X$ ) for each  $x \in X$ . Hence  $rgw\alpha\text{-}N(x) \neq \emptyset$  for every  $x \in X$ .

ii) If  $N \in rgw\alpha\text{-}N(x)$ , then  $N$  is a  $rgw\alpha$ -nbhd of  $x$ , so by definition of  $rgw\alpha$ -nbhd,  $x \in N$ . Let  $N \in rgw\alpha\text{-}N(x)$  and  $M \in N$ . Then there is a  $rgw\alpha$ -open set  $G$  such that  $x \in G \subset N$ . Since  $N \subset M$ ,  $x \in G \subset M$  and so  $M$  is  $rgw\alpha$ -nbhd of  $x$ . Hence  $M \in rgw\alpha\text{-}N(x)$ .

iv) Let  $N \in rgw\alpha\text{-}N(x)$  and  $M \in rgw\alpha\text{-}N(x)$ . Then by definition of  $rgw\alpha$ -nbhd there exists  $rgw\alpha$ -open sets  $G_1$  and  $G_2$  such that  $x \in G_1 \subset N$  and  $x \in G_2 \subset M$ .

Hence  $x \in G_1 \cap G_2 \subset N \cap M$ ---(1). Since  $G_1 \cap G_2$  is a  $rgw\alpha$ -open set (being the intersection of two  $rgw\alpha$ -open sets) it follows from (1) that  $N \cap M$  is also  $rgw\alpha$ -nbhd of  $x$ . Hence  $N \cap M \in rgw\alpha\text{-}N(x)$ .

v) If  $N \in rgw\alpha\text{-}N(x)$ , then there exists a  $rgw\alpha$ -open set  $M$  such that  $x \in M \subset N$ . Since  $M$  is  $rgw\alpha$ -open set, it is  $rgw\alpha$ -nbhd of each of its points. Therefore  $M \in rgw\alpha\text{-}N(y)$  for every  $y \in M$ .

**Theorem 6.12:** Let  $X$  be a non empty set, and for each  $x \in X$ , let  $rgw\alpha\text{-}N(x)$  be a nonempty collection of subsets of  $X$  satisfying following conditions.

- i)  $N \in rgw\alpha\text{-}N(x) \Rightarrow x \in N$
- ii)  $N \in rgw\alpha\text{-}N(x), M \in rgw\alpha\text{-}N(x) \Rightarrow N \cap M \in rgw\alpha\text{-}N(x)$

Let  $\tau$  consists of the empty set and all those non-empty subsets of  $G$  of  $X$  having the property that  $x \in G$  implies that there exists an  $N \in rgw\alpha\text{-}N(x)$  such that  $x \in N \subset G$ . Then  $\tau$  is a topology for  $X$ .

**Proof:**

- (i)  $\emptyset \in \tau$  by definition. We now show that  $X \in \tau$ . Let  $x$  be any arbitrary element of  $X$ . Since  $rgw\alpha\text{-}N(x)$  is nonempty, there is  $N \in rgw\alpha\text{-}N(x)$  and so  $x \in N$  by (i). Since  $N$  a subset of  $X$ , we have  $x \in N \subset X$ . Hence  $X \in \tau$ .
- (ii) Let  $G_1 \in \tau$  and  $G_2 \in \tau$ . if  $x \in G_1 \cap G_2$  Then  $x \in G_1, x \in G_2$ . Since  $G_1 \in \tau, G_2 \in \tau$ , there exist  $N \in rgw\alpha\text{-}N(x)$  and  $M \in rgw\alpha\text{-}N(x)$ , such that  $x \in N \subset G_1$  and  $x \in M \subset G_2$ . Then  $x \in N \cap M \subset G_1 \cap G_2$ . But  $N \cap M \in rgw\alpha\text{-}N(x)$  by theorem 6.11 (iv) Hence  $G_1 \cap G_2 \in \tau$ .
- (iii) Let  $G_\lambda \in \tau$  for every  $\lambda \in \Lambda$ . If  $x \in \bigcup \{G_\lambda : \lambda \in \Lambda\}$ , then  $x \in G_{\lambda_x}$  for some  $\lambda_x \in \Lambda$ . Since  $G_{\lambda_x} \in \tau$ , there exists an  $N \in rgw\alpha\text{-}N(x)$  such that  $x \in N \subset G_{\lambda_x}$  and consequently  $x \in N \subset \bigcup \{G_\lambda : \lambda \in \Lambda\}$ . Hence  $\bigcup \{G_\lambda : \lambda \in \Lambda\} \in \tau$ . It follows that  $\tau$  is a topology for  $X$ .

**Theorem 6.13:** Let  $X$  be a topological space then

- i)  $rgw\alpha d(A) = \emptyset$
- ii) If  $A \subset B \Rightarrow rgw\alpha d(A) \subset rgw\alpha d(B)$
- iii)  $rgw\alpha d(A \cup B) = rgw\alpha d(A) \cup rgw\alpha d(B)$

**Proof:** i) Suppose that  $rgw\alpha d(A) \neq \emptyset$  then  $rgw\alpha d(A)$  contains at least one element. Therefore let  $x \in rgw\alpha d(A)$  then  $x$  is a  $rgw\alpha$ -Limit point of  $\emptyset$  therefore for every  $rgw\alpha$ -open set  $G$  containing 'x',  $(G - \{x\}) \cap \emptyset \neq \emptyset$ , But this is not true. Since intersection of  $\emptyset$  with any set is again a  $\emptyset$ . Therefore  $rgw\alpha d(A) = \emptyset$ .

ii) Given  $A \subset B$  to prove  $rgw\alpha d(A) \subset rgw\alpha d(B)$ . let  $x \in rgw\alpha d(A)$ .  $\Rightarrow x$  is a  $rgw\alpha$ -limit point of  $A$ . Therefore by definition,  $\exists$  an  $rgw\alpha$ -open set  $G$  containing  $x$  such that  $(G - \{x\}) \cap A \neq \emptyset$ ---(1). But  $A \subset B \Rightarrow A - \{x\} \subset B - \{x\} \Rightarrow (G - \{x\}) \cap B \neq \emptyset$ .  $\Rightarrow x$  is a  $rgw\alpha$ -limit point of  $B \Rightarrow x \in rgw\alpha d(B)$ . Thus  $x \in rgw\alpha d(A) \Rightarrow x \in rgw\alpha d(B)$ . Therefore  $rgw\alpha d(A) \subset rgw\alpha d(B)$

iii) We have  $A \subset A \cup B$  and  $B \subset A \cup B$ . Therefore  $rgw\alpha d(A) \subset rgw\alpha d(A \cup B)$  and  $rgw\alpha d(B) \subset rgw\alpha d(A \cup B)$ . Therefore  $rgw\alpha d(A) \cup rgw\alpha d(B) \subset rgw\alpha d(A \cup B)$ . ---(1). To prove  $rgw\alpha d(A \cup B) \subset rgw\alpha d(A) \cup rgw\alpha d(B)$ . Let  $x \in rgw\alpha d(A \cup B) \Rightarrow x$  is  $rgw\alpha$ -limit point of  $(A \cup B)$ .

$\Rightarrow (G - \{x\}) \cap (A \cup B) \neq \emptyset$  for every  $rgw\alpha$ -open set  $G$  containing  $x$ .  $\Rightarrow [(G - \{x\}) \cap A] \cup [(G - \{x\}) \cap B] \neq \emptyset \Rightarrow (G - \{x\}) \cap A \neq \emptyset$  or  $(G - \{x\}) \cap B \neq \emptyset \Rightarrow x$  is a  $rgw\alpha$ -limit point of  $A$  or  $x$  is a  $rgw\alpha$ -limit point of  $B$ . i.e.  $x \in rgw\alpha d(A)$  or  $x \in rgw\alpha d(B)$  therefore  $x \in rgw\alpha d(A) \cup rgw\alpha d(B)$ .

For  $x \in rgw\alpha d(A \cup B) \Rightarrow x \in rgw\alpha d(A) \cup rgw\alpha d(B)$ .  $\Rightarrow rgw\alpha d(A \cup B) \subset rgw\alpha d(A) \cup rgw\alpha d(B)$ ---(2)

$\Rightarrow$  From (1) and (2)  $rgw\alpha d(A \cup B) = rgw\alpha d(A) \cup rgw\alpha d(B)$

**Theorem 6.14:** Let  $X$  be a topological space and  $A \subset X$ . Then  $AUrgwad(A)$  is  $rg\omega\alpha$ -closed set in  $X$ .

**Proof:** To prove  $AUrgwad(A)$  is a  $rg\omega\alpha$ -closed set in  $X$ . that is to prove  $X - AUrgwad(A)$  is an  $rg\omega\alpha$ -open set in  $X$ .  
 Let  $x \in X - AUrgwad(A) \Rightarrow x \in X$  &  $x \notin AUrgwad(A) \Rightarrow x \in X$  &  $(x \notin A$  &  $x \notin rgwad(A)) \Rightarrow x \in X$  &  $(x \notin A$  &  $x$  is not a limit point of  $A$ ).  
 $\Rightarrow x \in X, x \notin A$ , there exist an  $rg\omega\alpha$ -open set  $G$  containing  $x$  s.t.  $G \cap (A - \{x\}) = \emptyset$  i.e.  $G \cap A = \emptyset$ . Further,  $G \cap rgwad(A) = \emptyset$ . Let  $y \in G$ . then  $y \notin A$  because  $G \cap A = \emptyset$ . Now  $G$  is an  $rg\omega\alpha$ -open set containing  $y$  and  $G \cap A = \emptyset$  and  $y \in A$ . therefore  $G$  is an  $rg\omega\alpha$ -open set containing  $y$  s.t.  $G \cap (A - \{y\}) = \emptyset$ . Therefore there exist an  $rg\omega\alpha$ -open set  $G$  containing  $y$  s.t.  $G \cap (A - \{y\}) = \emptyset$ . Therefore  $y$  is not a limit point of  $A$ . i.e.  $y \notin rgwad(A)$ .  
 $y \in G, y \notin rgwad(A)$ . therefore  $G \cap rgwad(A) = \emptyset$ . Thus we have  $G \cap A = \emptyset$  and  $G \cap rgwad(A) = \emptyset$ .  
 $\Rightarrow (G \cap A) \cup (G \cap rgwad(A)) = \emptyset$ .  
 $\square G \cap AUrgwad(A) = \emptyset \Rightarrow G \subset X - AUrgwad(A)$ . Thus for all  $x \in \{X - (AUrgwad(A))\}$  there exist an open set  $G$  s.t.  $x \in G \subset \{X - (AUrgwad(A))\} \Rightarrow X - (AUrgwad(A))$  is an  $rg\omega\alpha$ -open set. Therefore  $AUrgwad(A)$  must be  $rg\omega\alpha$ -closed set in  $X$ .

**Theorem 6.15:** Let  $X$  be a topological space and  $A \subset X$ , then  $A$  is  $rg\omega\alpha$ -closed iff  $A \supset rgwad(A)$  i.e.  $A$  is  $rg\omega\alpha$ -closed if and only if  $A$  contains all its  $rg\omega\alpha$ -limit points. i.e.  $A$  is  $rg\omega\alpha$ -closed if and only if  $rgwad(A) \subset A$ .

**Proof:** Suppose  $A$  is  $rg\omega\alpha$ -closed set, To prove  $A \supset rgwad(A)$  i.e.  $rgwad(A) \subset A$ . Let  $x \notin A$ , we prove  $x \notin rgwad(A)$ . Since  $x \notin A$ , we have  $x \in X - A$ .

Now  $X - A$  is an  $rg\omega\alpha$ -open set containing  $x$  and  $(X - A) \cap A = \emptyset$ . i.e.  $(X - A) \cap (A - \{x\}) = \emptyset$ . There exist an  $rg\omega\alpha$ -open set  $(X - A)$  containing  $x$  s.t.  $(X - A) \cap (A - \{x\}) = \emptyset$ . Therefore  $x$  is not a limit point of  $A$ .  $x \notin rgwad(A)$ . Thus  $x \notin A \Rightarrow x \notin rgwad(A)$ . therefore  $A \supset rgwad(A)$  i.e.  $rgwad(A) \subset A$ .

Conversely, on the other hand suppose  $A \supset rgwad(A)$  i.e.  $rgwad(A) \subset A$ . we prove  $A$  is  $rg\omega\alpha$ -closed set i.e. we prove  $X - A$  is  $rg\omega\alpha$ -open set.

Let  $x \in X - A \Rightarrow x \notin A \Rightarrow x \notin rgwad(A)$ .  $\Rightarrow x$  is not a limit point of  $A$ .  $\Rightarrow$  there exist an  $rg\omega\alpha$ -open set  $G$  containing  $x$  s.t.  $G \cap (A - \{x\}) = \emptyset \Rightarrow$  there exist an  $rg\omega\alpha$ -open set  $G$  containing  $x$  s.t.  $G \cap A = \emptyset \Rightarrow$  there exist an  $rg\omega\alpha$ -open set  $G$  containing  $x$  s.t.  $G \subset X - A \Rightarrow$  there exist an  $rg\omega\alpha$ -open set  $G$  containing  $x$  s.t.  $x \in G \subset X - A$ . for all  $x \in X - A$  there exist an  $rg\omega\alpha$ -open set  $G$  containing  $x$  s.t.  $x \in G \subset X - A$ . therefore  $(X - A)$  must be an  $rg\omega\alpha$ -open set. Therefore  $A$  must be a  $rg\omega\alpha$ -closed set.

**Theorem 6.16:** Let  $X$  be topological space and  $A \subset X$  then  $rg\omega\alpha cl(A) = AUrgwad(A)$ .

**Proof:** w.k.t.  $AUrgwad(A)$  is  $rg\omega\alpha$ -closed set in  $X$ . Also we have  $A \subset AUrgwad(A)$ . Therefore  $AUrgwad(A)$  is a closed set containing  $A$ . But  $rg\omega\alpha cl(A)$  is the smallest closed set containing  $A$ . Therefore  $rg\omega\alpha cl(A) \subset AUrgwad(A)$ . (1)

Further we have  $A \subset rgwad(A)$  (i). To prove  $rgwad(A) \subset rg\omega\alpha cl(A)$ . Let  $x \in rgwad(A)$ .  $\Rightarrow x$  is a  $rg\omega\alpha$ -limit point of  $A$ . We prove that  $x \in rg\omega\alpha cl(A)$ . If possible let  $x \notin rg\omega\alpha cl(A)$ . Then  $x \in X - rg\omega\alpha cl(A)$ , Therefore  $X - rg\omega\alpha cl(A)$  is an  $rg\omega\alpha$ -open set containing  $x$  and  $[X - rg\omega\alpha cl(A)] \cap [A - \{x\}] = \emptyset$ . Therefore  $x$  is not a limit point of  $A$ . Which is wrong. Therefore  $x \in rg\omega\alpha cl(A)$ . If  $x \in rgwad(A)$  then  $x \in rg\omega\alpha cl(A) \Rightarrow rgwad(A) \subset rg\omega\alpha cl(A)$  (ii)

From (i) and (ii)  $AUrgwad(A) \subset rg\omega\alpha cl(A)$  (2)

From (1) and (2)  $rg\omega\alpha cl(A) = AUrgwad(A)$ .

## REFERENCES

1. A. pushpalatha studies on generalizations of Mappings in topological spaces, *Ph.D. Thesis, Bharattiar University Coimbatore, 2000*.
2. A.Vadivel and K. Vairamanickam  $rg\alpha$ -closed sets and  $rg\alpha$ -open sets in topological spaces *Int. Jl. math. Analysis Vol, 3, 2009, no. 37, 1803-1819*.
3. D. Andrijevic, Semi preopen sets, *math. vesnik, 38 (1986) 24-32*.
4. H.Maki, R.Devi and K Balachandran generalized  $\alpha$ -closed sets in topology *Bull. Fukuoka univ. Ed. Part III 42, (1993) 13-21*
5. J. Dontchev on generalizing semi-pre open sets, *Mem, fac. Sci. kochi univ. ser. Math. 16, 35-48 (1995)*.
6. M.Stone application of the theory of Boolean rings to general.
7. O. Njastad on some classes of Nearly open sets *pacific J. Math, 15, 1965, 961-970*.
8. P. Sundaram & M.Sheik John, On w-closed sets in topology. *Acta ciencia Indica 4, 389-39 (2000)*.
9. S.S. Benchelli, P.G.Patil and Pallavi S. Mirajakar  $g$  star  $\omega\alpha$ -closed sets in topological spaces. *Jl. New results in sciences, Vol. 9, (2015), 37-45*.
10. S.S. Benchelli, P.G.Patil and P.M.Nalwad,  $g\omega\alpha$ -closed sets in topological spaces. *Journal of New results in Sci.vol.7, (2014), 7-19*.

11. S.S. Benchelli, P.G.Patil and T.D. Rayanagaudar,  $w\alpha$ -closed sets in topological spaces. *The Global J Appl. Math. And Math, Sci*, 2, 2009.
12. S.S. Benchelli & R.S. Wali on  $rw$ -closed sets in topological spaces, *Bull Malays math, sci soc* 30 99-110(2007).
13. R.S.Wali & Vijayalaxmi R.Patil on  $rgw\alpha$ -closed sets in topological spaces, *Journal of comp. and Math.Sci*.vol.8, issue3, March 2017.

**Source of support: Nil, Conflict of interest: None Declared.**

***[Copy right © 2017. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]***