

ON THE SUM CONNECTIVITY GOURAVA INDEX

V. R. KULLI*

Department of Mathematics, Gulbarga University, Gulbarga - 585106, India.

Received On: 15-06-17; Revised & Accepted On: 30-06-17)

ABSTRACT

We introduce the sum connectivity Gourava index of a molecular graph. In this paper, we determine the sum connectivity Gourava index of some standard classes of graphs. We also compute the sum connectivity Gourava index of linear $[n]$ -Tetracene, V-Tetracenic nanotube, H-Tetracenic nanotube and Tetracenic nanotori.

Mathematics Subject Classification: 05C05, 05C12, 05C35.

Keywords: sum connectivity Gourava index, nanostructures.

1. INTRODUCTION

Let G be a finite, simple and connected graph with a vertex set $V(G)$ and an edge set $E(G)$. The degree $d_G(v)$ of a vertex v is the number of vertices adjacent to v . We refer to [1] for undefined term and notation.

A molecular graph is a simple graph related to the structure of a chemical compound. Each vertex of this graph represents an atom of the molecule and its edges to the bonds between atoms. A topological index is a numerical parameter mathematically derived from the graph structure. These indices are useful for establishing correlation between the structure of a molecular compound and its physico-chemical properties, see [2].

The first and second Zagreb indices [2] of a molecular graph G are defined as

$$M_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)], \quad M_2(G) = \sum_{uv \in E(G)} d_G(u)d_G(v).$$

Motivated by the definition of the Zagreb indices and their wide applications, Kulli introduced the first Gourava index of a molecular graph in [3] as follows:

The first Gourava index of a graph G is defined as

$$GO_1(G) = \sum_{uv \in E(G)} [(d_G(u) + d_G(v)) + (d_G(u)d_G(v))].$$

The second Gourava index [3] of a molecular graph G is defined as

$$GO_2(G) = \sum_{uv \in E(G)} (d_G(u) + d_G(v))(d_G(u)d_G(v)).$$

In [4], Kulli introduced the product connectivity Gourava index of a graph G and it is defined as

$$PGO(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{(d_G(u) + d_G(v))(d_G(u)d_G(v))}}.$$

In [5], Zhou and Trinajstić introduced the sum connectivity index of a graph G and it is defined as

$$S(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_G(u) + d_G(v)}}$$

Corresponding Author: V. R. Kulli*

Department of Mathematics, Gulbarga University, Gulbarga - 585106, India.

Motivated by the definition of the sum connectivity index, we define the sum connectivity Gourava index of a graph G and it is defined as

$$SGO(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{(d_G(u) + d_G(v)) + (d_G(u)d_G(v))}}.$$

Recently, many other topological indices were studied, for example, in [6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17].

In this paper, the sum connectivity Gourava index of certain nanostructures are determined. Also the sum connectivity Gourava index of some standard classes of graphs are determined.

2. RESULTS FOR SOME STANDARD CLASSES OF GRAPHS

Proposition 1: If C_n is a cycle with $n \geq 3$ vertices, then $SGO(C_n) = \frac{n}{2\sqrt{2}}$.

Proof: Let C_n be a cycle with n vertices and n edges. Then

$$SGO(C_n) = n \frac{1}{\sqrt{(2+2) + (2 \times 2)}} \frac{n}{2\sqrt{2}}$$

Proposition 2: If K_n is a complete graph with $n \geq 2$ edges, then $SGO(K_n) = \frac{n\sqrt{(n-1)}}{2\sqrt{(n+1)}}$

Proof: Let K_n be a complete graph. Then $E(K_n) = \frac{n(n-1)}{2}$.

$$SGO(K_n) = \frac{n(n-1)}{2} \frac{1}{\sqrt{[(n-1) + (n-1)] + (n-1)(n-1)}} = \frac{n\sqrt{(n-1)}}{2\sqrt{(n+1)}}.$$

Proposition 3: If $K_{m,n}$ is a complete bipartite graph with $1 \leq m \leq n$, then $SGO(K_{m,n}) = \frac{mn}{\sqrt{mn + m + n}}$.

Proof: Let $K_{m,n}$ be a complete bipartite graph with $m + n$ vertices and mn edges such that $|V_1| = m$, $|V_2| = n$, $V(K_{m,n}) = V_1 \cup V_2$. Every vertex of V_1 is adjacent with n vertices and every vertex of V_2 is adjacent with m vertices.

$$SGO(K_{m,n}) = \frac{mn}{\sqrt{(m+n) + mn}} = \frac{mn}{\sqrt{mn + m + n}}.$$

Corollary 3.1: Let $K_{n,n}$ be a complete bipartite graph. Then $SGO(K_{n,n}) = \frac{n\sqrt{n}}{\sqrt{2+n}}$.

Corollary 3.2: Let $K_{1,n}$ be a Star. Then $SGO(K_{1,n}) = \frac{n}{\sqrt{1+2n}}$.

Proposition 4: If G is an r -regular graph with n vertices, then $SGO(G) = \frac{n\sqrt{r}}{2\sqrt{2+r}}$.

Proof: Let G be an r -regular graph with n vertices and $\frac{nr}{2}$ edges. Then the degree of each vertex of G is r .

$$SGO(G) = \frac{nr}{2} \frac{1}{\sqrt{(r+r) + r^2}} = \frac{n\sqrt{r}}{2\sqrt{2+r}}.$$

Proposition 5: Let P_n be a path with $n \geq 3$ vertices. Then $SGO(P_n) = \frac{n}{2\sqrt{2}} + \frac{2}{\sqrt{5}} - \frac{3}{2\sqrt{2}}$.

Proof: Let $G=P_n$ be a path with $n \geq 3$ vertices. We obtain two partitions of the edge set of P_n as follows:

$$E_3 = \{uv \in E(G) \mid d_G(u)=1, d_G(v)=2\}, \mid E_3 \mid = 2.$$

$$E_4 = \{uv \in E(G) \mid d_G(u) = d_G(v)=2\}, \mid E_4 \mid = n - 3.$$

To compute $SGO(P_n)$, we see that

$$SGO(P_n) = \frac{1}{\sqrt{(1+2)+(1 \times 2)}} 2 + \frac{1}{\sqrt{(2+2)+(2 \times 2)}} (n-3) = \frac{n}{2\sqrt{2}} + \frac{2}{\sqrt{5}} - \frac{3}{2\sqrt{2}}.$$

3. SUM CONNECTIVITY GOURAVA INDEX OF SOME NANOSTRUCTURES

3.1 Linear $[n]$ – Tetracene.

The molecular graph of a linear $[n]$ -Tetracene is shown in Figure -1.




Figure-1: The molecular graph of a linear $[n]$ -Tetracene

We compute the sum connectivity Gourava index of a linear $[n]$ -Tetracene.

Theorem 1: Let T be a linear $[n]$ -Tetracene. Then

$$SGO(T) = \left(\frac{16}{\sqrt{11}} + \frac{7}{\sqrt{15}} \right) n + \left(\frac{3}{\sqrt{2}} - \frac{4}{\sqrt{11}} - \frac{4}{\sqrt{15}} \right).$$

Proof: From Figure 1, by algebraic method, we obtain $|V(T)|=18n$ and $|E(T)|=23n - 2$. Also we obtain three partitions of the edge set of T as follows:

$$E_{22} = \{uv \in E(T) \mid d_T(u) = d_T(v)=2\}, \mid E_{22} \mid = 6.$$

$$E_{23} = \{uv \in E(T) \mid d_T(u) = 2, d_T(v)=3\}, \mid E_{23} \mid = 16n - 4.$$

$$E_{33} = \{uv \in E(T) \mid d_T(u) = d_T(v)=3\}, \mid E_{33} \mid = 7n - 4.$$

To compute $SGO(T)$, we see that

$$SGO(T) = \sum_{uv \in E(T)} \frac{1}{\sqrt{(d_T(u) + d_T(v)) + (d_T(u)d_T(v))}}.$$

$$= 6 \frac{1}{\sqrt{(2+2)+(2 \times 2)}} + (16n - 4) \frac{1}{\sqrt{(2+3)+(2 \times 3)}} + (7n - 4) \frac{1}{\sqrt{(3+3)+(3 \times 3)}}$$

$$= \left(\frac{16}{\sqrt{11}} + \frac{7}{\sqrt{15}} \right) n + \left(\frac{3}{\sqrt{2}} - \frac{4}{\sqrt{11}} - \frac{4}{\sqrt{15}} \right).$$

3.2. Nanostructure $F = F[p,q]$

The molecular graph of 2-D lattice of $F = F[p,q]$ with $p = 2$ and $q = 4$ is shown in Figure 2.




Figure-2: The graph of 2-D lattice of $F = F[p, q]$ with $p = 2$ and $q = 4$

In the following theorem, we compute the sum connectivity Gourava index of a nanostructure $F = F[p, q]$.

Theorem 2: Let $F = F[p, q]$ be a nanostructure. Then

$$SGO(F[p, q]) = \frac{27}{\sqrt{15}} pq + \left(\frac{16}{\sqrt{11}} - \frac{20}{\sqrt{15}} \right) p + \left(\frac{1}{\sqrt{2}} + \frac{4}{\sqrt{11}} - \frac{8}{\sqrt{15}} \right) q + \left(\frac{2}{\sqrt{2}} - \frac{8}{\sqrt{11}} - \frac{4}{\sqrt{15}} \right)$$

Proof: Let $F = F[p, q]$ be a nanostructure. By algebraic method, we obtain three partitions of the edge set of F as follows:

$$E_{22} = \{uv \in E(F) \mid d_F(u)=d_F(v)=2\}, \mid E_{22} \mid = 2q+4.$$

$$E_{23} = \{uv \in E(F) \mid d_F(u) = 2, d_F(v)=3\}, \mid E_{23} \mid = 16p + 4q - 8.$$

$$E_{33} = \{uv \in E(F) \mid d_F(u) = d_F(v)=3\}, \mid E_{33} \mid = 27pq - 20p - 8q + 4.$$

To compute $SGO(F[p, q])$, we see that

$$\begin{aligned} SGO(F[p, q]) &= \sum_{uv \in E(F)} \frac{1}{\sqrt{(d_F(u) + d_F(v)) + (d_F(u)d_F(v))}} \\ &= (2q + 4) \frac{1}{\sqrt{(2+2) + (2 \times 2)}} + (16p + 4q - 8) \frac{1}{\sqrt{(2+3) + (2 \times 3)}} \\ &\quad + (27pq - 20p - 8q + 4) \frac{1}{\sqrt{(3+3) + (3 \times 3)}} \\ &= \frac{27}{\sqrt{15}} pq + \left(\frac{16}{\sqrt{11}} - \frac{20}{\sqrt{15}} \right) p + \left(\frac{1}{\sqrt{2}} + \frac{4}{\sqrt{11}} - \frac{8}{\sqrt{15}} \right) q + \left(\frac{2}{\sqrt{2}} - \frac{8}{\sqrt{11}} + \frac{4}{\sqrt{15}} \right). \end{aligned}$$

3.3. Nanostructure $G = G[p, q]$.

The molecular graph of 2-D lattice of $G=G[p, q]$ with $p = 2$ and $q = 4$ is shown in Figure 3.




Figure-3: The graph of 2-D lattice of $G = G[p, q]$ with $p = 2$ and $q = 4$

In the following theorem, we compute the sum connectivity Gourava index of a nanostructure $G = G[p, q]$.

Theorem 3: Let $G = G[p, q]$ be a nanostructure. Then

$$SGO(G[p, q]) = \frac{27}{\sqrt{15}} pq + \left(\frac{16}{\sqrt{11}} - \frac{20}{\sqrt{15}} \right) p.$$

Proof: Let $G = G[p, q]$ be a nanostructure. By algebraic method, we obtain $|V(G)| = 18pq$ and $|E(G)| = 27pq - 4p$. Further, we obtain two partitions of the edge set of G as follows:

$$E_{23} = \{uv \in E(G) \mid d_G(u)=2, d_G(v)=3\}, \mid E_{23} \mid = 16p.$$

$$E_{33} = \{uv \in E(G) \mid d_G(u) = d_G(v)=3\}, \mid E_{33} \mid = 27pq - 20p.$$

To compute $SGO(G[p, q])$, we see that

$$SGO(G[p, q]) = \sum \frac{1}{\sqrt{(d_G(u) + d_G(v)) + (d_G(u)d_G(v))}}.$$

$$\begin{aligned}
 &= 16p \frac{1}{\sqrt{(2+3)+(2 \times 3)}} + (27pq - 20p) \frac{1}{\sqrt{(3+3)+(3 \times 3)}} \\
 &= \frac{27}{\sqrt{15}} pq + \left(\frac{16}{\sqrt{11}} - \frac{20}{\sqrt{15}} \right) p.
 \end{aligned}$$

3.4. Nanostructure $K = K [p, q]$

The molecular graph of 2-D lattice of $K = K [p, q]$ with $p = 2$ and $q = 3$ is shown in Figure 4.




Figure-4: The graph of 2-D lattice of $K = K[p, q]$ with $p = 2$ and $q = 3$

In the following theorem, we compute the the sum connectivity Gourava index of a nanostructure $K = K[p, q]$.

Theorem 4: Let $K = K [p, q]$ be a nanostructure. Then

$$SGO(K[p, q]) = \frac{27}{\sqrt{15}} pq + \left(\frac{1}{\sqrt{2}} + \frac{4}{\sqrt{11}} - \frac{8}{\sqrt{15}} \right) q.$$

Proof: Let $K = K[p, q]$ be a nanostructure. By algebraic method, we obtain $|V(K)| = 18pq$ and $|E(K)| = 27pq - 2q$.

Further, we obtain three partitions of the edge set of K as follows:

$$E_{22} = \{uv \in E(K) \mid d_K(u) = d_K(v) = 2\}, \quad |E_{22}| = 2q.$$

$$E_{23} = \{uv \in E(K) \mid d_K(u) = 2, d_K(v) = 3\}, \quad |E_{23}| = 4q.$$

$$E_{33} = \{uv \in E(K) \mid d_K(u) = d_K(v) = 3\}, \quad |E_{33}| = 27pq - 8q.$$

To compute $SGO(K[p, q])$, we see that

$$\begin{aligned}
 SGO(K[p, q]) &= \sum_{uv \in E(K)} \frac{1}{\sqrt{(d_K(u) + d_K(v)) + (d_K(u)d_K(v))}} \\
 &= 2q \frac{1}{\sqrt{(2+2)+(2 \times 2)}} + 4q \frac{1}{\sqrt{(2+3)+(2 \times 3)}} + (27pq - 8q) \frac{1}{\sqrt{(3+3)+(3 \times 3)}} \\
 &= \frac{27}{\sqrt{15}} pq + \left(\frac{1}{\sqrt{2}} + \frac{4}{\sqrt{11}} - \frac{8}{\sqrt{15}} \right) q.
 \end{aligned}$$

3.5. Nanostructure $L = L [p, q]$

The molecular graph of 2-D lattice of $L = L [p, q]$ with $p = 2$ and $q = 4$ is shown in Figure 5.




Figure-5: The graph of 2-D lattice of $L = L[p, q]$ with $p = 2$ and $q = 4$.

In the next theorem, we compute the sum connectivity Gourava index of a nanostructure $L = L[p, q]$.

Theorem 5: Let $L = L[p, q]$ be a nanostructure. Then

$$SGO(L[p, q]) = \frac{27}{\sqrt{15}} pq$$

Proof: Let $L = L[p, q]$ be a nanostructure. By algebraic method, we obtain $|V(L)| = 18pq$ and $|E(L)| = 27pq$. Since the degree of each vertex of L is 3, the edge partition of L is as follows:

$$E_{33} = \{uv \in E(L) \mid d_L(u) = d_L(v) = 3\}, \quad |E_{33}| = 27pq.$$

To compute $SGO(L[p, q])$, we see that

$$\begin{aligned} SGO(L[p, q]) &= \sum \frac{1}{\sqrt{(d_L(u) + d_L(v)) + (d_L(u)d_L(v))}} \\ &= 27pq \frac{1}{\sqrt{(3+3) + (3 \times 3)}} = \frac{27}{\sqrt{15}} pq. \end{aligned}$$

REFERENCES

1. V.R.Kulli, *College Graph Theory*, Vishwa International Publications, Gulbarga, India (2012).
2. Gutman and N. Trinajstić, Graph theory and molecular orbitals. Total ϵ -electron energy of alternant hydrocarbons, *Chem. Phys. Lett.* 17, (1972) 535-538.
3. V.R.Kulli, The Gourava indices and coindices of graphs, *Annals of Pure and Applied Mathematics*, 14(1) (2017), 33-38.
4. V.R.Kulli, The product connectivity Gourava index, *J.Comp.& Math.Sci.*8(6) (2017) 235-242..
5. B. Zhou and N. Trinajstić, On a novel connectivity index, *J. Math. Chem.* 46(2009) 1252-1270.
6. V.R. Kulli, On K indices of graphs, *International Journal of Fuzzy Mathematical Archive*, 10(2), (2016) 105-109.
7. V.R. Kulli, Multiplicative connectivity indices of $TUC_4C_8[m, n]$ and $TUC_4[m, n]$ nanotubes, *Journal of Computer and Mathematical Sciences*, 7(11) (2016) 599-605.
8. V.R. Kulli, On multiplicative K Banhatti and multiplicative K hyper-Banhatti indices of V -Phenylenic nanotubes and nanotorus, *Annals of Pure and Applied Mathematics*, 11(2) (2016) 145-150.
9. V.R.Kulli, Computation of general topological indices for titania nanotubes, *International Journal of Mathematical Archive*, 7(12) (2016) 33-38.
10. V.R.Kulli, Multiplicative connectivity indices of nanostructures, *Journal of Ultra Scientist of Physical Sciences*, A 29(1) (2017) 1-10. DOI:http://dx.doi.org/10.22147/jusps-A/290101.
11. V.R.Kulli, Two new multiplicative atom bond connectivity indices, *Annals of Pure and Applied Mathematics*, 13(1) (2017) 1-7. DOI:http://dx.doi.org/10.22457/apam.v13n1a1.
12. V.R.Kulli, Some new multiplicative geometric-arithmetic indices, *Journal of Ultra Scientist of Physical Sciences*, A, 29(2) (2017) 52-57. DOI:http://dx.doi.org/10.22147/jusps-A/290201.
13. V.R.Kulli, Computation of some topological indices of certain networks, *International Journal of Mathematical Archive*, 8(2) (2017) 99-106.
14. V.R.Kulli, New K Banhatti topological indices, *International Journal of Fuzzy Mathematical Archive*, 12(1) (2017) 29-37. DOI:http://dx.doi.org/10.22457/ijfma.v12n1a4.

15. V.R.Kulli, New arithmetic-geometric indices, *Annals of Pure and Applied Mathematics*, 13(2) (2017) 165-172. DOI:<http://dx.doi.org/10.22457/apam.v13n2a2>.
16. V.R.Kulli, A new multiplicative arithmetic-geometric index, *International Journal of Fuzzy Mathematical Archive*, 12(2) (2017) 49-53. DOI:<http://dx.doi.org/10.22457/ijfma.v12n2a1>.
17. N.Soleimani, M.J.Nikmehr and H.A.Tavallaee, Computation of the different topological indices of nanostructures, *J.Natn.Sci.Foundation Sri Lanka*,43(2) (2015) 127-133.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2017. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]