

SOME OPERATIONS ON NEUTROSOPHIC FUZZY GRAPHS

G. DIVYA*

Research Scholar in Mathematics,
K. N. Govt. Arts College for women (A), Thanjavur – 7, Tamil Nadu, India.

Dr. J. MALARVIZHI

Research Advisor, Associate Professor in Mathematics,
K. N. Govt. Arts College for women (A), Thanjavur – 7, Tamil Nadu, India.

(Received On: 05-08-17; Revised & Accepted On: 01-09-17)

ABSTRACT

The concept of neutrosophic sets can be utilized as a Mathematical tool to deal with imprecise and unspecified information. The neutrosophic set is a generalization of fuzzy sets and intuitionistic fuzzy sets which is applied in Graph theory. The neutrosophic fuzzy graph is used in the problems, where the relation between vertices and hence edges are indeterminate. In this paper complement of Neutrosophic fuzzy graph is introduced and few operations on Neutrosophic fuzzy graph (union, join) are also defined.

Keywords: Neutrosophic Fuzzy Graph (NFG), Complement, Union, Join.

1. INTRODUCTION

Graph theory has been highly successful in certain academic fields, including natural sciences and engineering. Graph theoretic models can sometimes provide a useful structure upon which analytical techniques can be used. It is often convenient to depict the relationships between pairs of elements of a system by means of a graph or a digraph. The vertices of the graph represent the system elements and its edges or arcs represent the relationships between the elements. This approach is especially useful for transportation, scheduling, sequencing, allocation, assignment, and other problems which can be modelled as networks. Such a graph theoretic model is often useful as an aid in communicating. Zadeh [6] introduced the concept of fuzzy set. Atanassov [2] introduced the intuitionistic fuzzy sets which is a generalization of fuzzy sets. Fuzzy set theory and intuitionistic fuzzy set theory are useful models for dealing with uncertainty and incomplete information. But they may not be sufficient in modeling of indeterminate and inconsistent information encountered in real world. In order to cope with this issue, neutrosophic set theory was proposed by Smarandache [8] as a generalization of fuzzy sets and intuitionistic fuzzy sets.

Neutrosophic set proposed by Smarandache [12, 13] is a powerful tool to deal with incomplete, indeterminate and inconsistent information in real world. It is a generalization of the theory of fuzzy set [18], intuitionistic fuzzy sets [19], interval-valued fuzzy sets and interval-valued intuitionistic fuzzy sets, then the neutrosophic set is characterized by a truth-membership degree (t), an indeterminacy-membership degree (i) and a falsity-membership degree (f) independently, which are within the real standard or nonstandard unit interval $] -0, 1+[$. There are some difficulties in modeling of some problems in Engineering and Sciences. To overcome these difficulties, in 2010, concept of single-valued neutrosophic sets and its operations defined by Wang et al. [4] as a generalization of intuitionistic fuzzy sets.

In this research article we apply the concept of neutrosophic sets to graphs. We introduce the notion of Neutrosophic Fuzzy Graph and present few fundamental operations.

Corresponding Author: G. Divya*

**Research Scholar in Mathematics,
K. N. Govt. Arts College for women (A), Thanjavur – 7, Tamil Nadu, India.**

2. PRELIMINARIES

Definition 2.1: Let X be a space of points with generic elements in X denoted by x ; then the neutrosophic set A (NS A) is an object having the form


$$A = \{ \langle x: T_A(x), I_A(x), F_A(x) \rangle, x \in X \}$$

where the functions $T, I, F: X \rightarrow]0, 1^+[$ define respectively a truth-membership function, an indeterminacy-membership function, and a falsity-membership function of the element $x \in X$ to the set A with the condition

$$]0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3^+$$

The functions $T_A(x), I_A(x)$ and $F_A(x)$ are real standard or nonstandard subsets of $] -0, 1^+[$.

Definition 2.2: A fuzzy graph is a pair of functions $G = (\sigma, \mu)$ where σ is a fuzzy subset of a non empty set V and is a symmetric fuzzy relation on σ . i.e $\sigma: V \times V \rightarrow [0, 1]$ and $\mu: V \times V \rightarrow [0, 1]$ such that $\mu(uv) \leq \sigma(u) \wedge \sigma(v)$ for all $u, v \in V$ where uv denotes the edge between u and v and $\sigma(u) \wedge \sigma(v)$ denotes the minimum of $\sigma(u)$ and $\sigma(v)$. σ is called the fuzzy vertex set of V and is called the fuzzy edge set of E .



Definition 2.3: The fuzzy subgraph $H = (\tau, \rho)$ is called a fuzzy subgraph of $G = (\sigma, \mu)$

If $\tau(u) \leq \sigma(u)$ for all $u \in V$ and $\rho(u, v) \leq \mu(u, v)$ for all $u, v \in V$.

Definition 2.4: An intuitionistic fuzzy graph (IFG) is of the form $G = (V, E)$ where

(i) $V = \{v_1, v_2, \dots, v_n\}$ such that $\mu_1: V \rightarrow [0, 1]$ and $\gamma_1: V \rightarrow [0, 1]$ denote the degree of membership and non-membership of the element $v_i \in V$ respectively and

$$0 \leq \mu_1(v_i) + \gamma_1(v_i) \leq 1 \text{ for every } v_i \in V, (i = 1, 2, \dots, n),$$


(ii) $E \subseteq V \times V$ where $\mu_2: V \times V \rightarrow [0, 1]$ and $\gamma_2: V \times V \rightarrow [0, 1]$ are such that

$$\mu_2(v_i, v_j) \leq \mu_1(v_i) \cdot \mu_1(v_j)$$

$$\gamma_2(v_i, v_j) \leq \gamma_1(v_i) \cdot \gamma_1(v_j)$$

and $0 \leq \mu_2(v_i, v_j) + \gamma_2(v_i, v_j) \leq 1$

for every $(v_i, v_j) \in E$, where \cdot means the ordinary multiplication.



3. NEUTROSOPHIC FUZZY GRAPHS

Throughout this paper we denote $G = (V, E)$ a crisp graph, $N_G = (A, B)$ a neutrosophic fuzzy graph of Graph G .

Definition 3.1: Let X be a space of points with generic elements in X denoted by x . A neutrosophic fuzzy set A (NFS A) is characterized by truth-membership function $T_A(x)$, an indeterminacy-membership function $I_A(x)$, and a falsity-membership function $F_A(x)$. For each point x in X

$$T_A(x), I_A(x), F_A(x) \in [0, 1].$$

A NFS A can be written as $A = \{ \langle x: T_A(x), I_A(x), F_A(x) \rangle, x \in X \}$

Definition 3.2: Let $A = (T_A, I_A, F_A)$ and $B = (T_B, I_B, F_B)$ be neutrosophic fuzzy sets on a set X . If $A = (T_A, I_A, F_A)$ is a neutrosophic fuzzy relation on a set X , then $A = (T_A, I_A, F_A)$ is called a neutrosophic fuzzy relation on $B = (T_B, I_B, F_B)$ if

$$\begin{aligned} T_B(x, y) &\leq T_A(x) \cdot T_A(y) \\ I_B(x, y) &\leq I_A(x) \cdot I_A(y) \\ F_B(x, y) &\leq F_A(x) \cdot F_A(y) \text{ for all } x, y \in X \text{ where } \cdot \text{ means the ordinary multiplication} \end{aligned}$$

Definition 3.3: A neutrosophic fuzzy graph (NF-graph) with underlying set V is defined to be a pair $N_G = (A, B)$ where

1. The functions $T_A: V \rightarrow [0, 1]$, $I_A: V \rightarrow [0, 1]$ and $F_A: V \rightarrow [0, 1]$ denote the degree of truth-membership, degree of indeterminacy-membership and falsity-membership of the element $v_i \in V$, respectively, and

$$0 \leq T_A(v_i) + I_A(v_i) + F_A(v_i) \leq 3 \tag{1}$$

2. $E \subseteq V \times V$ where, functions $T_B: V \times V \rightarrow [0, 1]$, $I_B: V \times V \rightarrow [0, 1]$ and $F_B: V \times V \rightarrow [0, 1]$ are defined by

$$T_B(v_i, v_j) \leq T_A(v_i) \cdot T_A(v_j) \tag{2}$$


$$I_B(v_i, v_j) \leq I_A(v_i) \cdot I_A(v_j) \tag{3}$$

$$F_B(v_i, v_j) \leq F_A(v_i) \cdot F_A(v_j) \tag{4}$$

for all $v_i, v_j \in V$ where \cdot means the ordinary multiplication. Denotes the degree of truth-membership, indeterminacy-membership and falsity-membership of the edge $(v_i, v_j) \in E$ respectively, where

$$0 \leq T_B(v_i, v_j) + I_B(v_i, v_j) + F_B(v_i, v_j) \leq 3 \tag{5}$$

for all $(v_i, v_j) \in E$ ($i, j = 1, 2, \dots, n$)



Neutrosophic Fuzzy Graph

We call “A” the neutrosophic fuzzy vertex set of V , “B” the neutrosophic fuzzy edge set of E , respectively.

Notation:

- μ, λ, γ denote the truth membership value, indeterminacy membership value and falsity membership value respectively
- $(v_i, \mu_i, \lambda_i, \gamma_i)$ denotes the degree of truth membership, indeterminacy membership and falsity membership of the vertex v_i .
- $(e_{ij}, \mu'_{ij}, \lambda'_{ij}, \gamma'_{ij})$ denotes the degree of truth membership, indeterminacy membership and falsity membership of the edge e_{ij} .

Note 3.4:

- (i) When $\mu'_{ij} = \lambda'_{ij} = \gamma'_{ij} = 0$, for some i and j , then there is no edge between v_i and v_j . Otherwise there exists edge between v_i and v_j .
- (ii) If one of the inequalities (1) or (2) or (3) or (4) or (5) is not satisfied, then G is not a NFG.

Definition 3.5: A NF-graph $N_H = (A', B')$ of $H = (V', E')$ is partial NF-subgraph of NF-graph $N_G = (A, B)$ of $G = (V, E)$ if

- (i) $V' \subseteq V$, where $T_{A'}(v_i) \leq T_A(v_i)$, $I_{A'}(v_i) \leq I_A(v_i)$, $F_{A'}(v_i) \leq F_A(v_i)$ for all $v_i \in V'$.
- (ii) $E' \subseteq E$, where $T_{B'}(v_i, v_j) \leq T_B(v_i, v_j)$, $I_{B'}(v_i, v_j) \leq I_B(v_i, v_j)$, $F_{B'}(v_i, v_j) \leq F_B(v_i, v_j)$ for all $(v_i, v_j) \in E'$

Definition 3.6: A NF-graph $N_H = (A', B')$ of $H = (V', E')$ is NF-subgraph of NF-graph $N_G = (A, B)$ of $G = (V, E)$ if


- (i) $V' \subseteq V$, where $T_{A'}(v_i) = T_A(v_i)$, $I_{A'}(v_i) = I_A(v_i)$, $F_{A'}(v_i) = F_A(v_i)$ for all $v_i \in V'$.
- (ii) $E' \subseteq E$, where $T_{B'}(v_i, v_j) = T_B(v_i, v_j)$, $I_{B'}(v_i, v_j) = I_B(v_i, v_j)$, $F_{B'}(v_i, v_j) = F_B(v_i, v_j)$ for all $(v_i, v_j) \in E'$

Definition 3.7: A neutrosophic fuzzy graph $N_G = (A, B)$ is called complete if

$$\begin{aligned} T_B(v_i, v_j) &= T_A(v_i) \cdot T_A(v_j) \\ I_B(v_i, v_j) &= I_A(v_i) \cdot I_A(v_j) \\ F_B(v_i, v_j) &= F_A(v_i) \cdot F_A(v_j) \text{ for all } v_i, v_j \in V \end{aligned}$$

Definition 3.8: The complement of an NFG $N_G = (A, B)$ is a NFG, $\overline{N}_G = (\overline{A}, \overline{B})$, where

- $\overline{V} = V$
- $\overline{\mu}_i = \mu_i, \overline{\lambda}_i = \lambda_i, \overline{\gamma}_i = \gamma_i$
- $\overline{\mu}_{ij} = \mu_i \cdot \mu_j - \mu_{ij}, \overline{\lambda}_{ij} = \lambda_i \cdot \lambda_j - \lambda_{ij}, \overline{\gamma}_{ij} = \gamma_i \cdot \gamma_j - \gamma_{ij}$




Note: We can easily verify that $\overline{\overline{N}_G} = N_G$.

4. OPERATIONS ON NEUTROSOPHIC FUZZY GRAPH

Definition 4.1: Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two graphs, $N_{G_1} = (A_1, B_1)$ and $N_{G_2} = (A_2, B_2)$ be the neutrosophic fuzzy graphs of the crisp graphs G_1 and G_2 respectively with $V_1 \cap V_2 = \emptyset$. $G = G_1 \cup G_2 = (V_1 \cup V_2, E_1 \cup E_2)$ be the union of G_1 and G_2 .

The union of NFGs N_{G_1} and N_{G_2} is a NFG defined by


$$\begin{aligned}
 (\mu_1 \cup \mu_2)(v) &= \begin{cases} \mu_1(v) & \text{if } v \in V_1 - V_2 \\ \mu_2(v) & \text{if } v \in V_2 - V_1 \end{cases} \\
 (\lambda_1 \cup \lambda_2)(v) &= \begin{cases} \lambda_1(v) & \text{if } v \in V_1 - V_2 \\ \lambda_2(v) & \text{if } v \in V_2 - V_1 \end{cases} \\
 (\gamma_1 \cup \gamma_2)(v) &= \begin{cases} \gamma_1(v) & \text{if } v \in V_1 - V_2 \\ \gamma_2(v) & \text{if } v \in V_2 - V_1 \end{cases} \\
 (\mu'_1 \cup \mu'_2)(e_{ij}) &= \begin{cases} \mu'_1(e_{ij}) & \text{if } e_{ij} \in E_1 - E_2 \\ \mu'_2(e_{ij}) & \text{if } e_{ij} \in E_2 - E_1 \end{cases} \\
 (\lambda'_1 \cup \lambda'_2)(e_{ij}) &= \begin{cases} \lambda'_1(e_{ij}) & \text{if } e_{ij} \in E_1 - E_2 \\ \lambda'_2(e_{ij}) & \text{if } e_{ij} \in E_2 - E_1 \end{cases} \\
 (\gamma'_1 \cup \gamma'_2)(e_{ij}) &= \begin{cases} \gamma'_1(e_{ij}) & \text{if } e_{ij} \in E_1 - E_2 \\ \gamma'_2(e_{ij}) & \text{if } e_{ij} \in E_2 - E_1 \end{cases}
 \end{aligned}$$



Definition 4.2: Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two graphs, $N_{G_1} = (A_1, B_1)$ and $N_{G_2} = (A_2, B_2)$ be the neutrosophic fuzzy graphs of the crisp graphs G_1 and G_2 respectively with $V_1 \cap V_2 = \phi$. $G = G_1 \oplus G_2 = (V_1 \cup V_2, E_1 \cup E_2 \cup E')$ be the join of G_1 and G_2 .

The join of NFGs N_{G_1} and N_{G_2} is defined by

$$\begin{aligned}
 (\mu_1 \oplus \mu_2)(v) &= (\mu_1 \cup \mu_2)(v) && \text{if } v \in V_1 \cup V_2 \\
 (\lambda_1 \oplus \lambda_2)(v) &= (\lambda_1 \cup \lambda_2)(v) && \text{if } v \in V_1 \cup V_2 \\
 (\gamma_1 \oplus \gamma_2)(v) &= (\gamma_1 \cup \gamma_2)(v) && \text{if } v \in V_1 \cup V_2 \\
 (\mu'_1 \oplus \mu'_2)(e_{ij}) &= (\mu'_1 \cup \mu'_2)(e_{ij}) && \text{if } e_{ij} \in E_1 \cup E_2 \\
 (\mu'_1 \oplus \mu'_2)(e_{ij}) &= (\mu_1(v_i) \cdot \mu_2(v_j)) && \text{if } e_{ij} \in E' \\
 (\lambda'_1 \oplus \lambda'_2)(e_{ij}) &= (\lambda'_1 \cup \lambda'_2)(e_{ij}) && \text{if } e_{ij} \in E_1 \cup E_2 \\
 (\lambda'_1 \oplus \lambda'_2)(e_{ij}) &= (\lambda_1(v_i) \cdot \lambda_2(v_j)) && \text{if } e_{ij} \in E' \\
 (\gamma'_1 \oplus \gamma'_2)(e_{ij}) &= (\gamma'_1 \cup \gamma'_2)(e_{ij}) && \text{if } e_{ij} \in E_1 \cup E_2 \\
 (\gamma'_1 \oplus \gamma'_2)(e_{ij}) &= (\gamma_1(v_i) \cdot \gamma_2(v_j)) && \text{if } e_{ij} \in E'
 \end{aligned}$$



where, $u_1 = (0.5, 0.1, 0.4)$, $u_2 = (0.6, 0.3, 0.2)$, $u_3 = (0.4, 0.2, 0.5)$, $v_1 = (0.2, 0.3, 0.4)$, $v_2 = (0.3, 0.5, 0.7)$, $e_1 = (0.2, 0.03, 0.06)$, $e_2 = (0.1, 0.01, 0.15)$, $e_3 = (0.2, 0.05, 0.1)$, $e_4 = (0.06, 0.10, 0.2)$, $e_5 = (0.1, 0.03, 0.16)$, $e_6 = (0.12, 0.09, 0.08)$, $e_7 = (0.08, 0.05, 0.28)$, $e_8 = (0.15, 0.05, 0.28)$, $e_9 = (0.18, 0.15, 0.14)$, $e_{10} = (0.12, 0.10, 0.35)$

REFERENCES

1. Akram, M., Bipolar fuzzy graphs, Information Sciences, vol.181, no.24, pp. 5548 – 5564, 2011.
2. Atanassov, K.T., Intuitionistic fuzzy sets, VII ITKR's Session, Deposited in Central for Science-Technical Library of Bulgarian Academy of Sciences, 1697/84, 1983, Sofia, Bulgaria, June, (Bulgarian).
3. Parvathi Rangasamy, M.G. Karunambigai, Operations on intuitionistic fuzzy graphs, conference paper in international journal of computer Applications – January 2009
4. Wang, H., Smarandache, F., Y. Zhang, and R. Sunderraman, Single-valued neutrosophic sets, Multispace and Multistructure, vol. 4, pp.410-413, 2010.
5. Yang, H.-L., Guo, Z.-L., Y. She and X. Liao, On single valued neutrosophic relations, Journal of Intelligent Fuzzy Systems, vol.30, pp.1045-1056, 2016.
6. Zadeh, L.A., Fuzzy sets, Information and Control, vol. 8, pp.338-353, 1965.
7. Broumi, S., Talea, M., A. Bakali and F. Smarandache, Single-valued neutrosophic graphs, Journal of New Theory, vol.10, pp.86 – 101, 2016.
8. Smarandache, F., A Unifying field in logics neutrosophy: Neutrosophic probability, set and logic, Re-hoboth: American Research Press, 1998.
9. Muhammad Akram and Gulfam Shahzadi, Operations on Single Valued Neutrosophic Graphs , Journal of uncertain systems vol.11, No.1 , pp 1-26, 2017.
10. K.T. Atanassov, Intuitionistic fuzzy sets: Theory and applications, Physica-verlag, New-York, 1999.
11. A. Nagoor Gani and S.R Latha, On Irregular Fuzzy Graphs, Applied Mathematical Sciences, Vol.6, 2012, no.11, 517-523.
12. F. Smarandache, Neutrosophic set - a generalization of the intuitionistic fuzzy set, Granular Computing, 2006 IEEE International Conference, 38 – 42, 2006, DOI: 10.1109/GRC.2006.1635754.
13. F. Smarandache, A geometric interpretation of the neutrosophic set — A generalization of the intuitionistic fuzzy set Granular Computing (GrC), 2011 IEEE International Conference , 602 – 606, 2011, DOI 10.1109/GRC.2011.6122665.
14. H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman, Single valued Neutrosophic Sets, Multispace and Multistructure 4 (2010) 410-413.
15. K. Atanassov, —Intuitionistic fuzzy sets, Fuzzy Sets and Systems, vol. 20, p. 87-96 (1986).
16. L. Zadeh, Fuzzy sets, Inform and Control, 8(1965), 338-353
17. S. Broumi, M. Talea, F. Smarandache, Single Valued Neutrosophic Graphs: Degree, Order and Size, IEEE WCCI 2016.
18. H.J. Zimmermann, Fuzzy Set Theory and its Applications, Kluwer-Nijhoff, Boston, 1985.
19. H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman, Single valued Neutrosophic Sets, Multispace and Multistructure 4 (2010) 410-413.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2017. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]