

CRITICAL AND STABILITY CONCEPT FOR STRONG DOMINATING SET IN FUZZY GRAPHS

¹R. JAHIR HUSSAIN, ²S. AMEENA BANU*

¹PG and Research Department of Mathematics,
Jamal Mohamed College (Autonomous), Tiruchirappalli-620020, India.

²Research Scholar, Department of Mathematics,
Jamal Mohamed College (Autonomous), Tiruchirappalli-620020, India.

(Received On: 16-09-17; Revised & Accepted On: 09-10-17)

ABSTRACT

In this paper we introduce the critical and stability concept to strong fuzzy dominating set in fuzzy graphs. The strong fuzzy dominating critical node is a node whose removal increases (or) decreases the strong fuzzy domination number. The stability of strong fuzzy dominating set is the minimum number of nodes whose removal increases (or) decreases the strong domination number.

Key words: Fuzzy dominating set, strong fuzzy dominating set, strong fuzzy dominating critical node, strong fuzzy dominating stability, strong fuzzy domination number.

AMS Mathematics Subject Classification: 03E72, 05C72.

I. INTRODUCTION

Harary et al [1] explained an interesting application in voting situations using the concept of domination. Rosenfeld [5] introduced the notion of fuzzy graph and several fuzzy analogous of graph theoretic concepts such as paths, cycles, connectedness and etc. Somasundaram.A and Somasundaram.S [6] discussed domination in fuzzy graphs. Nagoor Gani.A and Basheer Ahamed.M [2] introduced strong and weak domination in fuzzy graphs. Ramachandra.S.R and Soner.N.D.[4] introduced strong domination critical and stability in graphs. In this paper we investigate the changes in the fuzzy cardinality of strong fuzzy dominating sets when we remove the nodes from G.

II. PRELIMINARIES

A fuzzy graph $G=(\sigma, \mu)$ is a non-empty set V together with a pair of functions $\sigma: V \rightarrow [0,1]$ and $\mu: V \times V \rightarrow [0,1]$ such that $\mu(u,v) \leq \sigma(u) \wedge \sigma(v)$ for all $u,v \in V$, where $\sigma(u) \wedge \sigma(v)$ is the minimum of $\sigma(u)$ and $\sigma(v)$. The underlying crisp graph of the fuzzy graph $G=(\sigma, \mu)$ is denoted as $G^*=(\sigma^*, \mu^*)$ where $\sigma^*=\{u \in V / \sigma(u) > 0\}$ and $\mu^*=\{(u,v) \in V \times V / \mu(u,v) > 0\}$. A fuzzy graph $G=(\sigma, \mu)$ is a complete fuzzy graph if $\mu(u,v) = \sigma(u) \wedge \sigma(v)$ for all $u,v \in \sigma^*$. Two nodes u and v are said to be neighbours if $\mu(u,v) > 0$. The strong neighbourhood of u is $N_S(u) = \{v \in V : (u,v) \text{ is a strong arc}\}$. $N_S[u] = N_S(u) \cup \{u\}$ is the closed strong neighbourhood of u . A path ρ in a fuzzy graph is a sequence of distinct nodes $u_0, u_1, u_2, \dots, u_n$ such that $\mu(u_{i-1}, u_i) > 0$; $1 \leq i \leq n$ here $n \geq 0$ is called the length of the path ρ . The consecutive pairs (u_{i-1}, u_i) are called the arcs of the path. A path ρ is called a cycle if $u_0 = u_n$ and $n \geq 3$. An arc (u,v) is said to be a strong arc if $\mu(u,v) \geq \mu^\infty(u,v)$ and the node v is said to be a strong neighbour of u .

If $\mu(u,v) = 0$ for every $v \in V$ then u is called isolated node. A fuzzy graph $G=(\sigma, \mu)$ is fuzzy bipartite if it has a spanning fuzzy sub graph $H=(\tau, \pi)$ which is bipartite where for all edges (u,v) not in H , weight of (u,v) in G is strictly less than the strength of pair (u,v) in H . i.e $\mu(u,v) < \pi^\infty(u,v)$. A fuzzy bipartite graph G with fuzzy bipartition (V_1, V_2) is said to be a complete fuzzy bipartite if for each node of V_1 , every node of V_2 is a strong neighbor. Let $G=(\sigma, \mu)$ be a fuzzy graph and u be a node in G then there exist a node v such that (u,v) is a strong arc then we say that u dominates v . Let $G=(\sigma, \mu)$ be a fuzzy graph. A set D of V is said to be fuzzy dominating set of G if every $v \in V-D$ there exist $u \in D$ such that u dominates v . A fuzzy dominating set D of G is called a minimal fuzzy dominating set of G if no proper subset of D is a fuzzy dominating set. The fuzzy domination number $\gamma_f(G)$ of the fuzzy graph G is the smallest number of nodes in any fuzzy dominating set of G . A fuzzy dominating set D of a fuzzy graph G such that $|D| = \gamma_f(G)$ is called minimum fuzzy dominating set.

Corresponding Author: ²S. Ameena Banu*

III. FUZZY DOMINATING CRITICAL NODES

Definition 3.1: Let $G=(\sigma, \mu)$ be a fuzzy graph. A node v of G is said to be fuzzy dominating critical node if its removal either increases (or) decreases the fuzzy domination number.

We partition the nodes of G into three disjoint sets according to how their removal affects $\gamma_f(G)$. Let $V = V_f^0 \cup V_f^+ \cup V_f^-$ for

$$V_f^0 = \{v \in V: \gamma_f(G-v) = \gamma_f(G)\}$$

$$V_f^+ = \{v \in V: \gamma_f(G-v) > \gamma_f(G)\}$$

$$V_f^- = \{v \in V: \gamma_f(G-v) < \gamma_f(G)\}$$

RESULTS

1. If I is the set of all isolated nodes of G then $I \subseteq V_f^-$.
2. If D is a fuzzy dominating set, removing any node in $V-D$ cannot increase the fuzzy domination number therefore removing any node in D increase the $\gamma_f(G)$. In other words $D \subseteq V_f^+$.

Definition 3.2: Let $G = (\sigma, \mu)$ be a fuzzy graph. Two nodes u and v of G are *strong adjacent* if (u,v) is strong arc. Otherwise they are said to be weak.

The *strong degree* of a node v is the number of nodes that are strong adjacent to v . It is denoted by $d_s(v)$.

Let $G = (\sigma, \mu)$ be a fuzzy graph. For any $u, v \in V$, u *strongly dominates* v (i) If u is strong adjacent to v and (ii) $d_s(u) \geq d_s(v)$.

A set $D \subseteq V$ is a *strong fuzzy dominating set* of G if every node in $V-D$ is strongly dominated by at least one node in D . A strong fuzzy dominating set D of G is called a minimal strong fuzzy dominating set if no proper subset of D is a strong fuzzy dominating set of G .

The smallest number of nodes in any strong fuzzy dominating set is called the *strong domination number* and is denoted by $\gamma_{sf}(G)$. A strong fuzzy dominating set D of G such that $|D| = \gamma_{sf}(G)$ is called minimum strong fuzzy dominating set.

IV. STRONG AND WEAK FUZZY DOMINATING CRITICAL NODES

Definition 4.1: Let $G=(\sigma, \mu)$ be a fuzzy graph. A node v of G is said to be strong fuzzy dominating critical node if its removal either increase (or) decrease the fuzzy strong domination number.

We partition the nodes of G into three disjoint sets according to how their removal affects $\gamma_{sf}(G)$.

$$\text{Let } V = V_{sf}^0 \cup V_{sf}^+ \cup V_{sf}^-$$

$$\text{For } V_{sf}^0 = \{v \in V: \gamma_{sf}(G-v) = \gamma_{sf}(G)\}$$

$$V_{sf}^+ = \{v \in V: \gamma_{sf}(G-v) > \gamma_{sf}(G)\}$$

$$V_{sf}^- = \{v \in V: \gamma_{sf}(G-v) < \gamma_{sf}(G)\}$$

Definition 4.2:

γ_{sf} - Stability of fuzzy graph G written γ_{sf} is the minimum number of nodes whose removal changes $\gamma_{sf}(G)$.

γ_{sf}^+ - Stability of a fuzzy graph G written γ_{sf}^+ is the minimum number of nodes whose removal increases $\gamma_{sf}(G)$.

γ_{sf}^- - Stability of a fuzzy graph G written γ_{sf}^- is the minimum number of nodes whose removal decreases $\gamma_{sf}(G)$.

V. STRONG FUZZY DOMINATING CRITICAL IN FUZZY GRAPHS

Proposition 5.1: If the removal of a node u from G increases $\gamma_{sf}(G)$, then (i) u is not isolated node and end node and (ii) there is no strong fuzzy dominating set for $G-N_s[u]$ having $\gamma_{sf}(G)$ nodes which also dominates $N_s[u]$ for some γ_{sf} - set D containing u .

Proof:

- (i) Suppose $\gamma_{sf}(G-v) > \gamma_{sf}(G)$ and $u \in D$. Then clearly u is not an isolated, and also u is not end -node, sine for any fuzzy graph G if u is end-node then $\gamma_{sf}(G-v) \leq \gamma_{sf}(G)$, a contradiction.
- (ii) Suppose there exists a weak fuzzy dominating set of $G-N_s[u]$ with $\gamma_{sf}(G)$ nodes. Then $\gamma_{sf}(G-v) \leq \gamma_{sf}(G)$, a contradiction.

VI. STRONG FUZZY DOMINATING STABILITY IN FUZZY PATHS

Proposition 6.1: For any fuzzy path $\rho_n, \gamma_{sf}^+(\rho_n) = \begin{cases} 1 \text{ if } n \equiv 0 \pmod{3}, n \geq 3 \\ 3 \text{ if } n \equiv 1 \pmod{3}, n \geq 7 \\ 2 \text{ if } n \equiv 2 \pmod{3}, n \geq 5 \end{cases}$

Proof: We consider the following three cases.

Case-(i): Let $n \equiv 0 \pmod{3}, n \geq 3$. Let v_1, v_2, \dots, v_n be the nodes of ρ_n then $\rho_n - \{v_2\}$ consists of an isolated node and a fuzzy path of order $n-2$. Thus, $\gamma_{sf}(\rho_n - \{v_2\}) = \gamma_{sf}(\rho_1) + \gamma_{sf}(\rho_{n-2}) = 1 + \left\lfloor \frac{(n-2)}{3} \right\rfloor > \gamma_{sf}(\rho_n)$. Hence $\gamma_{sf}^+(\rho_n) = 1$.

Case-(ii): Let $n \equiv 1 \pmod{3}, n \geq 7$. Let v_1, v_2, \dots, v_n be the nodes of ρ_n . Then $\rho_n - \{v_2, v_4, v_6\}$ consists of 3 isolated nodes and a path of order $n-6$. Thus $\gamma_{sf}(\rho_n - \{v_2, v_4, v_6\}) = 3(\gamma_{sf}(\rho_1)) + \gamma_{sf}(\rho_{n-6}) = 3 + \left\lfloor \frac{(n-6)}{3} \right\rfloor > \gamma_{sf}(\rho_n)$. Hence $\gamma_{sf}^+(\rho_n) = 3$.

Case-(iii): Let $n \equiv 2 \pmod{3}, n \geq 5$. Then $\rho_n - \{v_2, v_4\}$ consists of two isolated nodes and a path of order $n-4$. Thus $\gamma_{sf}(\rho_n - \{v_2, v_4\}) = 2(\gamma_{sf}(\rho_1)) + \gamma_{sf}(\rho_{n-4}) = 2 + \left\lfloor \frac{(n-4)}{3} \right\rfloor > \gamma_{sf}(\rho_n)$. Hence $\gamma_{sf}^+(\rho_n) = 2$.

Proposition 6.2: For any fuzzy path ρ_n with $n \geq 4$

$$\gamma_{sf}^-(\rho_n) = \begin{cases} 1 \text{ if } n \equiv 1 \pmod{3} \\ 2 \text{ if } n \equiv 2 \pmod{3} \\ 3 \text{ if } n \equiv 0 \pmod{3} \end{cases}$$

Proof:

Case-(i): Let $n \equiv 1 \pmod{3}$. Let v_1, v_2, \dots, v_n be the nodes of ρ_n . Then $\rho_n - \{v_1\}$ consists of a path of order $n-1$. Then $\gamma_{sf}(\rho_n - \{v_1\}) = \gamma_{sf}(\rho_{n-1}) = \left\lfloor \frac{(n-1)}{3} \right\rfloor$ since $n \equiv 1 \pmod{3}$ then $n-1 = 3K < \left\lfloor \frac{n}{3} \right\rfloor = \gamma_{sf}(\rho_n)$. Thus $\gamma_{sf}^-(\rho_n) = 1$.

Case-(ii): Let $n \equiv 2 \pmod{3}$ and Let v_1, v_2, \dots, v_n be the nodes of ρ_n . Then $\rho_n - \{v_1, v_2\}$ consists of a path of order $n-2$. Then $\gamma_{sf}(\rho_n - \{v_1, v_2\}) = \gamma_{sf}(\rho_{n-2}) = \left\lfloor \frac{(n-2)}{3} \right\rfloor < \left\lfloor \frac{n}{3} \right\rfloor = \gamma_{sf}(\rho_n)$. Hence $\gamma_{sf}^-(\rho_n) = 2$.

Case-(iii): Let $n \equiv 0 \pmod{3}$ and Let v_1, v_2, \dots, v_n be the nodes of ρ_n . Then $\rho_n - \{v_1, v_2, v_3\}$ consists of a path of order $n-3$. Then $\gamma_{sf}(\rho_n - \{v_1, v_2, v_3\}) = \gamma_{sf}(\rho_{n-3}) = \left\lfloor \frac{(n-3)}{3} \right\rfloor < \left\lfloor \frac{n}{3} \right\rfloor = \gamma_{sf}(\rho_n)$. Hence $\gamma_{sf}^-(\rho_n) = 3$.

VII. STRONG FUZZY DOMINATING STABILITY IN FUZZY CYCLES

Proposition 7.1: For any fuzzy cycle C_p with $n \geq 6$ nodes

$$\gamma_{sf}^+(C_n) = \begin{cases} 3 \text{ if } n \equiv 0 \pmod{3}, n \geq 6 \\ 5 \text{ if } n \equiv 1 \pmod{3}, n \geq 10 \\ 7 \text{ if } n \equiv 2 \pmod{3}, n \geq 14 \end{cases}$$

Proof: Let $C_n = v_1, v_2, v_3, \dots, v_n$. We consider the following three cases.

Case-(i): Let $n \equiv 0 \pmod{3}, n \geq 6$. Then the removal of the set of nodes $\{v_1, v_3, v_5\}$ leaves two isolated nodes and a path of order $n-5$. Thus $\gamma_{sf}(C_n - \{v_1, v_3, v_5\}) = 2(\gamma_{sf}(\rho_1)) + \gamma_{sf}(\rho_{n-5}) = 2 + \left\lfloor \frac{(n-5)}{3} \right\rfloor > \gamma_{sf}(C_n)$. Hence $\gamma_{sf}^+(C_n) = 3$.

Case-(ii): Let $n \equiv 1 \pmod{3}, n \geq 10$. Then the removal of the set of nodes $\{v_1, v_3, v_5, v_7, v_9\}$ leaves four isolated nodes and a path of order $n-9$. Thus $\gamma_{sf}(C_n - \{v_1, v_3, v_5, v_7, v_9\}) = 4(\gamma_{sf}(\rho_1)) + \gamma_{sf}(\rho_{n-9}) = 4 + \left\lfloor \frac{(n-9)}{3} \right\rfloor > \gamma_{sf}(C_n)$. Hence $\gamma_{sf}^+(C_n) = 5$.

Case-(iii): Let $n \equiv 2 \pmod{3}, n \geq 14$. Then the removal of the set of nodes $\{v_1, v_3, v_5, v_7, v_9, v_{11}, v_{13}\}$ leaves six isolated nodes and a path of order $n-13$. Thus $\gamma_{sf}(C_n - \{v_1, v_3, v_5, v_7, v_9, v_{11}, v_{13}\}) = 6(\gamma_{sf}(\rho_1)) + \gamma_{sf}(\rho_{n-13}) = 6 + \left\lfloor \frac{(n-13)}{3} \right\rfloor > \gamma_{sf}(C_n)$. Hence $\gamma_{sf}^+(C_n) = 7$.

Proposition 7.2: For any fuzzy cycle C_n with $n \geq 6$ nodes

$$\gamma_{sf}^-(C_n) = \begin{cases} 1 \text{ if } n \equiv 1 \pmod{3} \\ 2 \text{ if } n \equiv 2 \pmod{3} \\ 3 \text{ if } n \equiv 0 \pmod{3} \end{cases}$$

Proof:

Case-(i): Let $n \equiv 1 \pmod{3}$. Let v_1, v_2, \dots, v_p be the nodes of C_n . Then $C_n - \{v_1\}$ consists of a path of order $n-1$. Then $\gamma_{sf}(C_n - \{v_1\}) = \gamma_{sf}(\rho_{n-1}) = \left\lfloor \frac{(n-1)}{3} \right\rfloor < \left\lfloor \frac{n}{3} \right\rfloor = \gamma_{sf}(C_n)$. Hence $\gamma_{sf}^-(C_n) = 1$.

Case-(ii): Let $n \equiv 2 \pmod{3}$. Let v_1, v_2, \dots, v_p be the nodes of C_n . Then $C_n - \{v_1, v_2\}$ consists of a path of order $n-2$. Then $\gamma_{sf}(C_n - \{v_1, v_2\}) = \gamma_{sf}(\rho_{n-2}) = \left\lfloor \frac{(n-2)}{3} \right\rfloor < \left\lfloor \frac{n}{3} \right\rfloor = \gamma_{sf}(C_n)$. Hence $\gamma_{sf}^-(C_n) = 2$.

Case-(iii): Let $n \equiv 0 \pmod{3}$. Let v_1, v_2, \dots, v_p be the nodes of C_n . Then $C_n - \{v_1, v_2, v_3\}$ consists of a path of order $n-3$. Then $\gamma_{sf}(C_n - \{v_1, v_2, v_3\}) = \gamma_{sf}(\rho_{n-3}) = \left\lfloor \frac{(n-3)}{3} \right\rfloor < \left\lfloor \frac{n}{3} \right\rfloor = \gamma_{sf}(C_n)$. Hence $\gamma_{sf}^-(C_n) = 3$.

VIII. STRONG AND WEAK FUZZY DOMINATING STABILITY IN COMPLETE FUZZY GRAPHS AND COMPLETE FUZZY BIPARTITE GRAPHS

Proposition 8.1: Let $K_{m,n}$ be a complete fuzzy bipartite graph, which is neither K_2 nor $K_{2,2}$ then

$$\gamma_{sf}^+(K_{m,n}) = \begin{cases} 1 & \text{if } 1 = m < n \\ 1 & \text{if } m = n \geq 4 \\ m & \text{if } m < n \end{cases}$$

Proof: Suppose a complete fuzzy bipartite graph is star. Then $\gamma_{sf}(K_{1,n}) = 1$. Let v be a node of strong degree n . Then $K_{1,n} - v$ has at least two components. Hence $\gamma_{sf}(K_{1,n} - v) \geq 2 > 1 = \gamma_{sf}(K_{1,n})$. Therefore $\gamma_{sf}^+(K_{1,n}) = 1$.

Suppose $K_{m,n}$ is not a star and $m = n \geq 4$. We know that $\gamma_{sf}(K_{m,n}) = 2$ (if $m=n$). Let $V = V_1 \cup V_2$ be a vertex set of $K_{m,n}$ such that $|V_1| \leq m, |V_2| \leq n$. Let $v \in V_1$ so $\gamma_{sf}(K_{m,n} - v) = m-1 = 3 > \gamma_{sf}(K_{m,n}) = 2$.

Proposition 8.2: Let $K_{m,n}$ be a complete fuzzy bipartite graph which is neither K_2 nor $K_{2,2}$ then

$$\gamma_{sf}^-(K_{m,n}) = \begin{cases} m+n-1 & \text{if } m = n \\ 1 & \text{if } 3 \leq m \leq n \end{cases}$$

Proof: Suppose $m=n$ and $V = V_1 \cup V_2$ be the vertex set of $K_{m,n}$ such that $|V_1| = m, |V_2| = n$. Then $\gamma_{sf}(K_{m,n} - V_1 - V_2 + 1) = 1 < 2 = \gamma_{sf}(K_{m,n})$. Hence $\gamma_{sf}^-(K_{m,n}) = m+n-1$.

Suppose $3 \leq m \leq n$ $\gamma_{sf}(K_{m,n}) = m$ if $m < n$. Let $v \in V_1$ so $\gamma_{sf}(K_{m,n} - v) = m-1 < m = \gamma_{sf}(K_{m,n})$. Hence $\gamma_{sf}^-(K_{m,n}) = 1$.

Theorem 8.3: A vertex v of $V(G)$ is in V_{sf}^0 if and only if $G = \bigcup_{i=2}^t K_i$ where $t \geq 2$ and $K_i, i = 2, 3, \dots, t$ are complete fuzzy graphs.

Proof: Let D be an γ_{sf} -set of G . Clearly $|D| = t-1$. Since there is no path between $K_i, i = 2, 3, \dots, t$. Let $V = V_2 \cup V_3 \cup \dots \cup V_t$ where V_i are vertex sets of K_i respectively. Suppose $u \in V_r, 2 \leq r \leq t$. As $K_r - u$ is a complete fuzzy graph of order at least $r-1$ and $\gamma_{sf}(K_r - u) = \gamma_{sf}(K_r)$ then D will still γ_{sf} -set of G if such a node is removed thus for any $u \in V$. We have $u \in V_{sf}^0$. Conversely let $V(G) = V_{sf}^0(G)$ and D be an γ_{sf} -set of G . Note that every node of strong degree at least one must be in D . Also note that all nodes of $V-D$ must be in $\langle N_S[D] \rangle$. Furthermore there can be no path in G , connecting two nodes of D . A graph with these properties is only union of complete fuzzy graphs of order at least two.

Theorem 8.4: A vertex v of $V(G)$ is in V_{sf}^+ if and only if $G = \bigcup_{r=2}^t K_{r,r}$

Proof: Let D be an γ_{sf} -set of G . Clearly $|D| = 2(t-3)$. Since there is no path between $K_{i,i}, i = 4, 5, \dots, t$. Let $V = V_4 \cup V_5 \cup \dots \cup V_t$ where V_i are vertex sets of $K_{i,i}$ respectively. Suppose $u \in V_r, 4 \leq r \leq t$. As $K_{r,r} - u$ is a complete fuzzy graph of order at least $r-1$ and $\gamma_{sf}(K_{r,r} - u) = \gamma_{sf}(K_{r,r})$ then D will still γ_{sf} -set of G if such a vertex is removed thus for any $u \in V$. We have $u \in V_{sf}^+$. Conversely let $V(G) = V_{sf}^+(G)$ and D be an γ_{sf} -set of G . Note that every node of strong degree at least one must be in D . Also note that all nodes of $V-D$ must be in $\langle N_S[D] \rangle$. Furthermore there can be no path in G , connecting two nodes of D . A graph with these properties is only union of complete fuzzy graphs of order at least four.

CONCLUSION

In this paper we discussed critical and stability concept of strong fuzzy dominating set for several classes of fuzzy graphs and obtained some results. Here we considered all the complete fuzzy bipartite graphs $K_{m,n}$ except K_2 and $K_{2,2}$.

REFERENCES

1. Harary.F, Bauer.D, Nieminen.J and Suffel.C.L, Domination alteration sets in graphs, Discrete Math.47, 153-161(1983).
2. Nagoor Gani. A and Basheer Ahamed,M, Strong and weak domination in fuzzy graphs, East Asian Math J. 23(2007).
3. Nagoor Gani.A and Chandrasekaran.V.T , A First look at Fuzzy Graph Theory, Allied publishers (p) Ltd, 2010.
4. Ramachandra.S.R and Soner.N.D, Strong domination critical and stability in graphs, J.Comp, & Math.Sci. Vol 1(3), 294-299 (2010).
5. Rosenfeld A (1975) Fuzzy graphs: In: Zadeh LA, FU KS, Shimura M (eds) Fuzzy sets and their applications Academic press, New York.
6. Somasundaram.A and Somasundaram.S, Domination in fuzzy graphs, Pattern Recognition Letters, 19(9), 787-791, (1998).

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2017. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]