

NON EXISTENCE OF RELAXED SKOLEM MEAN LABELING FOR STAR GRAPHS

A. MANSHATH¹, V. BALAJI*² AND P. SEKAR³

¹Department of Mathematics, Sriram Engineering College, Thiruvallur-602024, India.

²Department of Mathematics, Sacred Heart College, Tirupattur-635601, India.

³Department of Mathematics, C. Kandasamy Naidu College, Chennai - 600102 India.

(Received On: 31-07-17; Revised & Accepted On: 25-09-17)

ABSTRACT

In this paper we prove that the two star graph $K_{1,m} \cup K_{1,n}$ is not a relaxed skolem mean graph if $|m - n| > 5$ and the three star graph $K_{1,\ell} \cup K_{1,m} \cup K_{1,n}$ is not a relaxed skolem mean graph if $|m - n| > \ell + 6$.

Keywords: Relaxed Skolem mean graph and star.

2010 Mathematical Subject Classification Number: 05C78.

1. INTRODUCTION

We proved that $K_{1,m}$ is not a relaxed skolem mean graph for $m \geq 5$. Also, we proved that the two star $K_{1,m} \cup K_{1,n}$ is a relaxed skolem mean graph if $|m-n| \leq 5$. Next, we proved the existence of relaxed skolem mean graphs. In [3], the three star $K_{1,\ell} \cup K_{1,m} \cup K_{1,n}$ is a relaxed skolem mean graph if $|m-n| \leq \ell + 6$ for $\ell = 1, 2, 3, \dots$; $m = 1, 2, 3, \dots$; if $n = \ell + m + 6$ when $\ell \leq m < n$.

2. [4] RELAXED SKOLEM MEAN LABELING

Definition 2.1: A graph $G = (V, E)$ with p vertices and q edges is said to be a relaxed skolem mean graph if there exists a function f from the vertex set of G to $\{1, 2, 3, \dots, p+1\}$ such that the induced map f^* from the edge set of G to $\{2, 3, 4, \dots, p+1\}$ defined by

$$f^*(e=uv) = \begin{cases} \frac{f(u)+f(v)}{2} & \text{if } f(u)+f(v) \text{ is even} \\ \frac{f(u)+f(v)+1}{2} & \text{if } f(u)+f(v) \text{ is odd, then} \end{cases}$$

the resulting edges get distinct labels from the set $\{2, 3, 4, \dots, p+1\}$.

Note 2.2: [4], In a Relaxed skolem mean graph, $p \geq q$.

Theorem 2.3: The two star $G = K_{1,m} \cup K_{1,n}$ is not a relaxed Skolem mean graph if $|m - n| > 5$.

Proof: Without loss of generality, let us consider $m \leq n$. Consider the primal graph under the condition $|m - n| > 5$ that is $m = 1$ and $n = 7$. That is $G = K_{1,1} \cup K_{1,7}$.

*Corresponding Author: V. Balaji*²*

²Department of Mathematics, Sacred Heart College, Tirupattur-635601, India.

Where $V(G) = \{v_{1,j} : 0 \leq j \leq 1\} \cup \{v_{2,j} : 0 \leq i \leq 7\}$

$$E(G) = \{v_{1,0}v_{1,1}\} \cup \{v_{2,0}v_{2,j} : 1 \leq j \leq 7\}$$

G has 10 vertices and 8 edges.

Suppose G is relaxed skolem mean graph.

Then there exists a function f from the vertex set of G to $\{1, 2, 3, \dots, 11\}$ such that the induced map f^* from the edge set of G to $\{2, 3, \dots, 11\}$ defined by

$$f^*(e = uv) = \begin{cases} \frac{f(u) + f(v)}{2} & \text{if } f(u) + f(v) \text{ is even} \\ \frac{f(u) + f(v) + 1}{2} & \text{if } f(u) + f(v) \text{ is odd} \end{cases}$$

Then the resulting edges get distinct labels from the set $\{2, 3, \dots, 11\}$.

Then the vertex and edge mappings of G is given by

$$f : V(G) \rightarrow \{1, 2, \dots, 11\}$$

$$f^* : E(G) \rightarrow \{2, 3, \dots, 11\}$$

Let, $t_{i,j} = f(v_{i,j})$ and $x_{i,j} = f^*(v_{i,0} v_{i,j}) \forall i$ and j .

Now let us consider the following cases,

Case-(a): $t_{2,0} = 11$.

If $t_{2,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then,

$$x_{2,j} = f^*(v_{2,0}v_{2,j}) = \left(\frac{11+2n}{2}\right) = 6+n = \left(\frac{11+2n+1}{2}\right) = f^*(v_{2,0}v_{2,k}) = x_{2,k}$$

Therefore, the possibilities for the pendent vertices are (1), (2 or 3), (4 or 5), (6 or 7), (8 or 9), (10). These six labels are not sufficient to label seven vertices, $t_{2,j}$ for $1 \leq j \leq 7$.

Suppose $t_{2,7}$ takes any of the remaining values.

Let $t_{2,0} = 11, t_{2,1} = 1, t_{2,2} = 3, t_{2,7} = 2$.

Then the corresponding edge labels are $x_{2,1} = 6, x_{2,2} = 7, x_{2,7} = 7$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{2,0} = 11$.

Case-(b): $t_{2,0} = 10$

If $t_{2,j} = 2n - 1$ and $t_{2,k} = 2n$ for some n, j and k then,

$$x_{2,j} = f^*(v_{2,0}v_{2,j}) = \left(\frac{10+2n-1}{2}\right) = 5+n = \left(\frac{10+2n}{2}\right) = f^*(v_{2,0}v_{2,k}) = x_{2,k}$$

Therefore, the possibilities for the pendent vertices are (1 or 2), (3 or 4), (5 or 6), (7 or 8), 9 and 11. These six labels are not sufficient to label seven vertices, $t_{2,j}$ for $1 \leq j \leq 7$.

Suppose $t_{2,7}$ takes any of the remaining values.

Let $t_{2,0} = 10$, $t_{2,1} = 2$, $t_{2,2} = 4$, $t_{2,7} = 1$.

Then the corresponding edge labels are $x_{2,1} = 6$, $x_{2,2} = 7$, $x_{2,7} = 6$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{2,0} = 10$.

Case-(c): $t_{2,0} = 9$.

If $t_{2,j} = 2n$ and $t_{2,k} = 2n+1$ for some n, j and k then,

$$x_{2,j} = f^*(v_{2,0}v_{2,j}) = \left(\frac{9+2n}{2}\right) = 5+n = \left(\frac{9+2n+1}{2}\right) = f^*(v_{2,0}v_{2,k}) = x_{2,k}$$

Therefore, the possibilities for the pendent vertices are (1), (2 or 3), (4 or 5), (6 or 7), 8, (10 or 11). These six labels are not sufficient to label seven vertices, $t_{2,0}$ for $1 \leq j \leq 7$.

Suppose $t_{2,7}$ takes any of the remaining values.

Let $t_{2,0} = 9$, $t_{2,1} = 1$, $t_{2,2} = 3$, $t_{2,7} = 2$.

Then the corresponding edge labels are $x_{2,1} = 5$, $x_{2,2} = 6$, $x_{2,7} = 6$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{2,0} = 9$.

Case- (d): $t_{2,0} = 8$.

If $t_{2,j} = 2n-1$ and $t_{2,k} = 2n$ for some n, j and k then,

$$x_{2,j} = f^*(v_{2,0}v_{2,j}) = \left(\frac{1+2n}{2}\right) = 1+n = \left(\frac{1+2n+1}{2}\right) = f^*(v_{2,0}v_{2,k}) = x_{2,k}$$

Therefore, the possibilities for the pendent vertices are (1 or 2), (3 or 4), (5 or 6), (7), (9 or 10), (11). These six labels are not sufficient to label seven vertices, $t_{2,j}$ for $1 \leq j \leq 7$.

Suppose $t_{2,7}$ takes any of the remaining values.

Let $t_{2,0} = 8$, $t_{2,1} = 2$, $t_{2,2} = 4$, $t_{2,7} = 1$.

Then the corresponding edge labels are $x_{2,1} = 5$, $x_{2,2} = 6$, $x_{2,7} = 5$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{2,0} = 8$.

Case-(e): $t_{2,0} = 7$.

If $t_{2,j} = 2n$ and $t_{2,k} = 2n+1$ for some n, j and k then,

$$x_{2,j} = f^*(v_{2,0}v_{2,j}) = \left(\frac{7+2n}{2}\right) = 4+n = \left(\frac{7+2n+1}{2}\right) = f^*(v_{2,0}v_{2,k}) = x_{2,k}$$

Therefore, the possibilities for the pendent vertices are (1), (2 or 3), (4 or 5), (6), (8 or 9), (10 or 11). These six labels are not sufficient to label seven vertices, $t_{2,j}$ for $1 \leq j \leq 7$.

Suppose $t_{2,7}$ takes any of the remaining values.

Let $t_{2,0} = 7, t_{2,1} = 1, t_{2,2} = 3, t_{2,7} = 2$.

Then the corresponding edge labels are $x_{2,1} = 4, x_{2,2} = 5, x_{2,7} = 5$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{2,0} = 7$.

Case-(f): $t_{2,0} = 6$.

If $t_{2,j} = 2n - 1$ and $t_{2,k} = 2n$ for some n, j and k then,

$$x_{2,j} = f^*(v_{2,0}v_{2,j}) = \left(\frac{6+2n-1}{2}\right) = 3+n = \left(\frac{6+2n}{2}\right) = f^*(v_{2,0}v_{2,k}) = x_{2,k}$$

Therefore, the possibilities for the pendent vertices are (1 or 2), (3 or 4), (5), (7 or 8), (9 or 10), (11). These six labels are not sufficient to labels seven vertices, $t_{2,j}$ for $1 \leq j \leq 7$.

Suppose $t_{2,7}$ takes any of the remaining values.

Let $t_{2,0} = 6, t_{2,1} = 1, t_{2,2} = 3, t_{2,7} = 2$.

Then the corresponding edge labels are $x_{2,1} = 4, x_{2,2} = 5, x_{2,7} = 4$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{2,0} = 6$.

Case-(g): $t_{2,0} = 5$.

If $t_{2,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then,

$$x_{2,j} = f^*(v_{2,0}v_{2,j}) = \left(\frac{5+2n}{2}\right) = 3+n = \left(\frac{5+2n+1}{2}\right) = f^*(v_{2,0}v_{2,k}) = x_{2,k}$$

Therefore, the possibilities for the pendent vertices are (1), (2 or 3), (4), (6 or 7), (8 or 9), (10 or 11). These six labels are not sufficient to labels seven vertices, $t_{2,j}$ for $1 \leq j \leq 7$.

Suppose $t_{2,7}$ takes any of the remaining values.

Let $t_{2,0} = 5, t_{2,1} = 1, t_{2,2} = 3, t_{2,7} = 2$.

Then the corresponding edge labels are $x_{2,1} = 3, x_{2,2} = 4, x_{2,7} = 4$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{2,0} = 5$.

Case-(h): $t_{2,0} = 4$.

If $t_{2,j} = 2n - 1$ and $t_{2,k} = 2n$ for some n, j and k then,

$$x_{2,j} = f^*(v_{2,0}v_{2,j}) = \left(\frac{4 + 2n - 1}{2}\right) = 2 + n = \left(\frac{4 + 2n}{2}\right) = f^*(v_{2,0}v_{2,k}) = x_{2,k}$$

Therefore, the possibilities for the pendent vertices are (1 or 2), (3), (5 or 6), (7 or 8), (9 or 10), (11). These six labels are not sufficient to label seven vertices, $t_{2,j}$ for $1 \leq j \leq 7$.

Suppose $t_{2,7}$ takes any of the remaining values.

Let $t_{2,0} = 4, t_{2,1} = 2, t_{2,2} = 3, t_{2,7} = 1$.

Then the corresponding edge labels are $x_{2,1} = 3, x_{2,2} = 4, x_{2,7} = 3$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{2,0} = 4$.

Case-(i): $t_{2,0} = 3$.

If $t_{2,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then

$$x_{2,j} = f^*(v_{2,0}v_{2,j}) = \left(\frac{3 + 2n}{2}\right) = 4 + n = \left(\frac{3 + 2n + 1}{2}\right) = f^*(v_{2,0}v_{2,k}) = x_{2,k}$$

Therefore, the possibilities for the pendent vertices are (1), (2), (4 or 5), (6 or 7), (8 or 9), (10 or 11). These six labels are not sufficient to label seven vertices, $t_{2,j}$ for $1 \leq j \leq 7$.

Suppose $t_{2,7}$ takes any of the remaining values.

Let $t_{2,0} = 3, t_{2,3} = 4, t_{2,4} = 6, t_{2,7} = 5$.

Then the corresponding edge labels are $x_{2,3} = 4, x_{2,4} = 5, x_{2,7} = 4$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{2,0} = 3$.

Case-(j): $t_{2,0} = 2$.

If $t_{2,j} = 2n - 1$ and $t_{2,k} = 2n$ for some n, j and k then,

$$x_{2,j} = f^*(v_{2,0}v_{2,j}) = \left(\frac{2 + 2n - 1}{2}\right) = 1 + n = \left(\frac{2 + 2n}{2}\right) = f^*(v_{2,0}v_{2,k}) = x_{2,k}$$

Therefore, the possibilities for the pendent vertices are (1), (3 or 4), (5 or 6), (7 or 8), (9 or 10), (11). These six labels are not sufficient to label even vertices, $t_{2,j}$ for $1 \leq j \leq 7$.

Suppose $t_{2,7}$ takes any of the remaining values.

Let $t_{2,0} = 2, t_{2,2} = 3, t_{2,3} = 5, t_{2,7} = 4$.

Then the corresponding edge labels are $x_{2,2} = 3, x_{2,3} = 4, x_{2,7} = 3$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{2,0} = 2$.

Case (k): $t_{2,0} = 1$.

If $t_{2,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{3+2n}{2}\right) = 2+n = \left(\frac{3+2n+1}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (2 or 3), (4 or 5), (6 or 7), (8 or 9), (10 or 11). These five labels are not sufficient to label seven vertices, $t_{2,j}$ for $1 \leq j \leq 7$.

Suppose $t_{2,7}$ takes any of the remaining values.

Let $t_{2,0} = 1, t_{2,1} = 2, t_{2,2} = 4, t_{2,7} = 3$.

Then the corresponding edge labels are $x_{2,1} = 2, x_{2,2} = 3, x_{2,7} = 2$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{2,0} = 1$.

\Rightarrow G is not a relaxed skolem mean graph for all values of $t_{2,0}$.

Therefore G is not a relaxed skolem mean graph.

In general, we see that G is not a relaxed skolem mean graph if $|m - n| = 6$.

Similar argument asserts that $|m - n| = 7$ is not a relaxed skolem mean graph.

Similarly, we can prove for all greater values of m.

Therefore, G is not a relaxed skolem mean graph if $|m - n| > 5$.

Definition 2.4: The three star is the disjoint union of $K_{1,\ell}, K_{1,m}$ and $K_{1,n}$. It is denoted by $K_{1,\ell} \cup K_{1,m} \cup K_{1,n}$.

Theorem 2.5: The three star $K_{1,\ell} \cup K_{1,m} \cup K_{1,n}$ is not a relaxed Skolem mean graph if $|m - n| > \ell + 6$.

Proof: Let us consider the primal graph under the condition $|m - n| > \ell + 6$ and $m = 1; n = 9$.

Therefore $G = K_{1,1} \cup K_{1,1} \cup K_{1,9} = 2K_{1,1} \cup K_{1,9}$

Where $V(G) = \{v_{i,j} : 1 \leq i \leq 2, 0 \leq j \leq 1\} \cup \{v_{3,j} : 0 \leq j \leq 9\}$

$$E(G) = \{v_{i,0}v_{i,1} : 1 \leq i \leq 2\} \cup \{v_{3,0}v_{3,j} : 1 \leq j \leq 9\}$$

G has 14 vertices and 11 edges.

Suppose G is relaxed skolem mean graph.

Then there exists a function f from the vertex set of G to $\{1,2,\dots,15\}$ such that the induced map f^* from the edge set of G to $\{2, 3,\dots,15\}$ defined by

$$f^*(e = uv) = \begin{cases} \frac{f(u) + f(v)}{2} & \text{if } f(u) + f(v) \text{ is even} \\ \frac{f(u) + f(v) + 1}{2} & \text{if } f(u) + f(v) \text{ is odd} \end{cases}$$

Then the resulting edges get distinct label from the set $\{2, 3, \dots, 15\}$.

Then the vertex and edge mappings of G is given by

$$f: V(G) \rightarrow \{1, 2, \dots, 15\}$$

$$f^*: V(G) \rightarrow \{2, 3, \dots, 15\}$$

Let $t_{i,j} = f(v_{i,j})$ and $x_{i,j} = f^*(v_{i,0} v_{i,j}) \forall i$ and j .

Now let us consider the following cases,

Case (a): $t_{3,0} = 15$.

If $t_{3,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0} v_{3,j}) = \left(\frac{15 + 2n}{2} \right) = 8 + n = \left(\frac{15 + 2n + 1}{2} \right) = f^*(v_{3,0} v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1), (2 or 3), (4 or 5), (6 or 7), (8 or 9), (10 or 11), (12 or 13), 14. These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 15, t_{3,1} = 1, t_{3,2} = 3, t_{3,9} = 2$.

Then the corresponding edge labels are $x_{3,1} = 8, x_{3,2} = 9, x_{3,9} = 9$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{3,0} = 15$.

Case (b): $t_{3,0} = 14$.

If $t_{3,j} = n - 1$ and $t_{2,k} = 2n$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0} v_{3,j}) = \left(\frac{14 + 2n - 1}{2} \right) = 7 + n = \left(\frac{14 + 2n}{2} \right) = f^*(v_{3,0} v_{3,k}) = x_{3,k}$$

Therefore the possibilities for the pendent vertices are (1 or 2), (3 or 4), (5 or 6), (7 or 8), (9 or 10), (11 or 12), (13), (15). These eight labels are not sufficient to label nine vertices $t_{3,9}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 14, t_{3,1} = 2, t_{3,2} = 4, t_{3,9} = 3$.

Then the corresponding edge labels are $x_{3,1} = 8, x_{3,2} = 9, x_{3,9} = 9$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{3,0} = 14$.

Case-(c): $t_{3,0} = 13$.

If $t_{3,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0} v_{3,j}) = \left(\frac{13 + 2n}{2} \right) = 7 + n = \left(\frac{13 + 2n + 1}{2} \right) = f^*(v_{3,0} v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1), (2 or 3), (4 or 5), (6 or 7), (8 or 9), (10 or 11), (12 or 13), 14. These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 13$, $t_{3,1} = 1$, $t_{3,2} = 3$, $t_{3,9} = 2$.

Then the corresponding edge labels are $x_{3,1} = 7$, $x_{3,2} = 8$, $x_{3,9} = 8$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore G is not a relaxed skolem mean graph when $t_{3,0} = 13$.

Case (d): $t_{3,0} = 12$.

If $t_{3,j} = 2n - 1$ and $t_{2,k} = 2n$ for some n, j and k then

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{12 + 2n - 1}{2}\right) = 6 + n = \left(\frac{12 + 2n}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1 or 2), (3 or 4), (5 or 6), (7 or 8), (9 or 10), (11), (13 or 14), (15). These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 12$, $t_{3,8} = 15$, $t_{3,7} = 13$, $t_{3,9} = 14$.

Then the corresponding edge labels are $x_{3,8} = 14$, $x_{3,7} = 13$, $x_{3,9} = 13$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore G is not a relaxed skolem mean graph when $t_{3,0} = 12$.

Case-(e): $t_{3,0} = 11$.

If $t_{3,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{11 + 2n}{2}\right) = 6 + n = \left(\frac{11 + 2n + 1}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}.$$

Therefore, the possibilities for the pendent vertices are (1), (2 or 3), (4 or 5), (6 or 7), (8 or 9), (10), (12 or 13), (14 or 15). These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 11$, $t_{3,1} = 1$, $t_{3,2} = 3$, $t_{3,9} = 2$.

Then the corresponding edge labels are $x_{3,1} = 6$, $x_{3,2} = 7$, $x_{3,9} = 7$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore G is not a relaxed skolem mean graph when $t_{3,0} = 11$.

Case-(f): $t_{3,0} = 10$.

If $t_{3,j} = 2n - 1$ and $t_{2,k} = 2n$ for some n, j and k then

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{10+2n-1}{2}\right) = 5+n = \left(\frac{10+2n}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1 or 2), (3 or 4), (5 or 6), (7 or 8), 9, (11 or 12), (13 or 14), (15). These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 10, t_{3,1} = 2, t_{3,2} = 4, t_{3,9} = 3$.

Then the corresponding edge labels are $x_{3,1} = 6, x_{3,2} = 7, x_{3,9} = 7$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore G is not a relaxed skolem mean graph when $t_{3,0} = 10$.

Case-(g): $t_{3,0} = 9$

If $t_{3,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{9+2n}{2}\right) = 5+n = \left(\frac{9+2n+1}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1), (2 or 3), (4 or 5), (6 or 7), (8), (10 or 11), (12 or 13), (14 or 15). These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 9, t_{3,1} = 1, t_{3,2} = 3, t_{3,9} = 2$.

Then the corresponding edge labels are $x_{3,1} = 5, x_{3,2} = 6, x_{3,9} = 6$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore G is not a relaxed skolem mean graph when $t_{3,0} = 9$.

Case-(h): $t_{3,0} = 8$.

If $t_{3,j} = 2n - 1$ and $t_{2,k} = 2n$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{8+2n-1}{2}\right) = 4+n = \left(\frac{8+2n}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1 or 2), (3 or 4), (5 or 6), 7, (9 or 10), (11 or 12), (13 or 14), (15). These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 8, t_{3,1} = 2, t_{3,2} = 4, t_{3,9} = 3$.

Then the corresponding edge labels are $x_{3,1} = 5, x_{3,2} = 6, x_{3,9} = 6$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{3,0} = 8$.

Case-(i): $t_{3,0} = 7$.

If $t_{3,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{7 + 2n}{2}\right) = 4 + n = \left(\frac{7 + 2n + 1}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1), (2 or 3), (4 or 5), 6, (8 or 9), (10 or 11), (12 or 13), (14 or 15). These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 7, t_{3,1} = 1, t_{3,2} = 3, t_{3,9} = 2$.

Then the corresponding edge labels are $x_{3,1} = 4, x_{3,2} = 5, x_{3,9} = 5$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{3,0} = 7$.

Case-(j): $t_{3,0} = 6$.

If $t_{3,j} = 2n - 1$ and $t_{2,k} = 2n$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{6 + 2n - 1}{2}\right) = 3 + n = \left(\frac{6 + 2n}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1 or 2), (3 or 4), (5), (7 or 8), (9 or 10), (11 or 12), (13 or 14), (15). These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 6, t_{3,2} = 3, t_{3,3} = 5, t_{3,9} = 4$.

Then the corresponding edge labels are $x_{3,2} = 5, x_{3,3} = 6, x_{3,9} = 5$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{3,0} = 6$.

Case-(k): $t_{3,0} = 5$.

If $t_{3,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{5 + 2n}{2}\right) = 3 + n = \left(\frac{5 + 2n + 1}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1), (2 or 3), 4, (6 or 7), (8 or 9), (10 or 11), (12 or 13), (14 or 15). These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 5, t_{3,1} = 1, t_{3,2} = 3, t_{3,9} = 2$.

Then the corresponding edge labels are $x_{3,1} = 3, x_{3,2} = 4, x_{3,9} = 4$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore G is not a relaxed skolem mean graph when $t_{3,0} = 5$.

Case-(l): $t_{3,0} = 4$.

If $t_{3,j} = 2n - 1$ and $t_{2,k} = 2n$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{4 + 2n - 1}{2}\right) = 2 + n = \left(\frac{4 + 2n}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1 or 2), (3), (5 or 6), (7 or 8), (9 or 10), (11 or 12), (13 or 14), (15). These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 4, t_{3,2} = 3, t_{3,3} = 5, t_{3,9} = 6$.

Then the corresponding edge labels are $x_{3,2} = 4, x_{3,3} = 5, x_{3,9} = 5$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore G is not a relaxed skolem mean graph when $t_{3,0} = 4$.

Case-(m): $t_{3,0} = 3$.

If $t_{3,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{3 + 2n}{2}\right) = 2 + n = \left(\frac{3 + 2n + 1}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1), (2), (4 or 5), (6 or 7), (8 or 9), (10 or 11), (12 or 13), (14 or 15). These eight labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 3, t_{3,3} = 4, t_{3,4} = 6, t_{3,9} = 5$.

Then the corresponding edge labels are $x_{3,3} = 4, x_{3,4} = 5, x_{3,9} = 4$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{3,0} = 3$.

Case-(n): $t_{3,0} = 2$.

If $t_{3,j} = 2n + 1$ and $t_{2,k} = 2n + 2$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{2 + 2n + 1}{2}\right) = 2 + n = \left(\frac{2 + 2n + 2}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (1), (3 or 4), (5 or 6), (7 or 8), (9 or 10), (11 or 12), (13 or 14), (15). These eight labels are not sufficient to label nine vertices $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 2, t_{3,2} = 3, t_{3,3} = 5, t_{3,9} = 4$.

Then the corresponding edge labels are $x_{3,2} = 3, x_{3,3} = 4, x_{3,9} = 3$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore, G is not a relaxed skolem mean graph when $t_{3,0} = 2$.

Case-(o): $t_{3,0} = 1$.

If $t_{3,j} = 2n$ and $t_{2,k} = 2n + 1$ for some n, j and k then,

$$x_{3,j} = f^*(v_{3,0}v_{3,j}) = \left(\frac{1+2n}{2}\right) = 1+n = \left(\frac{1+2n+1}{2}\right) = f^*(v_{3,0}v_{3,k}) = x_{3,k}$$

Therefore, the possibilities for the pendent vertices are (2 or 3), (4 or 5), (6 or 7), (8 or 9), (10 or 11), (12 or 13), (14 or 15). These seven labels are not sufficient to label nine vertices, $t_{3,j}$ for $1 \leq j \leq 9$.

Suppose $t_{3,9}$ takes any of the remaining values.

Let $t_{3,0} = 1, t_{3,1} = 3, t_{3,2} = 5, t_{3,8} = 2, t_{3,9} = 4$.

Then the corresponding edge labels are $x_{3,1} = 2, x_{3,2} = 3, x_{3,8} = 4, x_{3,9} = 3$. In this case it is not possible to label the vertices without labeling three of them as consecutive integers. If there are three consecutive integers then two of them will induce the same edge label.

Therefore G is not a relaxed skolem mean graph when $t_{3,0} = 1$.

G is not a relaxed skolem mean graph for all values of $t_{3,0}$.

Therefore, G is not a relaxed skolem mean graph.

Similarly $G = K_{1,2} \cup K_{1,2} \cup K_{1,11}$ is not a relaxed skolem mean graph.

In general we see that G is not a relaxed skolem mean graph if $|m - n| = \ell + 7$. With similar argument we can prove for all greater values

Therefore, G is not a relaxed skolem mean graph if $|m - n| > \ell + 6$.

3. APPLICATION OF GRAPH LABELING

The skolem mean labeling is applied on a graph (network), such as bus topology, mesh topology and star topology in order to solve the problems in establishing fastness, efficient communication and various issues in that area, in which the following will be taken into account.

1. A protocol, with secured communication can be achieved, provided the graph (network) is sufficiently connected.
2. To find an efficient way for safer transmissions in areas such as Cellular telephony, Wi – Fi, Security systems and many more.
3. Channel labeling can be used to determine the time at which sensor communicate.

CONCLUSION

Researchers may get some information related to graph labeling and its applications in communication field and work on some ideas related to their field of research.

ACKNOWLEDGEMENT

One of the authors (Dr. V. Balaji) acknowledges University Grants Commission, SERO, Hyderabad, India for financial assistance (No. F MRP 5766 / 15 (SERO / UGC)).

4. REFERENCES

1. Abraham K. Samuel, D.S.T. Ramesh, M. Elakiya and V. Balaji, *on Relaxed Skolem Mean Labeling for four star*, International Journal of Mathematics and its application vol.4,2016,17 – 23.
2. P. Alayamani, D.S.T. Ramesh, V. Balaji and M. Elakiya, *on Relaxed Skolem Mean Labeling for four star*. IOSR Journal of Mathematics vol.12, 2016, 19 – 26.
3. V. Blaji, D.S.T. Ramesh and A. Subramanian, *Some results on Relaxed Skolem Mean Graphs*, Bulletin of kerala mathematics Association vol.5 (2). Dec 2009, 33 – 44.
4. V. Blaji, D.S.T. Ramesh and A. Subramanian *Relaxed Skolem Mean Labeling, advances and applications in Discrete Mathematics*, vol.5 (1), Jan 2010, 11 – 22.
5. V. Blaji, D.S.T. Ramesh and K. Valarmathy, *on Relaxed Skolem Mean Labeling for three star*, International Journal of Mathematical Archieve, vol.7(2), 2016, 1 – 7.
6. J.A. Bondy, *Introduction to Analytic Number Theory*, Narosa publishing House, second edition 1991.
7. J.A. Gallian, *A Dynamic survey of Graph Labeling*, The Electronic Journal of combinatorics 14 (2007), #DS6.
8. F. Harary, *Graph Theory*, Addison – Wesley Reading Mars (1972).
9. V. Maheswari, D.S.T. Ramesh, Silviya Francis and V. Balaji, *Non Existence of Relaxed Mean Labeling for subdivision of star graphs*. Bulletin of Kerala Mathematics Association vol.13, No.1, June 2016, 57 – 73.
10. Rosa, *On certain valuation of the vertices of a graph, Theory of Graphs(International symposium, Rome, July 1966)* Gardon and Breach, N.Y and Dunod, paris (1967), 349 – 355.

**Source of support: University Grants Commission, SERO, Hyderabad, India,
Conflict of interest: None Declared.**

[Copy right © 2017. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]