

SOME FIXED POINT THEOREMS
FOR EXTENDED EXPANSION MAPPINGS ON CONE METRIC SPACES

SHERLY GEORGE*, SHAINI PULICKAKUNNEL

Department of Mathematics,
Sam Higginbottom University of Agriculture,
Technology and Sciences, Allahabad-211007, India.

(Received On: 08-07-17; Revised & Accepted On: 24-09-17)

ABSTRACT

In this paper, we prove some fixed point theorems for expansion mappings in the framework of cone metric spaces. Our results in this paper extends and improves upon, among others, the corresponding results of Aage and Salunke [1]

MSC: primary 47H09, 47H10, 54H25.

Keywords: Cone metric space, normal constants, fixed point.

1. INTRODUCTION

One of the most widely used fixed point theorems in all analysis is Banach contraction theorem. It has been generalized in different directions by Mathematicians over the years. In contemporary time, fixed point theory has evolved speedily in cone metric spaces equipped with partial ordering. Huang and Zhang introduced in [5] the concept of cone metric space as generalisation of metric space where the set of real numbers is replaced by an ordered Banach space. Thereafter various authors have generalised the results of Huang and Zhang and studied fixed point theorems for normal and non-normal cones [2, 3, 6, 11]. A new generalisation of contraction mappings called T-contraction mappings was introduced by Beiranvand on metric spaces [4]. Recently Morales and Rojas [7], [8], [9], [10] have extended the concept of T-contraction mappings to the cone metric spaces by proving fixed point theorems for T-Kannan, T-Zamfirescu, T-Weakly contraction mappings.

Motivated by that we generalize the theorems given in [1] by extending the expansive mappings.

The following definitions and results will be needed in the sequel.

Definition 1: Let E be a real Banach space. A subset P of E is called a cone if and only if

1. P is closed, nonempty and $P \neq \{0\}$;
2. $a, b \in \mathbb{R}, a, b \geq 0, x, y \in P$ imply that $ax + by \in P$;
3. $P \cap (-P) = \{0\}$.

Given a cone $P \subset E$, we define a partial ordering \leq with respect to P by $x \leq y$ if and only if $y - x \in P$. A cone P is called normal if there is a number $K > 0$ such that for all $x, y \in E$,

$$0 \leq x \leq y \text{ implies } \|x\| \leq K\|y\| \quad (1)$$

The least positive number satisfying the above inequality is called the normal constant of P . We shall write $x < y$ to indicate that $x \leq y$ but $x \neq y$, while $x \ll y$ stands for $y - x \in \text{int } P$ (interior of P).

In the following we always suppose that E is a Banach space, P is a cone in E with $\text{int } P \neq \emptyset$ and \leq is partial ordering with respect to P .

Corresponding Author: Sherly George*,
Department of Mathematics, Sam Higginbottom University of Agriculture,
Technology and Sciences, Allahabad-211007, India.

Definition 2: Let X be a nonempty set. Suppose that the mapping $d : X \times X \rightarrow E$ satisfies:

1. $0 \leq d(x,y)$ for all $x,y \in X$ and $d(x,y)=0$ if and only $x=y$;
2. $d(x,y)=d(y,x)$ for all $x,y \in X$;
3. $d(x,y) \leq d(x,z)+d(z,y)$ for all $x,y,z \in X$.

Then d is called a cone metric on X and (X,d) is called a cone metric space. The concept of cone metric space is more general than that of metric space.

Definition 3: Let (X,d) be a cone metric space, $\{x_n\}$ a sequence in X and $x \in X$. For every $c \in E$ with $0 \ll c$, we say that $\{x_n\}$ is :

1. a Cauchy sequence if there is an N such that, for $n, m, > N$, $d(x_n, x_m) \ll c$
2. a convergent sequence if there is an N such that, for all $n, > N$, $d(x_n, x) \ll c$ for some x in X

A cone metric space X is said to be complete if every Cauchy sequence in X is convergent in X . It is known that $\{x_n\}$ converges to $x \in X$ if and only if $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$. The limit of a convergent sequence is unique provided P is a normal cone with normal constant K [5].

Definition 4: Let (M,d) be a cone metric space, P be a normal cone with normal constant K and $T : M \rightarrow M$. Then

1. T is said to be continuous if $\lim_{n \rightarrow \infty} x_n = x$, implies that $\lim_{n \rightarrow \infty} Tx_n = Tx$ for every $\{x_n\}$ in M ;
2. T is said to be sequentially convergent if we have, for every sequence $\{x_n\}$, if $\{Tx_n\}$ is convergent, then $\{x_n\}$ is also convergent;
3. T is said to be subsequentially convergent if we have, for every sequence $\{x_n\}$, if $\{Tx_n\}$ is convergent, then $\{x_n\}$ has convergent subsequence.

Definition 5: Let (X,d) be a cone metric space and $T, S : X \rightarrow X$ be two functions. S is said to be T - contraction if there exists $a \in [0,1)$ such that

$$d(TSx, TSy) \leq ad(Tx, Ty), \forall x,y \in X.$$

The following two lemmas of Huang and Zhang [5] will be required in the sequel.

Lemma 1 [5, lemma 1]: Let (M,d) be a cone metric space, P be a normal cone with normal constant K . A sequence $\{x_n\}$ converges to $x \in X$ if and only if $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.

Lemma 2 [5, lemma 4]: Let (M,d) be a cone metric space, P be a normal cone with normal constant K . A sequence $\{x_n\}$ in X is Cauchy sequence if and only if $d(x_n, x_m) \rightarrow 0$ as $n, m \rightarrow \infty$

2. MAIN RESULTS

Theorem 1: Let (X,d) be a complete cone metric space, P be a normal cone with normal constant K and two continuous mappings $T, f : X \rightarrow X$ satisfies the contractive condition

$$d(fTx, fTy) \geq kd(fx, fy)$$

for all $x,y \in X$, where $k > 1$ is a constant. Assume that T is onto, f is one-one, subsequentially convergent, and T, f are commutative. Then T has a unique fixed point in X .

Proof: If $fT(x) = fT(y)$, then

$$0 \geq kd(f(x), f(y)) \Rightarrow 0 = d(f(x), f(y))$$

$$\text{i.e. } fT(x) = fT(y) \Rightarrow fx = fy$$

Since the maps commute, we get

$$Tf(x) = Tf(y) \Rightarrow f(x) = f(y).$$

Thus T is one-one. Define $G = T^{-1}$

$$\begin{aligned} d(f(x), f(y)) &= d(fTT^{-1}x, fTT^{-1}y) \\ &\geq kd(fT^{-1}x, fT^{-1}y) \\ &\geq kd(fGx, fG(y)). \end{aligned}$$

$$d(fG(x), fGy) \leq \frac{1}{k} d(fx, fy) \\ \leq Md(fx, fy) \text{ where } M = \frac{1}{k} \leq 1$$

From definition 5 and by Theorem 3.3 in [7], G has a unique fixed point u in X .

$$\text{i.e., } G(u) = u \Rightarrow T^{-1}u = u, \\ u = Tu.$$

$\Rightarrow T$ has a unique fixed point

When $T=I$ we get the following theorem

Corollary 1: [1, Theorem 2.1] Let (X,d) be a complete cone metric space and the mapping $T : X \rightarrow X$ is onto and satisfies the contractive condition

$$d(Tx, Ty) \geq kd(x, y),$$

for all $x, y \in X$, where $k > 1$ is a constant. Then T has a unique fixed point in X .

Theorem 2: Let (X,d) be a complete cone metric space, P be a normal cone with normal constant K and two commutative continuous mappings $f, T : X \rightarrow X$ satisfy the condition

$$d(fTx, fTy) \geq k[d(fTx, fx) + d(fTy, fy)],$$

for all $x, y \in X$ where $\frac{1}{2} < k \leq 1$. Assume that f is subsequentially convergent, injective and T is onto. Then T has fixed point in X .

Proof: Since T is onto, for each $x_0 \in X$, there exist $x_1 \in X$ such that $Tx_1 = x_0$. Similarly for each $n \geq 1$ there exist $x_{n+1} \in X$, such that

$x_n = Tx_{n+1}$. If $x_{n-1} = x_n$, then x_n is a fixed point of T . Thus, Suppose that $x_{n-1} \neq x_n$ for all $n \geq 1$. Then

$$d(fx_n, fx_{n-1}) = d(fTx_{n+1}, fTx_n) \\ \geq k[d(fTx_{n+1}, fx_{n+1}) + d(fTx_n, fx_n)] \\ \geq kd(fx_n, fx_{n+1}) + d(fx_{n-1}, fx_n). \\ d(fx_n, fx_{n+1}) \leq \frac{1-k}{k} d(fx_{n-1}, fx_n) = hd(fx_{n-1}, fx_n).$$

From this we get

$$d(fx_n, fx_{n+1}) \leq h^n d(fx_0, fx_1) \quad \text{where } h = \frac{1-k}{k} \text{ and } 0 \leq h < 1$$

Now, for $n < m$

$$d(fx_n, fx_m) \leq d(fx_n, fx_{n+1}) + d(fx_{n+1}, fx_{n+2}) + \dots + d(fx_{m-1}, fx_m) \\ \leq (h^n + h^{n+1} + \dots + h^{m-1})d(fx_0, fx_1) \\ \leq \frac{h^n}{1-h} d(fx_0, fx_1).$$

From (1) we get

$$\|d(fx_n, fx_m)\| \leq K \frac{h^n}{1-h} \|d(fx_0, fx_1)\| \rightarrow 0 \text{ as } n \rightarrow \infty$$

$\{fx_n\}$ is Cauchy sequence. Since (X,d) be complete cone metric space there exist $u \in X$ such that $\lim_{n \rightarrow \infty} fx_n = u$.

Since f is subsequentially convergent $\{x_n\}$ has a convergent subsequence $\{x_m\}$

$$\text{i.e., } \lim_{m \rightarrow \infty} x_m = x^*.$$

Since f is continuous

$$\lim_{m \rightarrow \infty} f(x_m) = fx^*.$$

By uniqueness of limit,

$$u = fx^*,$$

$$i.e., \lim_{n \rightarrow \infty} fx_n = fx^*$$

Since T is continuous

$$\lim_{n \rightarrow \infty} Tfx_n = Tfx^*.$$

$$d(fTx^*, fTx_n) \leq d(fTx^*, fTx_n) + d(fTx_n, fx^*) = 0 \text{ as } n \rightarrow \infty$$

Then we get

$$d(fTx^*, fx^*) = 0$$

Which implies that

$$fTx^* = fx^*.$$

Since f is injective, $Tx^* = x^*$ Thus T has a fixed point in X .

When $f=I$ we get the following result as corollary.

Corollary 2: [1, Theorem 2.3] Let (X, d) be a complete cone metric space and continuous, onto mapping $T : X \rightarrow \infty$ satisfies the condition

$$d(Tx, Ty) \geq kd[d(Tx, x) + d(Ty, y)],$$

for all $x, y \in X$ where $\frac{1}{2} < k \leq 1$. Then T has fixed point in X .

Theorem 3: Let (X, d) be a complete cone metric space with normal constant K and two commutative continuous mappings f, T satisfies the contractive condition

$$d(fTx, fTy) \geq kd(fx, fy) + ld(fTx, fy),$$

for all $x, y \in X$, where $l \geq 0, k > 1$. Assume that T is onto, f is injective and subsequentially convergent. Then T has a fixed point in X .

Proof: Since T is onto, for each $x_0 \in X$ there exist $x_1 \in X$ such that $Tx_1 = x_0$, Similarly for each $n \geq 1$ there exist $x_{n+1} \in X$ such that

$x_n = Tx_{n+1}$. If $x_{n-1} = x_n$, then x_n is a fixed point of T . Suppose that $x_{n-1} \neq x_n$ for all $n \geq 1$.

$$\begin{aligned} d(fx_n, fx_{n-1}) &= d(fTx_{n+1}, fTx_n) \\ &\geq kd(fx_{n+1}, fx_n) + ld(fTx_{n+1}, fx_n) \\ &= kd(fx_{n+1}, fx_n) + ld(fx_n, fx_n) \\ &\leq kd(fx_n, fx_{n+1}). \end{aligned}$$

$$d(fx_n, fx_{n+1}) \leq \frac{1}{k} d(fx_{n-1}, fx_n).$$

$$d(fx_n, fx_{n+1}) \leq hd(fx_{n-1}, fx_n) \text{ where } h = \frac{1}{k} < 1.$$

$$\begin{aligned} &\leq h^2 d(fx_{n-2}, fx_{n-1}) \\ &\dots\dots\dots \\ &\leq h^n d(fx_0, fx_1). \end{aligned}$$

Consider $n < m$

$$\begin{aligned} d(fx_n, fx_m) &\leq d(fx_n, fx_{n+1}) + d(fx_{n+1}, fx_{n+2}) + \dots + d(fx_{m-1}, fx_m) \\ &\leq (h^n + h^{n+1} + \dots + h^{m-1})d(fx_0, fx_1) \\ &\leq \frac{h^n}{1-h} d(fx_0, fx_1). \end{aligned}$$

From (1) we get

$$\|d(fx_n, fx_m)\| \leq K \frac{h^n}{1-h} \|d(fx_0, fx_1)\| \rightarrow 0 \text{ as } n \rightarrow \infty$$

This shows that $\{fx_n\}_{n=1}$ is Cauchy sequence.

Since (X, d) complete cone metric space,

$$\lim_{n \rightarrow \infty} fx_n = u.$$

Since f is subsequentially convergent, $\{x_n\}$ has a convergent subsequence $\{x_m\}$. i.e., $\lim_{m \rightarrow \infty} x_m = x^*$

Since f is continuous,

$$\lim_{m \rightarrow \infty} fx_m = fx^*.$$

By uniqueness of limit

$$\begin{aligned} u &= fx^*. \\ \text{i.e., } \lim_{n \rightarrow \infty} fx_n &= fx^*. \end{aligned}$$

Since T is continuous,

$$\lim_{n \rightarrow \infty} Tfx_n = Tfx^*.$$

Now,
$$d(fTx^*, fx^*) \leq d(fTx^*, fTx_n) + d(fTx_n, fx^*)$$

Which implies that

$$d(fTx^*, fx^*) = 0 \text{ as } n \rightarrow \infty$$

$$\text{Hence, } fTx^* = fx^*$$

Since f is injective, $Tx^* = x^*$. Thus T has a fixed point in X .

When $f=I$ we get the following result as corollary.

Corollary 3: [1, Theorem 2.5] Let (X, d) be a complete cone metric space and continuous, onto mapping T satisfies the contractive condition

$$d(Tx, Ty) \geq kd(x, y) + ld(Tx, y),$$

for all $x, y \in X$, where $l \geq 0, k > 0$. Then T has a fixed point in X .

Theorem 4: Let (X, d) be a complete cone metric space with normal constant K and two continuous commutative mappings $T, f : X \rightarrow X$ satisfies the condition

$$d(fTx, fTy) \geq kd(fx, fy) + ld(fx, fTx) + pfy, fTy),$$

for all $x, y \in X$ where $k \geq -1, p < 1$ and $l > 1$ with $k+l+p > 1$. Assume that T is onto and f is injective and subsequentially convergent. Then T has a fixed point in X .

Proof: Since T is onto, for each $x_0 \in X$, there exist $x_1 \in X$ such that $Tx_1 = x_0$. Similarly, for each $n \geq 1$ there exist $x_{n+1} \in X$, such that

$x_n = Tx_{n+1}$. If $x_{n-1} = x_n$, then x_n is a fixed point of T . Thus, suppose that $x_{n-1} \neq x_n$ for all $n \geq 1$. Then

$$\begin{aligned} d(fx_n, fx_{n-1}) &= d(fTx_{n+1}, fTx_n) \\ &\geq kd(fx_{n+1}, fx_n) + ld(fx_{n+1}, fTx_{n+1}) + pd(fx_n, fTx_n) \\ &= kd(fx_{n+1}, fx_n) + ld(fx_{n+1}, fx_n) + pd(fx_n, fx_{n-1}), \end{aligned}$$

$$\begin{aligned} d(fx_{n+1}, fx_n) &\leq \frac{1-p}{k+l} d(fx_n, fx_{n-1}) \\ &= hd(fx_n, fx_{n-1}) \text{ where } h = \frac{1-p}{k+l} \text{ and } 0 < h < 1 \\ &\leq h^2 d(fx_{n-1}, fx_{n-2}) \\ &\dots\dots\dots \\ &\dots\dots\dots \\ &\leq h^n d(fx_1, x_0). \end{aligned}$$

From this we get

$$d(fx_n, fx_{n+1}) \leq h^n d(fx_0, fx_1) \text{ where } h = \frac{1-p}{k+l} \text{ and } 0 < h < 1.$$

Consider $m > n$

$$\begin{aligned} d(fx_n, fx_m) &\leq h^n d(fx_0, fx_1) \text{ where } h = \frac{1-p}{k+l} \text{ and } 0 < h < 1. \\ &\leq d(fx_n, fx_{n+1}) + d(fx_{n+1}, fx_{n+2}) + \dots + d(fx_{m-1}, fx_m) \\ &\leq (h^n + h^{n+1} + \dots + h^{m-1}) d(fx_1, fx_0). \\ &\leq \frac{h^n}{1-h} d(fx_1, fx_0) \end{aligned}$$

From (1), we get

$$\|d(fx_n, fx_m)\| \leq K \frac{h^n}{1-h} \|d(fx_0, fx_1)\| \rightarrow 0 \text{ as } n \rightarrow \infty$$

Hence $\|d(fx_n, fx_m)\| \rightarrow 0$

Thus $\{fx_n\}_{n=1}$ is a Cauchy sequence.

Since (X, d) is a complete cone metric space, $\lim_{n \rightarrow \infty} fx_n = u$.

Since f is subsequentially convergent $\{x_n\}$ has a convergent subsequence $\{x_m\}$ i.e., $\lim_{m \rightarrow \infty} x_m = x^*$

Since f is continuous,

$$\lim_{m \rightarrow \infty} fx_m = fx^*,$$

By uniqueness of limit

$$\begin{aligned} u &= fx^*. \\ \text{i.e., } \lim_{n \rightarrow \infty} fx_n &= fx^* \end{aligned}$$

Since T is continuous

$$\lim_{n \rightarrow \infty} Tfx_n = Tfx^*$$

Now
$$d(fTx^*, fx^*) \leq d(fTx^*, Tfx_n) + d(fTx_n, fx^*)$$

Which implies that

$$d(fTx^*, fx^*) = 0 \text{ as } n \rightarrow \infty$$

Since f is injective, $Tx^* = x^*$. Then T has a fixed point.

When $f=I$ we get the following result as corollary.

Corollary 4: [1, Theorem 2.6] *Let (X, d) be complete cone metric space and continuous, onto mapping $T : X \rightarrow X$ satisfies the condition*

$$d(Tx, Ty) \geq kd(x, y) + ld(x, Tx) + pd(y, Ty),$$

for all $x, y \in X$ where $k \geq -1$, $p < 1$ and $l >$ with $k+l+p > 1$. Then T has a fixed point in X .

REFERENCES

- 1 C. T. Aage and J.N. Salunke *Some fixed point theorems onto mappings On cone metric spaces*, Acta Mathematica Sinica, English Series., J., 27, (2011), 1101-1106.
- 2 A. Azam, M. Arshad, I. Beg, *Banach Contraction Principle on Cone rectangular metric spaces*, Appl. Anal. Discrete Math., 3(2), (2009), 236-241.
- 3 M. Abbas, G. Jungck, *Common fixed point results for non commuting mappings without continuity in cone metric spaces*, J. Math. Anal. Appl. 341, (2008), 416-420.
- 4 A. Beiranvand, S. Moradi, M. Omid and H. Pazandeh, *Two fixed point theorems for special mappings*, arXiv:0903.1504v1 [math.FA].
- 5 L. G. Haung, X. Zhang, *Cone metric spaces and fixed point theorems of contractive mappings*, J. Math. Anal. Appl., 332, (2007), 1468-1476.
- 6 D. Ilic and V. Rakocevic, *Common fixed points maps on cone metric space*, J. Math. Anal. Appl., 341, (2008), 876-882.
- 7 J. R. Morales and E. Rojas, *Cone metric spaces and fixed point theorems of T-contractive mappings*, Revista Notas de mathematica, 5(1), 274, (2009), 64-71.
- 8 J. R. Morales and E. Rojas, *Some results on T-Zamfirescu operators*, Revista Notas de mathematica, 4(2), 269, (2009), 66-78.
- 9 J. R. Morales and E. Rojas, *T-Zamfirescu and T-weak contraction mappings on cone metric spaces*, arXiv: 0909.1255v1 [math.FA].
- 10 J. R. Morales and E. Rojas, *Cone metric spaces and fixed point theorems of T-Kannan contractive mappings*, Int. J. Math. Anal., 4(4), (2010), 175-184.
- 11 P. Raja, Vaezpour, S.M., *Some Extensions of Banach's Contraction Principle in complete cone metric spaces*, Fixed Point Theory Appl., 11, (2008) Article ID 768294.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2017. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]