

**$\delta\omega\alpha$ - CLOSED FUNCTIONS AND Quasi- $\delta\omega\alpha$ CLOSED FUNCTIONS
IN TOPOLOGICAL SPACES**

S. CHANDRASEKAR*¹, T. RAJESH KANNAN² AND R. SELVARAJ³

^{1,2}Department of Mathematics,
Arignar Anna Government Arts college, Namakkal (DT), Tamil Nadu, India.

³Department of Mathematics,
Pavai College of Technology, Namakkal (DT), Tamil Nadu, India.

(Received On: 22-09-17; Revised & Accepted On: 25-10-17)

ABSTRACT

Open and closed functions are most important concepts Mathematical Sciences. $\delta\omega\alpha$ -closed sets introduced by S.Chandrasekar, T.Rajesh Kannan *et.al.* In this paper we are introduced by $\delta\omega\alpha$ open functions, $\delta\omega\alpha$ closed functions, quasi $\delta\omega\alpha$ open functions, and quasi $\delta\omega\alpha$ closed functions in topological spaces.

Key Words: $\delta\omega\alpha$ -open functions, $\delta\omega\alpha$ -closed functions, quasi $\delta\omega\alpha$ -open functions and quasi $\delta\omega\alpha$ -closed functions

AMS subject classification: 54C55, 54A05.

1. INTRODUCTION

Velicko introduced δ -closed set in Topological spaces [11]. Using δ -closed set several results introduced by many researcher. $\omega\alpha$ - closed set[1] introduced by S.S.Benchalli, *et al.*, in the year 2009. Since the advent of these types of notions, several author have been introduced interesting results. $\delta\omega\alpha$ -closed sets [2] introduced by S.Chandrasekar, T.Rajesh Kannan *et.al.* In this paper we introduced $\delta\omega\alpha$ open functions, $\delta\omega\alpha$ closed functions, quasi $\delta\omega\alpha$ open functions, and quasi $\delta\omega\alpha$ closed functions in topological spaces and application properties are discussed detailed

2. PRELIMINARIES

Let us recall the following definition, which are useful in the sequel.

Definition 2.1: A subset A of a space (X, τ) is called

- 1) α - closed set [6] if $\text{cl}(\text{int}(\text{cl}(A))) \subseteq A$
 - 2) δ -closed [11] if $A = \text{cl}_\delta(A)$, where $\text{cl}_\delta(A) = \{x \in X: \text{int}(\text{cl}(U)) \cap A \neq \emptyset, U \in \tau \text{ and } x \in U\}$.
 - 3) ω -closed set [10] if $\text{cl}(A) \subseteq U$ whenever $A \subseteq U$, U is semi open in (X, τ) .
 - 4) $\omega\alpha$ -closed set [1] if $\alpha\text{cl}(A) \subseteq U$ whenever $A \subseteq U$, U is ω - open in (X, τ) .
 - 5) $\delta\omega\alpha$ -closed set[2] if $\text{cl}_\delta(A) \subseteq U$ whenever $A \subseteq U$, U is $\omega\alpha$ - open set in (X, τ) .
- the complement of above mentioned closed sets is called respective open sets.

Definition 2.3: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called

- (i) $\delta\omega\alpha$ -continuous if $f^{-1}(V)$ is $\delta\omega\alpha$ -closed in X for every closed subset V of Y;
- (ii) $\delta\omega\alpha$ -irresolute if $f^{-1}(V)$ is $\delta\omega\alpha$ -closed in X for every $\delta\omega\alpha$ -closed subset V of Y;

3. $\delta\omega\alpha$ -OPEN FUNCTIONS

Definition 3.1: Let (X, τ) and (Y, σ) be topological spaces. A function $f: X \rightarrow Y$ is called $\delta\omega\alpha$ -open map if for every open set G in X, $f(G)$ is a $\delta\omega\alpha$ -open set in Y.

**Corresponding Author: S. Chandrasekar*¹, ¹Department of Mathematics,
Arignar Anna Government Arts college, Namakkal (DT), Tamil Nadu, India.**

Theorem 3.2: Prove that a mapping $f: (X, \tau) \rightarrow (Y, \sigma)$ is $\delta\omega\alpha$ -open if and only if for each $x \in X$, and $U \in \tau$ such that $x \in U$, there exists a $\delta\omega\alpha$ -open set $W \subseteq Y$ containing $f(x)$ such that $W \subseteq f(U)$.

Proof: Follows immediately from Definition 3.1

Theorem 3.3: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be $\delta\omega\alpha$ -open. If $W \subseteq Y$ and $F \subseteq X$ is a closed set containing $f^{-1}(W)$, then there exists a $\delta\omega\alpha$ -closed $H \subseteq Y$ containing W such that $f^{-1}(H) \subseteq F$.

Proof: Let $H = Y - f(X - F)$. From the definition of H , H is a $\delta\omega\alpha$ -closed. By our assumption $f^{-1}(W) \subseteq F$, we have $f(X - F) \subseteq (Y - W)$. Hence $f^{-1}(H) = X - f^{-1}[f(X - F)] \subseteq X - (X - F) = F$.

Theorem 3.4: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be $\delta\omega\alpha$ -open and let $B \subseteq Y$. Then $f^{-1}[\delta\omega\alpha\text{-Cl}(\delta\omega\alpha\text{-Int}(\delta\omega\alpha\text{-Cl}(B)))] \subseteq \text{Cl}[f^{-1}(B)]$

Proof: Assume that $B \subseteq Y$. We know that $f^{-1}(B) \subseteq \text{Cl}[f^{-1}(B)]$ for any set. By Theorem 3.3, there exists a $\delta\omega\alpha$ -closed set $H \subseteq Y$, such that $f^{-1}(H) \subseteq \text{Cl}[f^{-1}(B)]$. Thus, $f^{-1}[\delta\omega\alpha\text{-Cl}(\delta\omega\alpha\text{-Int}(\delta\omega\alpha\text{-Cl}(B)))] \subseteq f^{-1}[\delta\omega\alpha\text{-Cl}(\delta\omega\alpha\text{-Int}(\delta\omega\alpha\text{-Cl}(H)))] \subseteq f^{-1}(H) \subseteq \text{Cl}[f^{-1}(B)]$.

Theorem 3.5: Prove that a function $f: (X, \tau) \rightarrow (Y, \sigma)$ is $\delta\omega\alpha$ -open if and only iff $[\text{Int}(A)] \subseteq \delta\omega\alpha\text{-Int}[f(A)]$, for all $A \subseteq X$.

Proof: Necessity: Let $A \subseteq X$. Let $x \in \text{Int}(A)$. Then there exists $U_x \in \tau$ such that $x \in U_x \subseteq A$. So $f(x) \in f(U_x) \subseteq f(A)$ and by hypothesis, $f(U_x) \in \delta\omega\alpha\text{-}\sigma$. Hence $f(x) \in \delta\omega\alpha\text{-Int}[f(A)]$. Thus $f[\text{Int}(A)] \subseteq \delta\omega\alpha\text{-Int}[f(A)]$.

Sufficiency: Let $U \in \tau$. Then by hypothesis, $f[\text{Int}(U)] \subseteq \delta\omega\alpha\text{-Int}[f(U)]$. Since $\text{Int}(U) = U$ as U is open. Also $\delta\omega\alpha\text{-Int}[f(U)] \subseteq f(U)$. Hence $f(U) = \delta\omega\alpha\text{-Int}[f(U)]$. Thus $f(U)$ is $\delta\omega\alpha$ -open in Y . So f is $\delta\omega\alpha$ -open.

Theorem 3.6: Prove that a function $f: (X, \tau) \rightarrow (Y, \sigma)$ is $\delta\omega\alpha$ -open if and only if $\text{Int}[f^{-1}(B)] \subseteq f^{-1}[\delta\omega\alpha\text{-Int}(B)]$, for all $B \subseteq Y$.

Proof:

Necessity: Let $B \subseteq Y$. Since $\text{Int}[f^{-1}(B)]$ is open in X and f is $\delta\omega\alpha$ -open, $f[\text{Int}(f^{-1}(B))]$ is $\delta\omega\alpha$ -open in Y . Also we have $f[\text{Int}(f^{-1}(B))] \subseteq f[f^{-1}(B)] \subseteq B$. Hence, $f[\text{Int}(f^{-1}(B))] \subseteq \delta\omega\alpha\text{-Int}(B)$. Therefore $\text{Int}[f^{-1}(B)] \subseteq f^{-1}[\delta\omega\alpha\text{-Int}(B)]$.

Sufficiency: Let $A \subseteq X$. Then $f(A) \subseteq Y$. Hence by hypothesis, we obtain $\text{Int}(A) \subseteq \text{Int}[f^{-1}(f(A))] \subseteq f^{-1}[\delta\omega\alpha\text{-Int}(f(A))]$. Thus $f[\text{Int}(A)] \subseteq \delta\omega\alpha\text{-Int}[f(A)]$, for all $A \subseteq X$. Hence, by Theorem 3.5, f is $\delta\omega\alpha$ -open.

Theorem 3.7: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a mapping. Then a necessary and sufficient condition for f to be $\delta\omega\alpha$ -open is that $f^{-1}[\delta\omega\alpha\text{-Cl}(B)] \subseteq \text{Cl}[f^{-1}(B)]$ for every subset B of Y .

Proof:

Necessity: Assume f is $\delta\omega\alpha$ -open. Let $B \subseteq Y$. Let $x \in f^{-1}[\delta\omega\alpha\text{-Cl}(B)]$. Then $f(x) \in \delta\omega\alpha\text{-Cl}(B)$. Let $U \in \tau$ such that $x \in U$. Since f is $\delta\omega\alpha$ -open, then $f(U)$ is a $\delta\omega\alpha$ -open set in Y . Therefore, $B \cap f(U) \neq \emptyset$. Then $U \cap f^{-1}(B) \neq \emptyset$. Hence $x \in \text{Cl}[f^{-1}(B)]$. We conclude that $f^{-1}[\delta\omega\alpha\text{-Cl}(B)] \subseteq \text{Cl}[f^{-1}(B)]$.

Sufficiency: Let $B \subseteq Y$. Then $(Y - B) \subseteq Y$. By hypothesis, $f^{-1}[\delta\omega\alpha\text{-Cl}(Y - B)] \subseteq \text{Cl}[f^{-1}(Y - B)]$. This implies $X - \text{Cl}[f^{-1}(Y - B)] \subseteq X - f^{-1}[\delta\omega\alpha\text{-Cl}(Y - B)]$. Hence $X - \text{Cl}[X - f^{-1}(B)] \subseteq f^{-1}[Y - \delta\omega\alpha\text{-Cl}(Y - B)]$. Then $\text{Int}[f^{-1}(B)] \subseteq f^{-1}[\delta\omega\alpha\text{-Int}(B)]$. Now from Theorem 3.6, it follows that f is $\delta\omega\alpha$ -open.

4. $\delta\omega\alpha$ -CLOSED FUNCTIONS

In this section we introduce $\delta\omega\alpha$ -closed functions and study certain properties and characterizations of this type of functions.

Definition 4.1: A mapping $f: (X, \tau) \rightarrow (Y, \sigma)$ is called $\delta\omega\alpha$ -closed if the image of each closed set in X is a $\delta\omega\alpha$ -closed set in Y .

Theorem 4.2: Prove that a mapping $f: (X, \tau) \rightarrow (Y, \sigma)$ is $\delta\omega\alpha$ -closed if and only $\delta\omega\alpha\text{-Cl}[f(A)] \subseteq f[\text{Cl}(A)]$ for each $A \subseteq X$.

Proof:

Necessity: Let f be $\delta\omega\alpha$ -closed and let $A \subseteq X$. Then $f(A) \subseteq f[\text{Cl}(A)]$ and $f[\text{Cl}(A)]$ is a $\delta\omega\alpha$ -closed set in Y . Thus $\delta\omega\alpha\text{-Cl}[f(A)] \subseteq f[\text{Cl}(A)]$.

Sufficiency: suppose that $\delta\omega\alpha\text{-Cl}[f(A)] \subseteq f[\text{Cl}(A)]$, for each $A \subseteq X$. Let $A \subseteq X$ be a closed set. Then $\delta\omega\alpha\text{-Cl}[f(A)] \subseteq f[\text{Cl}(A)] = f(A)$. This shows that $f(A)$ is a $\delta\omega\alpha$ -closed set. Hence f is $\delta\omega\alpha$ -closed.

Theorem 4.3: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be $\delta\omega\alpha$ -closed. If $V \subseteq Y$ and $E \subseteq X$ is an open set containing $f^{-1}(V)$, then there exists a $\delta\omega\alpha$ -open set $G \subseteq Y$ containing V such that $f^{-1}(G) \subseteq E$.

Proof: Let $G = Y - f(X - E)$. Since $f^{-1}(V) \subseteq E$, we have $f(X - E) \subseteq Y - V$. Since f is $\delta\omega\alpha$ -closed, then G is a $\delta\omega\alpha$ -open set and $f^{-1}(G) = X - f^{-1}[f(X - E)] \subseteq X - (X - E) = E$.

Theorem 4.4: Suppose that $f: (X, \tau) \rightarrow (Y, \sigma)$ is a $\delta\omega\alpha$ -closed mapping. Then $\delta\omega\alpha\text{-Int}[\delta\omega\alpha\text{-Cl}(f(A))] \subseteq f[\text{Cl}(A)]$ for every subset A of X .

Proof: Suppose f is a $\delta\omega\alpha$ -closed mapping and A is an arbitrary subset of X . Then $f[\text{Cl}(A)]$ is $\delta\omega\alpha$ -closed in Y . Then $\delta\omega\alpha\text{-Int}[\delta\omega\alpha\text{-Cl}(f[\text{Cl}(A)])] \subseteq f[\text{Cl}(A)]$. But also $\delta\omega\alpha\text{-Int}[\delta\omega\alpha\text{-Cl}(f(A))] \subseteq \delta\omega\alpha\text{-Int}[\delta\omega\alpha\text{-Cl}(f[\text{Cl}(A)])]$. Hence $\delta\omega\alpha\text{-Int}[\delta\omega\alpha\text{-Cl}(f(A))] \subseteq f[\text{Cl}(A)]$.

Theorem 4.5: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a $\delta\omega\alpha$ -closed function, and $B, C \subseteq Y$.

- (i) If U is an open neighborhood of $f^{-1}(B)$, then there exists a $\delta\omega\alpha$ -open neighborhood V of B such that $f^{-1}(B) \subseteq f^{-1}(V) \subseteq U$.
- (ii) If f is also onto, then if $f^{-1}(B)$ and $f^{-1}(C)$ have disjoint open neighborhoods, so have B and C .

Proof:

- (i) Let $V = Y - f(X - U)$. Then $V^c = Y - V = f(U^c)$. Since f is $\delta\omega\alpha$ -closed, so V is a $\delta\omega\alpha$ -open set. Since $f^{-1}(B) \subseteq U$, we have $V^c = f(U^c) \subseteq f[f^{-1}(B^c)] \subseteq B^c$. Hence, $B \subseteq V$, and thus V is a $\delta\omega\alpha$ -open neighborhood of B . Further $U^c \subseteq f^{-1}[f(U^c)] = f^{-1}(V^c) = [f^{-1}(V)]^c$. This proves that $f^{-1}(V) \subseteq U$.
- (ii) If $f^{-1}(B)$ and $f^{-1}(C)$ have disjoint open neighborhoods M and N , then by (i), we have $\delta\omega\alpha$ -open neighborhoods U and V of B and C respectively such that $f^{-1}(B) \subseteq f^{-1}(U) \subseteq \delta\omega\alpha\text{-Int}(M)$ and $f^{-1}(C) \subseteq f^{-1}(V) \subseteq \delta\omega\alpha\text{-Int}(N)$. Since M and N are disjoint, so are $\delta\omega\alpha\text{-Int}(M)$ and $\delta\omega\alpha\text{-Int}(N)$, and hence so $f^{-1}(U)$ and $f^{-1}(V)$ are disjoint as well. It follows that U and V are disjoint too as f is onto.

Theorem 4.6: Prove that a surjective mapping $f: (X, \tau) \rightarrow (Y, \sigma)$ is $\delta\omega\alpha$ -closed if and only if for each subset B of Y and each open set U in X containing $f^{-1}(B)$, there exists a $\delta\omega\alpha$ -open set V in Y containing B such that $f^{-1}(V) \subseteq U$.

Proof:

Necessity: This follows from (1) of Theorem 4.5.

Sufficiency: Suppose F is an arbitrary closed set in X . Let y be an arbitrary point in $Y - f(F)$. Then $f^{-1}(y) \subseteq X - f^{-1}[f(F)] \subseteq (X - F)$ and $(X - F)$ is open in X . Hence by hypothesis, there exists a $\delta\omega\alpha$ -open set V_y containing y such that $f^{-1}(V_y) \subseteq (X - F)$. This implies that $y \in V_y \subseteq [Y - f(F)]$. Thus $Y - f(F) = \cup\{V_y: y \in Y - f(F)\}$. Hence $Y - f(F)$, being a union of $\delta\omega\alpha$ -open sets, is $\delta\omega\alpha$ -open. Thus its complement $f(F)$ is $\delta\omega\alpha$ -closed. This shows that f is $\delta\omega\alpha$ -closed.

Theorem 4.6: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a bijection. Then the following are equivalent:

- (i) f is $\delta\omega\alpha$ -closed.
- (ii) f is $\delta\omega\alpha$ -open.
- (iii) f^{-1} is $\delta\omega\alpha$ -continuous.

Proof:

(i) = (ii): Let $U \in \tau$. Then $X - U$ is closed in X . By (i), $f(X - U)$ is $\delta\omega\alpha$ -closed in Y . But $f(X - U) = f(X) - f(U) = Y - f(U)$. Thus $f(U)$ is $\delta\omega\alpha$ -open in Y . This shows that f is $\delta\omega\alpha$ -open.

(ii) = (iii): Let $U \subseteq X$ is an open set. Since f is $\delta\omega\alpha$ -open. So $f(U) = (f^{-1})^{-1}(U)$ is $\delta\omega\alpha$ -open in Y . Hence f^{-1} is $\delta\omega\alpha$ -continuous.

(iii) = (i): Let A be an arbitrary closed set in X . Then $X - A$ is open in X . Since f^{-1} is $\delta\omega\alpha$ -continuous, $(f^{-1})^{-1}(X - A)$ is $\delta\omega\alpha$ -open in Y . But $(f^{-1})^{-1}(X - A) = f(X - A) = Y - f(A)$. Thus $f(A)$ is $\delta\omega\alpha$ -closed in Y . This shows that f is $\delta\omega\alpha$ -closed.

Remark 4.7: A bijection $f: (X, \tau) \rightarrow (Y, \sigma)$ may be open and closed but neither $\delta\omega\alpha$ -open nor $\delta\omega\alpha$ -closed.

5. QUASI $\delta\omega\alpha$ -OPEN FUNCTIONS

We introduce a new definition as follows:

Definition 5.1: A function $f: X \rightarrow Y$ is said to be quasi $\delta\omega\alpha$ -open if the image of every $\delta\omega\alpha$ -open set in X is open in Y .

It is evident that, the concepts of quasi $\delta\omega\alpha$ -openness and $\delta\omega\alpha$ -continuity coincide if the function is a bijection.

Theorem 5.2: A function $f: X \rightarrow Y$ is quasi $\delta\omega\alpha$ -open if and only if for every subset U of X , $f(\delta\omega\alpha\text{-int}(U)) \subset \text{int}(f(U))$.

Proof: Let f be a quasi $\delta\omega\alpha$ -open function and U be a subset of X . Now, we have $\delta\omega\alpha\text{-int}(U) \subset U$ and $\delta\omega\alpha\text{-int}(U)$ is a $\delta\omega\alpha$ -open set. Hence, we obtain that $f(\delta\omega\alpha\text{-int}(U)) \subset f(U)$. As $f(\delta\omega\alpha\text{-int}(U))$ is open, $f(\delta\omega\alpha\text{-int}(U)) \subset \text{int}(f(U))$. Conversely, assume that U is a $\delta\omega\alpha$ -open set in X . Then, $f(U) = f(\delta\omega\alpha\text{-int}(U)) \subset \text{int}(f(U))$ but $\text{int}(f(U)) \subset f(U)$. Consequently, $f(U) = \text{int}(f(U))$ and $f(U)$ is open in Y . Hence f is quasi $\delta\omega\alpha$ -open.

Lemma 5.3: If a function $f: X \rightarrow Y$ is quasi $\delta\omega\alpha$ -open, then $\delta\omega\alpha\text{-int}(f^{-1}(G)) \subset f^{-1}(\text{int}(G))$ for every subset G of Y .

Proof: Let G be any arbitrary subset of Y . Then, $\delta\omega\alpha\text{-int}(f^{-1}(G))$ is a $\delta\omega\alpha$ -open set in X and since f is quasi $\delta\omega\alpha$ -open, then $f(\delta\omega\alpha\text{-int}(f^{-1}(G))) \subset \text{int}(f(f^{-1}(G))) \subset \text{int}(G)$. Thus, $\delta\omega\alpha\text{-int}(f^{-1}(G)) \subset f^{-1}(\text{int}(G))$.

Recall that a subset S is called a $\delta\omega\alpha$ -neighbourhood of a point x of X if there exists a $\delta\omega\alpha$ -open set U such that $x \in U \subset S$.

Theorem 5.4: For a function $f: X \rightarrow Y$, the following are equivalent:

- (i) f is quasi $\delta\omega\alpha$ -open;
- (ii) For each subset U of X , $f(\delta\omega\alpha\text{-int}(U)) \subset \text{int}(f(U))$;
- (iii) For each $x \in X$ and each $\delta\omega\alpha$ -neighbourhood U of x in X , there exists a neighbourhood V of $f(x)$ in Y such that $V \subset f(U)$.

Proof

(i) \Rightarrow (ii): It follows from Theorem 5.2.

(ii) \Rightarrow (iii): Let $x \in X$ and U be an arbitrary $\delta\omega\alpha$ -neighbourhood of x in X .

Then there exists a $\delta\omega\alpha$ -open set V in X such that $x \in V \subset U$. Then by (ii), we have $f(V) = f(\delta\omega\alpha\text{-int}(V)) \subset \text{int}(f(V))$ and hence $f(V) = \text{int}(f(V))$. Therefore, it follows that $f(V)$ is open in Y such that $f(x) \in f(V) \subset f(U)$.

(iii) \Rightarrow (i): Let U be an arbitrary $\delta\omega\alpha$ -open set in X such that $x \in U$. Then for each $f(x) = y \in f(U)$, by (iii) there exists a neighbourhood V_y of y in Y such that $V_y \subset f(U)$. As V_y is a neighbourhood of y , there exists an open set W_y in Y such that $y \in W_y \subset V_y$. Thus $f(U) = \cup \{W_y: y \in f(U)\}$ which is an open set in Y . This implies that f is quasi $\delta\omega\alpha$ -open function.

Theorem 5.5: A function $f: X \rightarrow Y$ is quasi $\delta\omega\alpha$ -open if and only if for any subset B of Y and for any $\delta\omega\alpha$ -closed set F of X containing $f^{-1}(B)$, there exists a closed set G of Y containing B such that $f^{-1}(G) \subset F$.

Proof: Suppose f is quasi $\delta\omega\alpha$ -open. Let $B \subset Y$ and F be a $\delta\omega\alpha$ -closed set of X containing $f^{-1}(B)$. Now, put $G = Y - f(X - F)$. It is clear that $f^{-1}(B) \subset F$ implies $B \subset G$. Since f is quasi $\delta\omega\alpha$ -open, we obtain G is a closed set of Y . Moreover, we have $f^{-1}(G) \subset F$.

Conversely, let U be a $\delta\omega\alpha$ -open set of X and put $B = Y - f(X - U)$. Then $X - U$ is a $\delta\omega\alpha$ -closed set in X containing $f^{-1}(B)$. By hypothesis, there exists a closed set F of Y such that $B \subset F$ and $f^{-1}(F) \subset X - U$. Hence, we obtain $f(U) \subset Y - F$. On the other hand, it follows that $B \subset F$, $Y - F \subset Y - B = f(U)$. Thus, we obtain $f(U) = Y - F$ which is open and hence f is a quasi $\delta\omega\alpha$ -open function.

Theorem 5.6: A function $f: X \rightarrow Y$ is quasi $\delta\omega\alpha$ -open if and only if $f^{-1}(\text{cl}(B)) \subset \delta\omega\alpha\text{-cl}(f^{-1}(B))$ for every subset B of Y .

Proof: Suppose that f is quasi $\delta\omega\alpha$ -open. For any subset B of Y , $f^{-1}(B) \subset \delta\omega\alpha\text{-cl}(f^{-1}(B))$. Therefore by Theorem 5.5, there exists a closed set F in Y such that $B \subset F$ and $f^{-1}(F) \subset \delta\omega\alpha\text{-cl}(f^{-1}(B))$. Therefore, we obtain $f^{-1}(\text{cl}(B)) \subset f^{-1}(F) \subset \delta\omega\alpha\text{-cl}(f^{-1}(B))$.

Conversely, let $B \subset Y$ and F be a $\delta\omega\alpha$ -closed of X containing $f^{-1}(B)$. Put $W = \text{cl}_Y(B)$, then we have $B \subset W$ and W is closed in Y and $f^{-1}(W) \subset \delta\omega\alpha\text{-cl}(f^{-1}(B)) \subset F$. Then, by Theorem 5.5, f is quasi $\delta\omega\alpha$ -open.

Lemma 5.7: Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be two functions and $g \circ f: X \rightarrow Z$ is quasi $\delta\omega\alpha$ -open. If g is continuous and injective, then f is quasi $\delta\omega\alpha$ -open.

Proof: Let U be a $\delta\omega\alpha$ -open set in X . Then $(g \circ f)(U)$ is open in Z since $g \circ f$ is quasi $\delta\omega\alpha$ -open. Again g is an injective continuous function, $f(U) = g^{-1}(g \circ f(U))$ is open in Y . This shows that f is quasi $\delta\omega\alpha$ -open.

6. QUASI $\delta\omega\alpha$ -CLOSED FUNCTIONS

Definition 6.1: A function $f: X \rightarrow Y$ is said to be quasi $\delta\omega\alpha$ -closed if the image of each $\delta\omega\alpha$ -closed set in X is closed in Y .

Remark 6.2: Quasi $\delta\omega\alpha$ -closed function is independent to closed map.

Example 6.3: Let $X = Y = \{a, b, c\}$, $\tau = \{\phi, \{a\}, \{a, b\}, X\}$ and $\sigma = \{\phi, \{a\}, Y\}$. Define a function $f: (X, \tau) \rightarrow (Y, \sigma)$ be identity map. Then f is quasi $\delta\omega\alpha$ -closed but not closed.

Example 6.4: Let $X = Y = \{a, b, c\}$, $\tau = \sigma = \{\phi, \{a\}, \{b, c\}, X\}$ and closed set $\{\phi, \{a\}, \{b, c\}, X\}$ and $\delta\omega\alpha$ -closed set $\{\text{all power set in } X\}$. Define a function $f: (X, \tau) \rightarrow (Y, \sigma)$ be identity map. Clearly closed but not quasi $\delta\omega\alpha$ -closed function

Remark 6.5: Quasi $\delta\omega\alpha$ -closed function is independent to $\delta\omega\alpha$ -closed map as shown by the following example.

Example 6.6: Let $X = Y = \{a, b, c\}$, $\tau = \sigma = \{\phi, \{a\}, \{a, b\}, \{a, c\}, X\}$ and closed set $\{\phi, \{b\}, \{c\}, \{b, c\}, X\}$ and $\delta\omega\alpha$ -closed set $\{\phi, \{b, c\}, X\}$.

Example 6.7: Let function $f: (X, \tau) \rightarrow (Y, \sigma)$ be identity map. Clearly quasi $\delta\omega\alpha$ -closed function but not $\delta\omega\alpha$ -closed map let $X = Y = \{a, b, c\}$, $\tau = \sigma = \{\phi, \{a\}, \{b, c\}, X\}$ and closed set $\{\phi, \{a\}, \{b, c\}, X\}$ and $\delta\omega\alpha$ -closed set $\{\text{all power set in } X\}$.

Define a function $f: (X, \tau) \rightarrow (Y, \sigma)$ be identity map. Clearly $\delta\omega\alpha$ closed but not quasi $\delta\omega\alpha$ -closed function

Lemma 6.8: If a function $f: X \rightarrow Y$ is quasi $\delta\omega\alpha$ -closed, then $f^{-1}(\text{cl}(B)) \subset \delta\omega\alpha\text{-cl}(f^{-1}(B))$ for every subset B of Y .

Proof: This proof is similar to the proof of Lemma 5.3.

Theorem 6.9: A function $f: X \rightarrow Y$ is quasi $\delta\omega\alpha$ -closed if and only if for any subset B of Y and for any $\delta\omega\alpha$ -open set G of X containing $f^{-1}(B)$, there exists an open set U of Y containing B such that $f^{-1}(U) \subset G$.

Proof: This proof is similar to that of Theorem 5.5.

Definition 6.10: A function $f: X \rightarrow Y$ is called $\delta\omega\alpha^*$ -closed if the image of every $\delta\omega\alpha$ -closed subset of X is $\delta\omega\alpha$ -closed in Y .

Theorem 6.11: If $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are two quasi $\delta\omega\alpha$ -closed functions, then $f: X \rightarrow Z$ is a need not be quasi $\delta\omega\alpha$ -closed function.

Proof: Obvious. Furthermore, we have the above example.

Theorem 6.12: Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be any two functions. Then

- (i) if f is $\delta\omega\alpha$ -closed and g is quasi $\delta\omega\alpha$ -closed, then $g \circ f$ is closed;
- (ii) if f is quasi $\delta\omega\alpha$ -closed and g is $\delta\omega\alpha$ -closed, then $g \circ f$ is $\delta\omega\alpha^*$ -closed;
- (iii) if f is $\delta\omega\alpha^*$ -closed and g is quasi $\delta\omega\alpha$ -closed, then $g \circ f$ is quasi $\delta\omega\alpha$ -closed.

Proof: Obvious.

Theorem 6.13: Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be two functions such that $g \circ f: X \rightarrow Z$ is quasi $\delta\omega\alpha$ -closed. Then

- (i) if f is $\delta\omega\alpha$ -irresolute surjective, then g is $\delta\omega\alpha$ -closed.
- (ii) if g is $\delta\omega\alpha$ -continuous injective, then f is $\delta\omega\alpha^*$ -closed.

Proof:

- (i) Suppose that F is an arbitrary $\delta\omega\alpha$ -closed in Y . As f is $\delta\omega\alpha$ -irresolute, $f^{-1}(F)$ is $\delta\omega\alpha$ -closed in X . Since $g \circ f$ is quasi $\delta\omega\alpha$ -closed and f is surjective, $g \circ f(f^{-1}(F)) = g(F)$, which is closed in Z . This implies that g is a $\delta\omega\alpha$ -closed function.
- (ii) Suppose that F is any $\delta\omega\alpha$ -closed set in X . Since $g \circ f$ is quasi $\delta\omega\alpha$ -closed, $(g \circ f)(F)$ is closed in Z . Again g is a $\delta\omega\alpha$ -continuous injective function, $g^{-1}(g \circ f(F)) = f(F)$, which is $\delta\omega\alpha$ -closed in Y . This shows that f is $\delta\omega\alpha^*$ -closed.

Theorem 6.14: Let X and Y be topological spaces. Then the function $g : X \rightarrow Y$ is a quasi $\delta\omega\alpha$ -closed if and only if $g(X)$ is closed in Y and $g(V) - g(X-V)$ is open in $g(X)$ whenever V is $\delta\omega\alpha$ -open in X .

Proof: Necessity: Suppose $g : X \rightarrow Y$ is a quasi $\delta\omega\alpha$ -closed function. Since X is $\delta\omega\alpha$ -closed, $g(X)$ is closed in Y and $g(V) - g(X-V) = g(V) \cap g(X) - g(X-V)$ is open in $g(X)$ when V is $\delta\omega\alpha$ -open in X .

Sufficiency: Suppose $g(X)$ is closed in Y , $g(V) - g(X-V)$ is open in $g(X)$ when V is $\delta\omega\alpha$ -open in X , and let C be closed in X . Then $g(C) = g(X) - (g(X-C) - g(C))$ is closed in $g(X)$ and hence, closed in Y .

Corollary 6.15: Let X and Y be topological spaces. Then a surjective function $g : X \rightarrow Y$ is quasi $\delta\omega\alpha$ -closed if and only if $g(V) - g(X-V)$ is open in Y whenever V is $\delta\omega\alpha$ -open in X .

Proof: Obvious.

Corollary 6.16: Let X and Y be topological spaces and let $g : X \rightarrow Y$ be a $\delta\omega\alpha$ -continuous quasi $\delta\omega\alpha$ -closed surjective function. Then the topology on Y is $\{g(V) - g(X-V) : V \text{ is } \delta\omega\alpha\text{-open in } X\}$.

Proof: Let W be open in Y . Then $g^{-1}(W)$ is $\delta\omega\alpha$ -open in X , and $g(g^{-1}(W)) - g(X - g^{-1}(W)) = W$. Hence, all open sets in Y are of the form $g(V) - g(X-V)$, V is $\delta\omega\alpha$ -open in X . On the other hand, all sets of the form $g(V) - g(X-V)$, V is $\delta\omega\alpha$ -open in X , are open in Y from Corollary 6.15.

Definition 6.17: A topological space (X, τ) is said to be $\delta\omega\alpha^*$ -normal if for any pair of disjoint $\delta\omega\alpha$ -closed subsets F_1 and F_2 of X , there exist disjoint open sets U and V such that $F_1 \subset U$ and $F_2 \subset V$.

Theorem 6.18: Let X and Y be topological spaces with X is $\delta\omega\alpha^*$ -normal. If $g : X \rightarrow Y$ is a $\delta\omega\alpha$ -continuous quasi $\delta\omega\alpha$ -closed surjective function, then Y is normal.

Proof: Let K and M be disjoint closed subsets of Y . Then $g^{-1}(K)$ and $g^{-1}(M)$ are disjoint $\delta\omega\alpha$ -closed subsets of X . Since X is $\delta\omega\alpha^*$ -normal, there exist disjoint open sets V and W such that $g^{-1}(K) \subset V$ and $g^{-1}(M) \subset W$. Then $K \subset (g(V) - g(X-V))$ and $M \subset (g(W) - g(X-W))$. Further by Corollary 6.15, $(g(V) - g(X-V))$ and $(g(W) - g(X-W))$ are open sets in Y and clearly $(g(V) - g(X-V)) \cap (g(W) - g(X-W)) = \phi$. This shows that Y is normal.

REFERENCES

1. S. S. Benchalli, P. G. Patil and T. D. Rayanagaudar, $\omega\alpha$ -Closed Sets in Topological Spaces, The Global J. Appl. Math. and Math. Sci., 2, No. 1-2, (2009), 53-63.
2. S.Chandrasekar T.Rajesh Kannan and, M.Suresh, $\delta\omega\alpha$ -closed sets in Topological Spaces Global Journal of Mathematical Sciences, Theory and Practical. Vol. 9, N 2 (2017), pp. 103-116.
3. S.Chandrasekar, A.Atkinswestley and M.Sathyabama, $g\#b$ -closed sets in Topological Spaces, Int.Journal of Advanced Research, 5(7) (2017), 2722-2730.
4. S.Chandrasekar, M.Suresh and T.Rajesh Kannan, On Quasi Soft sg -open and Quasi Soft sg -closed Functions in soft topological spaces, International Journal of Mathematics And its Applications, 5(2-B)(2017), 239-246.
5. S.Chandrasekar, M.Sathyabama and A.Atkinswestley, $g\#b$ -interior and $g\#b$ closure in Topological Spaces International Journal of Mathematics And its Applications, Vol.5, Issue3-B (2017), 151-157.
6. O. Njastad, On some classes of nearly open sets, Pacific J. Math. 15(1965), 961-970.
7. O.Ravi, S.Ganesan and S.Chandrasekar, Almost αg_s -closed functions and separation axioms. Bulletin of Mathematical Analysis and Applications, Vol.3 Issue 1(2011), 165-177.
8. O.Ravi, S.Ganesan and S.Chandrasekar - \tilde{g} -pre-closed sets in topology International urnal of Mathematical Archive-2(2), Feb. - 2011, Page: 294- 299
9. O.Ravi, S.Chandrasekar and S.Ganesan (\tilde{g}, s)-Continuous Functions between Topological Spaces, Kyungpook Math. J. 51(2011), 323-338
10. P.Sundaram and M.Sheik John, On w -closed sets in topology, Acta Ciencia Indica 4(2000), 389-92.
11. N.V.Velicko, H-Closed Topological Spaces, Amer. Math. Soc. Transl, 78, No.2 (1968), 103-18.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2017. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]