

ON SEMI ROUGH SETS

R. PANDURANGA RAO\*

Assistant Professor, Department of Humanities & Sciences,  
Ellenki College of Engineering and Technology, Patelguda Village, Ameenpur,  
Sangareddy -502319, Telangana State.

(Received On: 13-11-17; Revised & Accepted On: 04-12-17)

---

ABSTRACT

This present research article introduces semi Rough sets by using Rough membership function. Some interesting properties of semi Rough sets and Rough membership functions are studied. A Rough membership function was introduced by Zdzislaw Pawlak together with Skowron [4] in 1991 and it is used to generate semi Rough sets in this work in a finite universe set. Some of their properties are discussed in this context.

AMS Subject Classification: 06B10, 16P70, 37A20, 46J20.

Key Words: Universe set, Information system, Equivalence relation, Lower Rough Approximation, Upper Rough Approximation, Rough set, Rough Membership function, Semi Rough sets.

---

INTRODUCTION

The problem of imperfect knowledge has been tackled for a long time by philosophers, logicians and mathematicians. Recently it became also a crucial issue for computer scientists, particularly in the area of Artificial Intelligence.

There are many approaches to the problem of how to understand and manipulate imperfect knowledge. The most successful approaches to tackle this problem are the Fuzzy set theory and the Rough set theory. Theories of Fuzzy sets and Rough sets are powerful mathematical tools for modeling various types of uncertainties. Fuzzy set theory was introduced by L. A. Zadeh in his classical paper [5] of 1965.

A polish applied mathematician and computer scientist Zdzislaw Pawlak introduced Rough set theory in his classical paper [2] of 1982. Rough set theory is a new mathematical approach to imperfect knowledge. This theory presents still another attempt to deal with uncertainty or vagueness.

The Rough set theory has attracted the attention of many researchers and practitioners who contributed essentially to its development and application. Rough sets have been proposed for a very wide variety of applications. In particular, the Rough set approach seems to be important for Artificial Intelligence and cognitive sciences, especially for machine learning, knowledge discovery, data mining, pattern recognition and approximate reasoning.

Throughout this research article, let  $\phi$  and  $\mathcal{U}$  stand for the empty set and a finite universe set respectively. Let  $|A|$  denote the number of elements in  $A$ , where  $A$  is any subset of  $\mathcal{U}$ .

1. PRELIMINARIES

This section is devoted to some basic definitions which are needed for the further study of this Article.

**Definition 1.1:** A relation  $R$  on  $\mathcal{U}$  is said to be an *equivalence relation* on  $\mathcal{U}$  if

- (a)  $(x, x) \in R$  for every  $x \in \mathcal{U}$  (reflexivity)
  - (b)  $(x, y) \in R \Leftrightarrow (y, x) \in R$  for every  $x, y \in \mathcal{U}$  (symmetry)
  - (c)  $(x, y) \in R$  and  $(y, z) \in R \Rightarrow (x, z) \in R$  for every  $x, y, z \in \mathcal{U}$  (transitivity)
- 

Corresponding Author: R. Panduranga Rao\*

**Definition 1.2:** If  $R$  is an equivalence relation on  $\mathcal{U}$  then the *equivalence class* of an element  $x \in \mathcal{U}$  is denoted by the symbol  $[x]_R$  and it is defined by  $[x]_R = \{y \in \mathcal{U} : yRx\}$ .

**Definition- 1.3:** An *information system* is a pair  $(\mathcal{U}, \mathcal{A})$  where  $\mathcal{A}$  is a set of attributes. Each attribute  $f \in \mathcal{A}$  is a mapping  $f : \mathcal{U} \rightarrow V_f$  where  $V_f$  is the range set of the attribute  $f \in \mathcal{A}$ . Each attribute  $f \in \mathcal{A}$  generates an equivalence relation on  $\mathcal{U}$ .

Corresponding to each attribute  $f \in \mathcal{A}$ , a relation  $R_f$  is defined on  $\mathcal{U}$  such that  $(x, y) \in R_f \Leftrightarrow f(x) = f(y)$ . It is easy to verify that  $R_f$  is an equivalence relation on  $\mathcal{U}$  and  $R_f$  is called an *indiscernible relation*. If  $x \in \mathcal{U}$  then  $[x]_{R_f} = \{y \in \mathcal{U} : f(x) = f(y)\}$ .

**Definition 1.4:** Let  $(\mathcal{U}, \mathcal{A})$  be an information system and  $R_f$ , an indiscernible relation on  $\mathcal{U}$  for some  $f \in \mathcal{A}$ . If  $X$  is any subset of  $\mathcal{U}$  then

- a) The *lower Rough approximation* of  $X$  is defined to be the set

$$R_*(X) = \{x \in \mathcal{U} : [x]_{R_f} \subseteq X\}$$

- b) The *upper Rough approximation* of  $X$  is defined to be the set

$$R^*(X) = \{x \in \mathcal{U} : [x]_{R_f} \cap X \neq \emptyset\}$$

- c) The *boundary region* of  $X$  with respect to  $R_f$  is defined to be the set

$$\mathfrak{B}_{R_f}(X) = R^*(X) - R_*(X)$$

**Definition 1.5:** A subset  $X$  of  $\mathcal{U}$  is said to be a *Rough set*, if the boundary region  $\mathfrak{B}_{R_f}(X) = R^*(X) - R_*(X)$  is non- empty. Sometimes, a Rough set  $X$  can also be represented as a pair  $(R_*(X), R^*(X))$  using Rough approximations.

**Definition 1.6:** A subset  $X$  of  $\mathcal{U}$  is said to be a *Crisp set* if  $R_*(X) = X = R^*(X)$ .

It is easy to observe that a subset  $X$  of  $\mathcal{U}$  is a Crisp set if and only if it is not a Rough set.

## 2. ROUGH MEMBERSHIP FUNCTION

In most of the cases, the universe set is finite. The Rough membership function seems to be a very useful tool to deal with such conditions. The lower and upper Rough approximations can be obtained by using a rough membership function when the universe set is finite. Let the universe set  $\mathcal{U}$  be a non-empty finite set.

**Definition 2.1:** Let  $(\mathcal{U}, \mathcal{A})$  be a finite information system. Fix an indiscernible relation  $R_f$  corresponding to an attribute  $f \in \mathcal{A}$ . If  $A$  is any subset of  $\mathcal{U}$  then the Rough membership function  $\lambda_A : \mathcal{U} \rightarrow [0,1]$  is defined as follows.

$$\lambda_A(x) = \frac{|[x]_{R_f} \cap A|}{|[x]_{R_f}|} \quad \forall x \in \mathcal{U}.$$

The Rough membership function expresses conditional probability that  $x \in A$  given  $R_f$  and can be interpreted as a degree that  $x \in A$  in view of information about  $x$  expressed by  $R_f$ .

**Theorem 2.2:** If  $A \subseteq \mathcal{U}$ , then

- a)  $R^*(A) = \{x \in \mathcal{U} : \lambda_A(x) > 0\}$
- b)  $R_*(A) = \{x \in \mathcal{U} : \lambda_A(x) = 1\}$
- c)  $\mathfrak{B}_{R_f}(A) = \{x \in \mathcal{U} : 0 < \lambda_A(x) < 1\}$

**Proof:** suppose that  $A \subseteq \mathcal{U}$  and  $R_f$  is an indiscernible relation on  $\mathcal{U}$ .

$$\begin{aligned} \text{a) } x \in \{x \in \mathcal{U} : \lambda_A(x) > 0\} &\Leftrightarrow \lambda_A(x) > 0 \\ &\Leftrightarrow \frac{|[x]_{R_f} \cap A|}{|[x]_{R_f}|} > 0 \\ &\Leftrightarrow |[x]_{R_f} \cap A| > 0 \\ &\Leftrightarrow [x]_{R_f} \cap A \neq \phi \\ &\Leftrightarrow x \in R^*(A) \end{aligned}$$

Thus  $R^*(A) = \{x \in \mathcal{U} : \lambda_A(x) > 0\}$ .

$$\begin{aligned} \text{b) } x \in \{x \in \mathcal{U} : \lambda_A(x) = 1\} &\Leftrightarrow \lambda_A(x) = 1 \\ &\Leftrightarrow \frac{|[x]_{R_f} \cap A|}{|[x]_{R_f}|} = 1 \\ &\Leftrightarrow |[x]_{R_f} \cap A| = |[x]_{R_f}| \\ &\Leftrightarrow [x]_{R_f} \cap A = [x]_{R_f} \\ &\Leftrightarrow [x]_{R_f} \subseteq A \\ &\Leftrightarrow x \in R_*(A) \end{aligned}$$

Thus  $R_*(A) = \{x \in \mathcal{U} : \lambda_A(x) = 1\}$ .

$$\begin{aligned} \text{c) Clearly, } \mathfrak{B}_{R_f}(A) &= R^*(A) - R_*(A) \\ &= \{x \in \mathcal{U} : \lambda_A(x) > 0\} - \{x \in \mathcal{U} : \lambda_A(x) = 1\} \\ &= \{x \in \mathcal{U} : 0 < \lambda_A(x) < 1\}. \end{aligned}$$

**Remark 2.3:** Suppose that  $A \subseteq \mathcal{U}$  and  $R_f$  is an indiscernible relation on  $\mathcal{U}$ . We write  $\lambda$  instead of writing  $\lambda_A$  for simplicity. If we define another indiscernible relation  $R_\lambda$  on  $\mathcal{U}$  corresponding to the Rough membership function  $\lambda$  such that

$$(x, y) \in R_\lambda \Leftrightarrow \lambda(x) = \lambda(y)$$

Then we can observe the following.

$$\begin{aligned} (x, y) \in R_f &\Leftrightarrow [x]_{R_f} = [y]_{R_f} \\ &\Leftrightarrow \frac{|[x]_{R_f} \cap A|}{|[x]_{R_f}|} = \frac{|[y]_{R_f} \cap A|}{|[y]_{R_f}|} \\ &\Leftrightarrow \lambda(x) = \lambda(y) \\ &\Leftrightarrow (x, y) \in R_\lambda \end{aligned}$$

Hence  $R_f = R_\lambda$ . Thus the process of generating indiscernible relation using a Rough membership function terminates at the first stage itself.

**Theorem 2.4:** If  $A$  and  $B$  are any two subsets of  $\mathcal{U}$ , then

- a)  $\lambda_\phi = 0$
- b)  $\lambda_{\mathcal{U}} = 1$
- c)  $\lambda_{A \cup B} = \lambda_A + \lambda_B - \lambda_{A \cap B}$
- d)  $\lambda_{A \cup B} = \lambda_A + \lambda_B$  provided  $A \cap B = \phi$
- e)  $\lambda_{A^c} = 1 - \lambda_A$  where  $A^c = \mathcal{U} - A$

**Proof:** Clearly,  $\lambda_\phi(x) = \frac{|[x]_{R_f} \cap \phi|}{|[x]_{R_f}|} = \frac{|\phi|}{|[x]_{R_f}|} = 0$  and

$$\lambda_{\mathcal{U}}(x) = \frac{|[x]_{R_f} \cap \mathcal{U}|}{|[x]_{R_f}|} = \frac{|[x]_{R_f}|}{|[x]_{R_f}|} = 1.$$

Now we prove (c).

$$\begin{aligned} \lambda_{A \cup B}(x) &= \frac{|[x]_{R_f} \cap (A \cup B)|}{|[x]_{R_f}|} = \frac{|([x]_{R_f} \cap A) \cup ([x]_{R_f} \cap B)|}{|[x]_{R_f}|} \\ &= \frac{|[x]_{R_f} \cap A| + |[x]_{R_f} \cap B| - |[x]_{R_f} \cap A \cap B|}{|[x]_{R_f}|} \\ &= \lambda_A(x) + \lambda_B(x) - \lambda_{A \cap B}(x) \quad \forall x \in \mathcal{U} \\ \Rightarrow \lambda_{A \cup B} &= \lambda_A + \lambda_B - \lambda_{A \cap B}. \end{aligned}$$

Now, suppose that  $A \cap B = \phi$ . Then  $\lambda_{A \cup B} = \lambda_A + \lambda_B - \lambda_{A \cap B}$ .

$$\begin{aligned} &= \lambda_A + \lambda_B - \lambda_\phi \\ &= \lambda_A + \lambda_B. \end{aligned}$$

Since  $A \cup A^c = \mathcal{U}$  and  $A \cap A^c = \phi$ ,  $\lambda_{A \cup A^c} = \lambda_A + \lambda_{A^c}$ .

$$\begin{aligned} \Rightarrow \lambda_{\mathcal{U}} &= \lambda_A + \lambda_{A^c} \\ \Rightarrow 1 &= \lambda_A + \lambda_{A^c} \\ \Rightarrow \lambda_{A^c} &= 1 - \lambda_A. \end{aligned}$$

### 3. SEMI ROUGH SETS

In this section, we introduce the notions of lower and upper semi Rough sets. The concept of exact rough set is also defined. A few properties of these notions are established.

**Definition 3.1:** Let  $A$  be any subset of a finite universe set  $\mathcal{U}$  and  $\lambda_A : \mathcal{U} \rightarrow [0,1]$  be the Rough membership function corresponding to  $A$ . Put

$$\theta_A = \min \{ \lambda_A(x) : x \in \mathcal{U} \text{ and } \lambda_A(x) \neq 0 \}$$

Then  $A$  is said to be

- (a) A lower semi Rough set if,  $0 < \theta_A \leq 1/2$ .

(b) An upper semi Rough set if,  $\frac{1}{2} \leq \theta_A \leq 1$ .

(c) An exact Rough set if,  $\theta_A = \frac{1}{2}$ .

**Example 3.2:** Let  $\mathcal{U} = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$  and let  $d$  be defined on  $\mathcal{U}$  by  $d(n) =$  The number of divisors of  $n$ .

The indiscernible relation  $R_d$  is given by  $(m, n) \in R_d \Leftrightarrow d(m) = d(n)$ .

Then we have  $[1]_{R_d} = \{1\}$ ,  $[2]_{R_d} = \{2, 3, 5, 7, 11\}$ ,  $[4]_{R_d} = \{4, 9\}$ ,  $[6]_{R_d} = \{6, 8, 10\}$  and  $[12]_{R_d} = \{12\}$

Let  $A = \{1, 3, 5\}$ . Then  $\lambda_A(1) = 1$ ,  $\lambda_A(2) = \lambda_A(3) = \lambda_A(5) = \lambda_A(7) = \lambda_A(11) = \frac{2}{5}$  and

$\lambda_A(4) = \lambda_A(6) = \lambda_A(8) = \lambda_A(9) = \lambda_A(10) = \lambda_A(12) = 0$ . Then the non-zero values of  $\lambda_A(x)$  are 1 and  $\frac{2}{5}$ .

$\Rightarrow \theta_A = \frac{2}{5}$  hence  $A$  is a lower semi Rough set.

Let  $B = \{1, 4\}$ . Then  $\lambda_B(1) = 1$ ,

$\lambda_B(2) = \lambda_B(3) = \lambda_B(5) = \lambda_B(7) = \lambda_B(11) = \lambda_B(6) = \lambda_B(8) = \lambda_B(10) = \lambda_B(12) = 0$  and

$\lambda_B(4) = \lambda_B(9) = \frac{1}{2}$ .

The non-zero values of  $\lambda_B(x)$  are 1 and  $\frac{1}{2}$ .

$\Rightarrow \theta_B = \frac{1}{2}$

$\Rightarrow B$  is an exact Rough set.

If  $C = \{1\}$  then  $C$  is an upper semi Rough set.

**Proposition 3.3:** If  $A \subseteq B$ , then  $\theta_A \leq \theta_B$

## REFERENCES

1. Herstein I.N., *Topics in Algebra*, 2<sup>nd</sup> edition, Wiley&Sons, New York, 1975.
2. Pawlak Z., *Rough Sets*, International Journal of Computer and Information Sciences, Vol-11, pp.341-356, 1982.
3. Pawlak Z., *Rough Sets Theoretical Aspects of Reasoning about Data*, Kluwer Academic Publishers, Dordrecht, 1991.
4. Pawlak. Z., and Skowron.A., *Rough Membership functions*, ICS Research Report, 10-91, Warsaw University of Technology, 1991.
5. Zadeh L.A., *Fuzzy sets*, Inform. Control., vol-8, pp. 338-353, 1965.

**Source of support: Nil, Conflict of interest: None Declared.**

**[Copy right © 2017. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]**