

INTUITIONISTIC FUZZY IDEALS OF M- Γ GROUPS

S.K MALA*¹, Dr. MM SHANMUGAPRIYA*²

¹PhD Scholar in Mathematics,
Karpagam University, Coimbatore, Tamil Nadu- 641021, India.

&
Assistant Professor of Mathematics,
KG College of Arts and Science, Saravanampatti, Tamil Nadu – 6410351, India.

²Asst. Professor and Head of Department (i/c) in Mathematics,
Karpagam Academy of Higher Education, Coimbatore, Tamil Nadu- 641021, India.

ABSTRACT

In this paper we define and derive lemmas and theorems of Intuitionistic Fuzzy Ideals of M- Γ Groups analogous to Fuzzy Ideals.

Keywords: Intuitionistic fuzzy rings, Near ring, Fuzzy M- Γ sub group, Intuitionistic fuzzy ideals, of Intuitionistic Fuzzy Ideals of M- Γ Groups.

INTRODUCTION

In 1986 Atanassov [1] introduced the notion of an Intuitionistic Fuzzy set as a generalization of Zadeh [9] fuzzy set. Atanassov [1] also described different operations of Intuitionistic Fuzzy set and their properties elaborately. Clay [3] discussed various properties and applications of near rings. The theories and applications of Fuzzy sets and Fuzzy logics were put forth by Klir and Yuan [9]. Characteristics and properties of Intuitionistic Fuzzy Ideals of Near Rings were discussed in 2004 by Ma and Zhan [10]. Based on the introductory study of M- Γ Groups, various results of Fuzzy Ideals of M- Γ Groups were defined, discussed and derived by Satyanarayana and Prasad [14]. An analogous definition, theorems and properties of Intuitionistic Fuzzy Ideals of M- Γ Groups are explored.

PRELIMINARIES

Definition 2.1-Crisp Sets [19]: The Crisp set is defined to classify the individuals in the Universe in two groups: Members and Non- Members

Definition 2.2-Fuzzy Sets [19]: A fuzzy set has objects with a continuum of grades of membership. Such a set is characterized by a membership (characteristic) function which assigns to each object a grade of membership ranging between zero and one.

Definition 2.3-Fuzzy Subsets [19]: Let S be any non-empty set, A mapping μ from S to [0,1] is called a Fuzzy sub set of S.

Definition 2.4-Intuitionistic Fuzzy sets [4, 9]: Let X be any non-empty set, a fuzzy subset A of X is of the form $A = \{ \langle x, \mu_A(x), \gamma_A(x) \rangle / x \in X \}$ where the functions $\mu_A : X \rightarrow [0,1]$ and $\gamma_A : X \rightarrow [0,1]$ denote the degrees of membership and non-membership of the element $x \in X$ to A respectively and satisfy $0 \leq \mu_A(x) + \gamma_A(x) \leq 1$ for all $x \in X$. The family of all intuitionistic fuzzy sets in X denoted by IFS (X).

Definition 2.5 [9]: Let A be any fuzzy set in X. Then for any $\alpha \in [0, 1]$, α cut of A, denoted by ${}^{\alpha}A$, is defined as ${}^{\alpha}A = \{x: x \in X \text{ such that } \mu_A(x) \geq \alpha\}$ and the strong α cut of A denoted by ${}^{\alpha+}A$, is defined as ${}^{\alpha+}A = \{x: x \in X \text{ such that } \mu_A(x) > \alpha\}$

Definition 2.6 [9]: Let A be any fuzzy set in X , then denote ${}_aA$ by ${}_aA(x) = \alpha \cdot {}_aA(x)$. Also ${}_{a+}A(x) = \alpha \cdot {}_{a+}A(x)$; where ${}_aA$ and ${}_{a+}A$ are fuzzy sets.

Definition 2.7 [9]: Let A be any fuzzy set in X . Then the level set of A denoted by $\wedge(A)$, is defined as

$$\wedge(A) = \{\alpha \mid A(x) = \alpha : x \in X\}$$

Definition 2.8 [14]: By a near-ring we mean a non-empty set R with two binary operations “+” and “ \cdot ” satisfying the following axioms:

- (i) $(R, +)$ is a group
- (ii) (R, \cdot) is a semigroup
- (iii) $x \cdot (y + z) = x \cdot y + x \cdot z$ for all $x, y, z \in R$.

Precisely speaking, it is a left near-ring because it satisfies the left distributive law. We will use the word “near-ring” instead of “left near-ring”. We denote xy instead of $x \cdot y$.

Definition 2.9 [14]: An ideal of a near-ring R is a subset I of R such that

- (i) $(I, +)$ is a normal subgroup of $(R, +)$
- (ii) $RI \subseteq I$
- (iii) $(x + i)y - xy \in I$ for all $i \in I$ and $x, y \in R$.

Definition 2.10 [8]: A fuzzy set μ in a near-ring R is called a fuzzy ideal of R if it satisfies:

- (i) $\mu(x - y) \geq \min\{\mu(x), \mu(y)\}$
- (ii) $\mu(y + x - y) \geq \mu(x)$
- (iii) $\mu(xy) \geq \mu(y)$
- (iv) $\mu((x + z)y - xy) \geq \mu(z)$ for all $x, y, z \in R$.

Definition 2.11 [7]: A fuzzy set μ in a near-ring R is an anti-fuzzy ideal of R if it satisfies:

- (i) $\mu(x - y) \leq \max\{\mu(x), \mu(y)\}$
- (ii) $\mu(y + x - y) \leq \mu(x)$
- (iii) $\mu(xy) \leq \mu(y)$
- (iv) $\mu((x + z)y - xy) \leq \mu(z)$ for all $x, y, z \in R$.

Definition 2.12 [12]: Let R be a ring. An IFS $A = \{ \langle x, \mu_A(x), \gamma_A(x) \rangle : x \in R \}$ of R is said to be **intuitionistic fuzzy subring** of R (In short IFSR) of R if

- (i) $\mu_A(x - y) \geq \text{Min} \{ \mu_A(x), \mu_A(y) \}$
- (ii) $\mu_A(xy) \geq \text{Min} \{ \mu_A(x), \mu_A(y) \}$
- (iii) $\gamma_A(x - y) \leq \text{Max} \{ \gamma_A(x), \gamma_A(y) \}$
- (iv) $\gamma_A(xy) \leq \text{Max} \{ \gamma_A(x), \gamma_A(y) \}$, for all $x, y \in R$

Definition 2.13 [12]: Let R be a ring. An IFSR A of R is said to be intuitionistic fuzzy normal subring (in short IFNSR) of R if

- (i) $\mu_A(xy) = \mu_A(yx)$
- (ii) $\gamma_A(xy) = \gamma_A(yx)$, for all $x, y \in R$

Definition 2.14 [15]: An IFS $A = \{ \langle x, \mu_A(x), \gamma_A(x) \rangle : x \in R \}$ of a ring R said to be

(a) **intuitionistic fuzzy Left Ideal** of R (In short IFLI) of R if

- (i) $\mu_A(x - y) \geq \text{Min} \{ \mu_A(x), \mu_A(y) \}$
- (ii) $\mu_A(xy) \geq \mu_A(y)$
- (iii) $\gamma_A(x - y) \leq \text{Max} \{ \gamma_A(x), \gamma_A(y) \}$
- (iv) $\gamma_A(xy) \leq \gamma_A(y)$, for all $x, y \in R$

(b) **intuitionistic fuzzy Right Ideal** of R (In short IFRI) of R if

- (i) $\mu_A(x - y) \geq \text{Min} \{ \mu_A(x), \mu_A(y) \}$
- (ii) $\mu_A(xy) \geq \mu_A(x)$
- (iii) $\gamma_A(x - y) \leq \text{Max} \{ \gamma_A(x), \gamma_A(y) \}$
- (iv) $\gamma_A(xy) \leq \gamma_A(x)$, for all $x, y \in R$

(c) **intuitionistic fuzzy Ideal** of R (In short IFI) of R if

- (i) $\mu_A(x - y) \geq \text{Min} \{ \mu_A(x), \mu_A(y) \}$
- (ii) $\mu_A(xy) \geq \text{Max} \{ \mu_A(x), \mu_A(y) \}$
- (iii) $\gamma_A(x - y) \leq \text{Max} \{ \gamma_A(x), \gamma_A(y) \}$
- (iv) $\gamma_A(xy) \leq \text{Min} \{ \gamma_A(x), \gamma_A(y) \}$, for all $x, y \in R$

Theorem 2.15 [15]: If $A = \{ \langle x, \mu_A(x), \square A(x) \rangle : x \in R \}$ be IFSR of ring R , then

- (i) $\mu_A(0) \geq \mu_A(x)$ and $\gamma_A(0) \leq \gamma_A(x)$
- (ii) $\mu_A(-x) = \mu_A(x)$ and $\gamma_A(-x) = \gamma_A(x)$, for all $x, y \in R$
- (iii) If R is ring with unity 1, then $\mu_A(1) \leq \mu_A(x)$ and $\gamma_A(1) \geq \gamma_A(x)$, for all $x \in R$

Definition 2.16 [11]: An intuitionistic fuzzy groupoid, denoted by $((X, I, I), F)$, is an IFS (X, I, I) together with an intuitionistic fuzzy binary operation F defined over it. An intuitionistic fuzzy semigroup is an intuitionistic fuzzy groupoid that is associative. An intuitionistic fuzzy monoid is an intuitionistic fuzzy semigroup that admits an identity. An intuitionistic fuzzy group is an intuitionistic fuzzy monoid in which each intuitionistic fuzzy element has an inverse. An intuitionistic fuzzy group $((G, I, I), F)$ is called an abelian (commutative) intuitionistic fuzzy group if and only if for all $(x, I, I), (y, I, I) \in ((G, I, I), F)$ $(x, I, I) F (y, I, I) = (y, I, I) F (x, I, I)$.

Definition 2.17 [6]: A fuzzy set μ of G is called a fuzzy $M\Gamma$ -subgroup of G if it satisfies the following two conditions:

- (i) $\mu(x - y) \geq \min \{ \mu(x), \mu(y) \}$ and
- (ii) $\mu(a\alpha y) \geq \mu(y)$ for all $x, y \in G, a \in M$, and $\alpha \in \Gamma$.

In this section, M denotes a gamma near ring, and G stands for an $M\Gamma$ -group.

Definition 2.18 [14]: Let M be a Γ -near ring. An additive group G is said to be a Γ -near ring module (or $M\Gamma$ -module) if there exists a mapping $M \times \Gamma \times G \rightarrow G$ (denote the image of (m, α, g) by $m\alpha g$ for $m \in M, \alpha \in \Gamma, g \in G$) satisfying the conditions

- (i) $(m_1 + m_2) \alpha_1 g = m_1 \alpha_1 g + m_2 \alpha_1 g$ and
- (ii) $(m_1 \alpha_1 m_2) \alpha_2 g = m_1 \alpha_1 (m_2 \alpha_2 g)$, for all $m_1, m_2 \in M, \alpha_1, \alpha_2 \in \Gamma$, and $g \in G$.

Definition 2.19 [14]:

- (i) An additive subgroup H of G is said to be a $M\Gamma$ -subgroup if $m\alpha h \in H$ for all $m \in M, \alpha \in \Gamma$, and $h \in H$. (Note that (0) and G are the trivial $M\Gamma$ -subgroups.)
- (ii) A normal subgroup H of G is said to be a submodule (or ideal) of G if $m\alpha (g + h) - m\alpha g \in H$ for $m \in M, \alpha \in \Gamma, g \in G$, and $h \in H$.
- (iii) For two $M\Gamma$ -modules G_1 and G_2 , a group homomorphism $\theta: G_1 \rightarrow G_2$ is said to be a module homomorphism (or Γ -module homomorphism) if $\theta(m\alpha g) = m\alpha(\theta g)$ for $m \in M, \alpha \in \Gamma$, and $g \in G_1$.

INTUITIONISTIC FUZZY IDEALS OF M- Γ GROUPS:

Definition 3.1: An intuitionistic fuzzy set $A (\mu_A, \gamma_A)$ of G is called an intuitionistic fuzzy $M\Gamma$ subgroup of G if it satisfies the following two conditions.

- a. $\mu(x - y) \geq \min \{ \mu(x), \mu(y) \}$
 $\gamma(x - y) \leq \max \{ \gamma(x), \gamma(y) \}$
- b. $\mu(a \alpha y) \geq \mu(y)$
 $\gamma(a \alpha y) \leq \gamma(y)$ for all $x, y \in G, a \in M$ and $\alpha \in \Gamma$.

Where M denotes a gamma near ring and G stands for an $M\Gamma$ group.

Definition 3.2: If G is said to be an intuitionistic fuzzy ideal if $M: G \rightarrow [0, 1]$ and $\gamma: G \rightarrow [0, 1]$ satisfying the following properties.

- i. $\mu(x + y) \geq \min \{ \mu(x), \mu(y) \}$
- ii. $\mu(x + y - x) \geq \mu(y)$
- iii. $\mu(x) = \mu(-x)$
- iv. $\mu(n\alpha(a + x) - n\alpha a) \geq \mu(x)$
- v. $\gamma(x + y) \leq \max \{ \gamma(x), \gamma(y) \}$
- vi. $\gamma(x + y - x) \leq \gamma(y)$
- vii. $\gamma(x) = \gamma(-x)$
- viii. $\gamma(n\alpha(a + x) - n\alpha a) \leq \gamma(x)$ for all $n \in M, \alpha \in \Gamma, a, x, y \in G$.

Remark 3.3: If μ and γ satisfies (i), (ii), (iii), (v), (vi), (vii) then $A (\mu_A, \gamma_A)$ is the intuitionistic fuzzy normal $M\Gamma$ subgroup of G .

Definition 3.4: For any family $\{A_i \in i \quad J\}$ of intuitionistic fuzzy of a set S , we define the intersection of the intuitionistic fuzzy sets A_i for $i \in J$ as

$$\begin{aligned} \mu(x) &= \min \{ \mu_i(x) / x \in A_i \} \\ \gamma(x) &= \min \{ \gamma_i(x) / x \in A_i \} \end{aligned}$$

$A_i (\mu_{A_i}, \gamma_{A_i})$ is an intuitionistic fuzzy subset of a set S .

Result 3.5: If $A (\mu_A, \gamma_A)$ is an intuitionistic fuzzy ideal of $M\Gamma$ group G then

- a. $\mu(g - g^1) \geq \min \{ \mu(g), \mu(g^1) \}$
 $\gamma(g - g^1) \leq \max \{ \gamma(g), \gamma(g^1) \}$
- b. $\mu(g + g^1) = \mu(g^1 + g)$
 $\gamma(g + g^1) = \gamma(g^1 + g)$ for all $g, g^1 \in G$.

Proposition 3.6: If $A (\mu_A, \gamma_A)$ is an intuitionistic fuzzy of the $M\Gamma$ group G for $x, y \in G$ with $\mu(x) > \mu(y)$ and $\gamma(x) > \gamma(y)$ then, $\mu(x + y) = \mu(y)$ and $\gamma(x + y) = \gamma(x)$.

Moreover, if $x, y \in G$, $\mu(x) \neq \mu(y)$ and $\gamma(x) \neq \gamma(y)$ then,

$$\begin{aligned} \mu(x + y) &= \min \{ \mu(x), \mu(y) \} \\ \gamma(x + y) &= \max \{ \gamma(x), \gamma(y) \} \end{aligned}$$

Proof: Suppose $\mu(x) > \mu(y)$ and $\gamma(x) > \gamma(y)$

By definition,

$$\begin{aligned} \mu(x + y) &\geq \min \{ \mu(x), \mu(y) \} \\ &= \mu(y) \text{ (since } \mu(x) > \mu(y) \text{)} \\ \Rightarrow \mu(x + y) &\geq \mu(y) \end{aligned} \tag{1}$$

$$\begin{aligned} \mu(y) &= \mu(y + x - x) \\ &\geq \min \{ \mu(y + x), \mu(-x) \} \text{ [since } \mu(-x) = \mu(x) \text{]} \\ &= \min \{ \mu(y + x), \mu(x) \} \\ &= \mu(x) \text{ if } \mu(y + x) > \mu(x) \\ \Rightarrow \mu(y) &\geq \mu(x) \text{ which is a contradiction.} \end{aligned} \tag{2}$$

Therefore, $\mu(y + x) < \mu(x)$.

Substituting in 2

$$\mu(y) \geq \mu(x + y) \tag{3}$$

1 and 3 $\Rightarrow \mu(x + y) = \mu(y)$

Consider $\gamma(x + y) \leq \max \{ \gamma(x), \gamma(y) \}$
 $\leq \gamma(x)$ [since $\gamma(x) > \gamma(y)$]

i.e. $\gamma(x + y) \leq \gamma(x)$ (4)

Consider $\gamma(x) = \gamma(x + y - y)$
 $\leq \max \{ \gamma(x + y), \gamma(-y) \}$
 $= \max \{ \gamma(x + y), \gamma(y) \}$ [since $\gamma(-y) = \gamma(y)$] (5)

If $\gamma(x + y) < \gamma(y)$
 $\Rightarrow \gamma(x) \leq \gamma(y)$ which is a contradiction to the given data.

Therefore, $\gamma(x + y) > \gamma(y)$
5 $\Rightarrow \gamma(x) \leq \gamma(x + y)$ (6)
4 and 6 $\Rightarrow \gamma(x) = \gamma(x + y)$

This completes the preposition

Theorem 3.7: Let M be a zero symmetric Γ near ring and G be a $M\Gamma$ group. If A (μ_A, γ_A) is an intuitionistic fuzzy ideal of G then $\mu(m \alpha x) \geq \mu(x)$ for all $x \in G, m \in M$ and $\alpha \in \Gamma$ and $\gamma(m \alpha x) \leq \gamma(x)$

Proof: Consider $\mu(m \alpha x) = \mu(m \alpha (x + 0))$
 $= \mu(m \alpha (0 + x - 0))$
 $= \mu(m \alpha (0 + x) - m \alpha)$
 $\geq \mu(x)$

Since M is a zero symmetric Γ near ring.

Similarly, $\gamma(m \alpha x) = \gamma(m \alpha (x + 0))$
 $= \gamma(m \alpha (0 + x) - 0)$
 $= \gamma(m \alpha (0 + x) - m \alpha)$
 $\leq \gamma(x)$

Thus, the theorem is proved.

Remark 3.8: If A (μ_A, γ_A) is an intuitionistic fuzzy ideal then

- $\mu(0) \geq \mu(g)$ and $\gamma(1) \leq \gamma(g)$ for all $g \in G$.
- $\mu(0) = \sup_{g \in G} \mu(g)$
 $\gamma(1) = \sup_{g \in G} \gamma(g)$.

As A (μ_A, γ_A) is an intuitionistic fuzzy, the image of A under $\mu, \mu(A)$ is a subset of $[0, \mu(0)]$ and the image of A under $\gamma, \gamma(A)$ is a subset of $[\gamma(1), 1]$.

Examples of Intuitionistic Fuzzy Ideal 3.9:

- Every constant intuitionistic fuzzy set A (μ_A, γ_A) where $\mu_A: A \rightarrow [0, 1]$ and,
- $\gamma_A: A \rightarrow [0, 1]$ as $\mu_A(x) = 0.6$ and $\gamma_A(x) = 0.2$ for all $x \in A$ is an intuitionistic fuzzy ideal of G.
- Take $N = Z$ with usual operations '+' and '.'. Then $(N, +, \cdot)$ forms a near ring. Let $\Gamma = \{.\}$ and $M = N$. then M is a near ring with $G = Z$. Then $(G, +)$ is a group of integers with usual addition. Now G is a N- group and a $M\Gamma$ group.

Let $\mu: A \rightarrow [0, 1]$ and $\gamma: A \rightarrow [0, 1]$ be defined by

$$\mu(x) = \begin{cases} 0.5 & \text{if } x = 4n \text{ for some } n \in Z \\ 0.3 & \text{if } x = 2n \text{ for some } n \in Z \text{ and } x \neq 4m \text{ for some } m \in Z \\ 0 & \text{if } x \text{ is odd} \\ 1 & \text{if } x = 0 \text{ and otherwise.} \end{cases}$$

$$\gamma(x) = \begin{cases} 0.3 & \text{if } x = 4n \text{ for some } n \in Z \\ 0.4 & \text{if } x = 2n \text{ for some } n \in Z \text{ and } x \neq 4m \text{ for some } m \in Z \\ 1 & \text{if } x \text{ is odd} \\ 0 & \text{if } x = 0 \text{ and otherwise.} \end{cases}$$

Here $0 \leq \mu(x) + \gamma(x) \leq 1$ for all $x \in A$ Then A satisfies all the axioms of intuitionistic fuzzy ideal of G.

Lemma 3.10: If A (μ_A, γ_A) is an intuitionistic fuzzy of the $M\Gamma$ group G and if $\mu(x - y) = \mu(0)$ and $\gamma(x - y) = \gamma(1)$ then $\mu(x) = \mu(y)$ and $\gamma(x) = \gamma(y)$ for all $x, y \in G$.

Proof: Assume $\mu(x - y) = \mu(0)$

Consider $\mu(x) = \mu(x - y + y)$
 $\geq \min\{\mu(x - y), \mu(y)\}$
 $= \min\{\mu(0), \mu(y)\}$ by * our assumption.
 $= \mu(y)$
 $\mu(x) \geq \mu(y)$ -----> **I**

On the other hand,

$$\mu(y) = \mu(y - x + x)$$

$$\geq \min\{\mu(y - x), \mu(x)\} [\mu(y - x) = \mu(-(x - y))]$$

$$= \mu(x - y)$$

$$= \min\{\mu(0), \mu(x)\}$$

$$= \mu(x) \quad [\mu(0) = 1, \gamma(1) = 0]$$

$$\mu(y) \geq \mu(x)$$
 -----> **II**

I & II $\Rightarrow \mu(x) = \mu(y)$

Assume $\gamma(x - y) = \gamma(1)$
 $= \gamma(x - y + y)$

Consider, $\gamma(x) = \gamma(x - y + y)$
 $\leq \max\{\gamma(x - y), \gamma(y)\}$
 $\leq \max\{\gamma(1), \gamma(y)\}$ by our assumption
 $= \gamma(y)$
 $\gamma(x) \leq \gamma(y)$ -----> **III**

On the other hand,
 $\gamma(y) = \gamma(y - x + x)$
 $\leq \max\{\gamma(y - x), \gamma(x)\}$ [$\gamma(y - x) = \gamma(-(x - y))$]
 $= \max\{\gamma(1), \gamma(x)\} = \gamma(x - y)$
 $= \gamma(x)$
 $\gamma(y) \leq \gamma(x)$ -----> **IV**

III & IV $\Rightarrow \gamma(x) = \gamma(y)$

The converse of the above lemma 3.10 is not true. That is, we can find an $M\Gamma$ group G and a intuitionistic fuzzy ideal H of G such that $\mu(x) = \mu(y)$ and $\gamma(x) = \gamma(y)$ but $\mu(x - y) \neq 0$ and $\gamma(x - y) \neq 0$. For this consider the following example:

Consider the example of intuitionistic fuzzy ideal.

Let $x = 3, y = 7$ then by μ, γ defined
 $\mu(3) = 0$ and $\mu(7) = 0 \Rightarrow \mu(x) = \mu(y)$

But $\mu(x - y) = \mu(3 - 7) = \mu(-4) = 0.5 \neq 1 = \mu(0)$
 And $\gamma(3) = 1$ and $\gamma(7) = 1 \Rightarrow \gamma(x) = \gamma(y)$

But $\gamma(x - y) = \gamma(3 - 7) = 0.3 \neq 0 = \gamma(1)$.

Theorem 3.11: Let G be an $M\Gamma$ group an $A (\mu_A, \gamma_A)$ be an intuitionistic fuzzy subset of G . then the following conditions are equivalent.

1. A is an intuitionistic fuzzy ideal of the $M\Gamma$ group G and
2. $U(\mu_A, t)$ and $L(\gamma_A, s)$ are ideals of the $M\Gamma$ group G for all $s, t \in [0, 1]$.

Proof:

1 \Rightarrow 2: Let $s, t \in [0, 1]$

Let $x, y \in U(\mu_A, t) \Rightarrow \mu_A(x) > t$ and $\mu_A(y) > t$.
 $\mu_A(x - y) \geq \min\{\mu_A(x), \mu_A(-y)\}$
 $= \min\{\mu_A(x), \mu_A(y)\}$
 $= \min\{t, t\} = t$
 $\Rightarrow \mu_A(x - y) \geq t$
 $\Rightarrow x - y \in U(\mu_A, t)$.
 $\Rightarrow \mu_A(t)$ is a subgroup of $(G, +)$

Let $g \in G$.

Consider, $\mu_A(g + x - g) \geq \mu_A(x) \geq t$
 $\Rightarrow g + x - g \in U(\mu_A, t)$
 $\Rightarrow U(\mu_A, t)$ is a normal subgroup of $(G, +)$.

Let $m \in M$ and $\alpha \in \Gamma$.

Consider, $\mu_A(m \alpha (g + x) - m \alpha g) \geq \mu_A(x) \geq t$
 $\Rightarrow m \alpha (g + x - m \alpha g) \in U(\mu_A, t)$
 $\Rightarrow U(\mu_A, t)$ is an intuitionistic fuzzy of the $M\Gamma$ group G .

Similarly, $L(\gamma_A, t)$ is an intuitionistic fuzzy of the $M\Gamma$ group G .

Let $x, y \in L(\gamma_A, s) \Rightarrow \gamma_A(x) < s$ and $\gamma_A(y) < s$.

Therefore, $\gamma_A(x - y) \leq \max\{\gamma_A(x), \gamma_A(-y)\}$
 $= \max\{\gamma_A(x), \gamma_A(y)\}$
 $= \max\{s, s\} = s$
 $\Rightarrow \gamma_A(x - y) \leq s$
 $\Rightarrow x - y \in L(\gamma_A, s)$.
 $\Rightarrow L(\gamma_A, s)$ is a subgroup of $(G, +)$

Let $g \in G$. Consider

$\gamma_A(g + x - g) \leq \gamma_A(x) \leq s$
 $\Rightarrow g + x - g \in L(\gamma_A, s)$
 $\Rightarrow L(\gamma_A, s)$ is a normal subgroup of $(G, +)$.

Let $m \in M$ and $\alpha \in \Gamma$.

Consider $\gamma_A(m \alpha (g + x) - m \alpha g) \leq \gamma_A(x) \leq s$.
 $\Rightarrow m \alpha (g + x) - m \alpha g \in L(\gamma_A, s)$
 $\Rightarrow L(\gamma_A, s)$ is an intuitionistic fuzzy ideal of the $M\Gamma$ group G .

2 \Rightarrow 1 Assume $U(\mu_A, t)$ and $L(\gamma_A, t)$ are the ideals of the $M\Gamma$ group of the group G .

Let $x, y \in U(\mu_A, t) \Rightarrow \mu_A(x) \geq t$ and $\mu_A(y) \geq t$

With $\mu_A(x - y) \geq t$ as $U(\mu_A, t)$ is an intuitionistic fuzzy ideal.

As $x \in U(\mu_A, t)$

$\Rightarrow g + x - g \in U(\mu_A, t)$ for all $g \in G$. Since $U(\mu_A, t)$ is a normal subgroup.
 $\Rightarrow \mu_A(g + x - g) \geq t$.

To prove A is intuitionistic fuzzy let us check all the conditions of definition 2.

(*) Let $x, y \in U(\mu_A, t)$ such that

$$\mu_A(x - y) < \min\{\mu_A(x), \mu_A(y)\}$$

Take $t = [\mu(x - y) + \min\{\mu_A(x), \mu_A(y)\}] / 2$

Then $0 \leq \mu(x - y) < t < \min\{\mu_A(x), \mu(y)\} \leq 1$
 \Rightarrow for $x, y \in U(\mu_A, t)$ which is a contradiction to $U(\mu_A, t)$ is an ideal.

Therefore $\mu_A(x - y) \geq t$

$\Rightarrow \mu_A(x + y) = \mu_A(x - (-y))$
 $= \min\{\mu_A(x), \mu_A(-y)\}$
 $= \min\{\mu_A(x), \mu_A(y)\}$
 $\geq \min\{t, t\} = t$
 $\Rightarrow \mu_A(x + y) \geq t$ -----> **(i)**

(**) Let $x, y \in G$ such that $\mu_A(y + x - y) < \mu_A(x)$

Take $t = [\mu_A(y + x - y) + \mu_A(x)] / 2$

$\Rightarrow 0 \leq \mu_A(y + x - y) < t < \mu_A(x) \leq 1$
 \Rightarrow for $x, y \in U(\mu_A, t)$, $y + x - y$ does not belong to $U(\mu_A, t)$ which is a contradiction to $U(\mu_A, t)$ is an ideal
 $\Rightarrow \mu_A(y + x - y) \geq \mu(x)$ -----> **(ii)**

(***) Let $y \in G$ and $x \in U(\mu_A, t)$ such that

$$\mu_A(xy) < \mu_A(x)$$

Take $t = [\mu_A(xy) + \mu_A(x)] / 2$

$\Rightarrow 0 \leq \mu_A(x) < t < \mu_A(xy) \leq 1$
 \Rightarrow for $x \in U(\mu_A, t)$, xy does not belong to $U(\mu_A, t)$ which is a contradiction to $U(\mu_A, t)$ is an ideal.

Therefore $\mu_A(xy) > \mu_A(x)$ -----> (iii)

(****) Let $x, y, z \in G$ such that
 $\mu_A[(x+z)y - xy] < \mu_A(z)$

Take $t = \{\mu_A[(x+z)y - xy] + \mu_A(z)\} / 2$
 $\Rightarrow 0 \leq \mu_A[(x+z)y - xy] < t < \mu_A(z) \leq 1$
 \Rightarrow for $z \in U(\mu_A, t)$ but $(x+z)y - xy$ does not belongs to $U(\mu_A, t)$ which is a contradiction to $U(\mu_A, t)$ is an ideal.
 $\Rightarrow \mu_A[(x+z)z - xy] \geq \mu_A(z)$ -----> (iv)

From (i), (ii), (iii), (iv) μ_A satisfies first four conditions of definition 2.

Similarly, γ_A satisfies the last four conditions of definition 2.

Hence A is an intuitionistic fuzzy ideal of the $M\Gamma$ group.

Proposition 3.12: Let H be an intuitionistic fuzzy ideal of the $M\Gamma$ group G and $A(\alpha_1, \beta_1)$ and $B(\alpha_2, \beta_2)$ are the two-level ideals of H . Then the following two conditions are equivalent.

- $A(\alpha_1, \beta_1) = B(\alpha_2, \beta_2)$ and
- There is no $x \in G$ such that $\alpha_1 < \mu(x) < \alpha_2$ and $\beta_1 > \gamma(x) > \beta_2$.

Proof:

a \Rightarrow b

let us assume there exists an element $x \in G$ such that

$\alpha_1 < \mu(x) < \alpha_2$ and $\beta_1 > \gamma(x) > \beta_2$
 $\Rightarrow \mu(x) > \alpha_1$ and $\gamma(x) < \beta_1$ and $\mu(x) < \alpha_2$ and $\gamma(x) > \beta_2$
 $\Rightarrow x \in A(\alpha_1, \beta_1)$ and x does not belongs to $B(\alpha_2, \beta_2)$ which is a contradiction to (i). Hence our assumption is wrong and so there is no $x \in G$ such that $\alpha_1 < \mu(x) < \alpha_2$ and $\beta_1 > \gamma(x) > \beta_2$.

b \Rightarrow a

Let $x \in B(\alpha_2, \beta_2)$

$\Rightarrow \mu(x) > \alpha_2$ and $\gamma(x) < \beta_2$

Since $\alpha_2 > \alpha_1$ and $\beta_2 < \beta_1$ we have $\mu(x) > \alpha_1$ and $\gamma(x) < \beta_1$
 $\Rightarrow x \in B(\alpha_1, \beta_1)$

Hence $B(\alpha_2, \beta_2) \subset B(\alpha_1, \beta_1)$ -----> I

Let $x \in B(\alpha_1, \beta_1) \Rightarrow \mu(x) > \alpha_1$ and $\gamma(x) < \beta_1$

By our assumption there is no x such that

$\alpha_1 < \mu(x) < \alpha_2$ and $\beta_1 > \gamma(x) > \beta_2$
 $\Rightarrow \mu(x)$ does not less than α_2 and $\gamma(x)$ does not greater than β_2 .
 $\Rightarrow \mu(x) > \alpha_2$ and $\gamma(x) < \beta_2$
 $\Rightarrow x \in B(\alpha_2, \beta_2)$

And so, on $B(\alpha_1, \beta_1) \subset B(\alpha_2, \beta_2)$ -----> II

From I & II

$B(\alpha_1, \beta_1) = B(\alpha_2, \beta_2)$ and hence (ii) \Rightarrow (i)

Hence the Proof is completed.

CONCLUSION

We defined Intuitionistic Fuzzy Ideals of $M-\Gamma$ Groups by extending its Fuzzy Ideals and discussed equivalent conditions on Intuitionistic Fuzzy Γ Groups. We have also explained the properties and illustrated an example to prove Intuitionistic Fuzzy Ideals of $M-\Gamma$ Groups.

REFERENCES

1. Atanassov, K. T. (1986). Intuitionistic fuzzy sets. *Fuzzy sets and Systems*, 20(1), 87-96.
2. Batulan, R. A., & Veeramani, V. (1991). Some characterisations of alpha cut in Intuitionistic fuzzy set theory.
3. Clay, J. (1993). *Nearrings*. Oxford: Oxford University Press.
4. Dubois, D., & Prade, H. (2012). *Fundamentals of fuzzy sets (Vol. 7)*. Springer Science & Business Media.
5. Jianming, Z., & Xueling, M. (2005). Intuitionistic fuzzy ideals of near-rings. *International Society for Mathematical Sciences*, 61, 219-223.
6. Jun Y.B., Kwon Y.I., and Park J.W. (1995). Fuzzy $M\Gamma$ -groups, *Kyungpook Mathematical Journal*. 35, 259-265.
7. Kim, K.B, Jun, Y.B, & Yon, Y.H (2001). Abnormalization of anti-fuzzy ideals in near-rings. *Far East Journal of Mathematical Sciences*, 15-25.
8. Kim, S. D., & Kim, H. S. (1996). On fuzzy ideals of near-rings. *Bull. Korean Mathematical Society*, 33(4), 593-601.
9. Klir, G. J., & Yuan, B. (1996). *Fuzzy Sets and Fuzzy Logic: Theory and Applications*.
10. Ma, X., Zhan, J., & Leoreanu-Fotea, V. (2013). On (fuzzy) isomorphism theorems of soft Γ -hypermoudles. *UPB Scientific Bulletin*, 75(1), 65-76.
11. Marashdeh, M. F., & Salleh, A. R. (2011). Intuitionistic fuzzy rings. *International Journal of Algebra*, 5(1), 37-47.
12. Meena, K., & Thomas, K. V. (2011). Intuitionistic L-fuzzy subrings. In *International Mathematical Forum Vol. 6, No. 52*, pp. 2561-2572.
13. Palaniappan, N& Ramachandran, M. (2010). A Note on Characterization of Intuitionistic Fuzzy Ideals in Γ -Rings. *International Mathematical Forum*. 5. 2553-2562.
14. Satyanarayana, B., & Prasad, K. S. (2013). Near rings, fuzzy ideals, and graph theory. *CRC press*, 305 – 312, 335 – 344.
15. Sharma, P. K. (2011). Translates of Intuitionistic fuzzy subring. *International Review of Fuzzy Mathematics*, 6(2), 77-84.
16. Sharma, P. K. (2013). Cut of Intuitionistic fuzzy modules-II. *International Journal of Mathematical Sciences and Applications*, 3(1), 11-17.
17. Sharma, P. K. (2015). Intuitionistic fuzzy representations of intuitionistic fuzzy groups. *Asian Journal of Fuzzy and Applied Mathematics*.
18. Sharma, P. K., & Bahl, A. (2012). Translates of Intuitionistic fuzzy submodules. *International Journal of Mathematics Sciences*, 11(3-4), 277-287.
19. Zadeh, L. A. (1996). Fuzzy sets. In *Fuzzy Sets, Fuzzy Logic, And Fuzzy Systems*, 338-353.

Source of support: Proceedings of UGC Funded International Conference on Intuitionistic Fuzzy Sets and Systems (ICIFSS-2018), Organized by: Vellalar College for Women (Autonomous), Erode, Tamil Nadu, India.