

NOTE ON COMPLEMENT GRAPHS

P. SREENIVASULU REDDY*, MELAKU ABERA EJERSA

Department of Mathematics,
Samara University, Semera, Afar Regional State, Ethiopia. Post Box No.131.

(Received On: 10-12-17; Revised & Accepted On: 03-01-18)

ABSTRACT

In this article we generalize the complement for all types of graphs and we proved some primary results on complement of graphs.


Key words: Subgraph, Complement of graph.

1. INTRODUCTION

There are various types of graphs depending upon the number of vertices, number of edges, interconnectivity, and their overall structure. We will discuss only a complement of graphs in this chapter. Note that the edges in graph-I are not present in graph-II and vice versa. Hence, the combination of both the graphs gives a final graph of 'n' vertices.

1.1. Definition: A graph 'G' is defined as $G = (V, E)$ Where V is a set of all vertices and E is a set of all edges in the graph.

Example:



In the above example, AB, AC, CD, and BD are the edges of the graph. Similarly, A, B, C, and D are the vertices of the graph.

1.2. Definition: A graph having no edges is called a Null Graph.

Example:




1.3. Definition: A simple graph with 'n' mutual vertices is called a complete graph and it is denoted by ' K_n '. In the graph, a vertex should have edges with all other vertices, and then it is called a complete graph.

In other words, if a vertex is connected to all other vertices in a graph, then it is called a complete graph.

Example: In the following graphs I and II, each vertex in the graph is connected with all the remaining vertices in the graph except by itself

Corresponding Author: P. Sreenivasulu Reddy*,
Department of mathematics, Samara University,
Semera, Afar Regional State, Ethiopia. Post Box No.131.




1.4. Definition: Let H be a graph with vertex set $V(H)$ and edge set $E(H)$, and similarly let G be a graph with vertex set $V(G)$ and edge set $E(G)$. Then, we say that H is a subgraph of G if $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$. In such a case, we also say that G is a supergraph of H.

Example: Last four graphs are subgraphs for first graph in the below diagram.




1.5. Definition: The union of 2 simple graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ is the simple graph $G = (V, E)$ with vertex set $V = V_1 \cup V_2$ and edge set $E = E_1 \cup E_2$. The union is denoted by $G_1 \cup G_2$.

Example:




1.6. Definition: The intersection of graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ is the graph $G = (V, E)$ with vertex set $V = V_1 \cap V_2$ and edge set $E = E_1 \cap E_2$. The intersection is denoted by $G_1 \cap G_2$.

Example:



1.7. Definition: The complement of a graph G is the graph having the same vertex set as G such that two vertices are adjacent if and only if the same two vertices are non-adjacent in G. We denote the complement of a graph G by G^c or G^1 or \overline{G} .

Example:



1.8. Notation: Since the complete graph on n vertices has nc_2 edges, it follows that if G is a graph on n vertices with m edges, then G^c is also a graph on n vertices but with $nc_2 - m$ edges.

2. OUR APPROACH ON COMPLEMENT OF GRAPHS


2.1. Lemma: Let G be a graph and H is a subgraph of G with the same vertices set of G then H^c is also subgraph of G.

The subgraphs with the same vertices set of graph G contains only its complement. So every subgraph of graph G has may not have its complement.

2.2. Theorem: Let G be a graph and H is a subgraph of G with the same vertices set of G then $G = H \cup H^c$

Proof: Let G be a graph with vertices set V and edges set E. H be a subgraph of G with vertices set V and edges set E_1 then H^c has with the same vertices set V and edges set E_2 which is different from E_1 . Since all the edges of E contained in any one of the edge sets E_1 and E_2 , $E = E_1 \cup E_2$. Therefore, $G = H \cup H^c$.


Example:



2.3. Theorem: Let G be a graph and H is a subgraph of G with the same vertices set of G then $H \cap H^c$ is null graph

Proof: Let G be a graph with vertices set V and edges set E. H be a subgraph of G with vertices set V and edges set E_1 then H^c has with the same vertices set V and edges set E_2 which is different from E_1 implies $E_1 \cap E_2$ is empty set. So, $H \cap H^c$ has vertices set V and no edges. $H \cap H^c$ is null graph.

Example: Consider the above example



2.4. Notations:

- 1) The complement of graph G is null graph and null graph complement is graph G. G and null graphs are improper subgraphs of graph G.
- 2) Let G be a graph and H is a subgraph of G with the same vertices set of G then $(H^c)^c = H$.
- 3) Let G be graph with subgraphs $H_1, H_2, H_3, \dots, H_n$ then $(H_1 \cup H_2 \cup H_3 \cup \dots \cup H_n)^c = H_1^c \cap H_2^c \cap H_3^c \cap \dots \cap H_n^c$
- 4) Let G be graph with subgraphs $H_1, H_2, H_3, \dots, H_n$ then $(H_1 \cap H_2 \cap H_3 \cap \dots \cap H_n)^c = H_1^c \cup H_2^c \cup H_3^c \cup \dots \cup H_n^c$.
- 5) Observe that the trivial graph on 1 vertex and no edges is clearly self-complement.

Similarly, we can prove the following theorems

2.5. Theorem: Let G be graph with all possible n proper subgraphs $H_1, H_2, H_3, \dots, H_n$ then $G = H_1^c \cup H_2^c \cup H_3^c \cup \dots \cup H_n^c$.

2.6. Theorem: Let G be graph withal possible n proper subgraphs $H_1, H_2, H_3, \dots, H_n$ then $H_1^c \cap H_2^c \cap H_3^c \cap \dots \cap H_n^c$ is null graph.

REFERENCES

1. J. Akiyama, K. Ando and F. Harary, A graph and its complement with specified properties. VIII. Interval graphs, Math. Japon. 29 (1984) 659-670. MR 86b:05061.
2. J. Akiyama and F. Harary, A graph and its complement with specified properties. I. Connectivity, Internat. J. Math. Math. Sci. 2 (1979a) 223-228. MR 82i:05062a.
3. M. Aigner, Graphs whose complement and line graph are isomorphic, J. Combin. Theory 7 (1969) 273{275. MR 39:6765.
4. M. Chudnovsky, N. Robertson, P. Seymour, R. Thomas, The strong perfect graph theorem,
5. Annals of Mathematics 164 (1) (2006) 51-229.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]