

$G^*\alpha$ -LOCALLY CLOSED SETS AND $G^*\alpha$ -LOCALLY CLOSED FUNCTIONS

Dr. A. PUNITHA THARANI
 Associate professor, St. Mary's College, Thoothukudi, India.

T. DELCIA*
 Research Scholar, Manonmaniam Sundaranar University, Tirunelveli, India.

(Received On: 20-01-18; Revised & Accepted On: 15-02-18)

ABSTRACT

The purpose of this paper is to introduce the concepts of g^α -locally closed sets and $g^*\alpha$ -locally closed functions. We investigate their basic properties. We also discuss their relationship with already existing concepts.*

INTRODUCTION

The notion of locally closed sets in topological space was introduced by Bourbaki [3]. Ganster and Reilly [6] further studied the properties of locally closed sets and defined the LC-continuity and LC-irresoluteness. In Literature many general topologists introduced the studies of locally closed sets. Balachandran *et al.* [1] introduced the concept of generalized locally closed sets and seven different notions of generalized continuities. In this paper we continue the study of generalizations locally closed sets and investigate the classes of $G^*\alpha$ -Locally closed functions and study some of their properties.

PRELIMINARIES

Throughout this paper (X, τ) denotes a topological space with a topology τ on which no separation axioms are assumed unless otherwise mentioned. For a subset A of a space (X, τ) , $cl(A)$, $int(A)$, A^c , $P(X)$ denote the closure of A , the interior of A , the complement of A , the power set of X . We recall the following Definitions, Remarks, Corollary and Theorem which are prerequisite for this paper.

Definition 2.1: A subset A of a topological space (X, τ) is called

- (1) a semi-open set [9] if $A \subseteq cl(int(A))$ and a semiclosed set if $int(cl(A)) \subseteq A$
- (2) an α -open set if [12] $A \subseteq int(cl(int(A)))$ and a α -closed set if $cl(int(cl(A))) \subseteq A$
- (3) a regular open set [16] if $int(cl(A)) = A$ and regular closed set if $A = int(cl(A))$

Definition 2.2: A subset A of a topological space (X, τ) is called

- (1) generalized closed (briefly g -closed) set [8] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is open; the complement of g -closed set is g -open set.
- (2) regular generalised closed set (briefly rg -closed) [13] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and regular open in (X, τ) ; the complement of rg -closed set is rg -open set.
- (3) α -generalised closed set (briefly αg -closed) [10] if $\alpha cl(A) \subseteq U$ whenever $A \subseteq U$ and U is Open in (X, τ) ; the complement of αg -closed set is αg -open set.
- (4) g^\wedge -closed set [20] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is semi open in (X, τ) ; the complement of g^\wedge -closed set is g^\wedge -open set.
- (5) complement of g^\wedge -closed set is g^\wedge -open set.
- (6) g^* -closed set [22] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is g^* -open in (X, τ) ; the complement of g^* -closed set is g^* -open set.
- (7) $g^\#$ -closed set [18] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is αg -open in (X, τ) ; the complement of $g^\#$ -closed set is $g^\#$ -open set.
- (8) $g^*\alpha$ closed set if $\alpha cl(A) \subseteq U$ whenever $A \subseteq U$ and U is g^* open in X .

Corresponding Author: T. Delcia*
 Research Scholar, Manonmaniam Sundaranar University, Tirunelveli, India.

Definition 2.3: A subset A of a topological space (X, τ) is called

1. locally closed (briefly lc) [6] if $S = U \cap F$, where U is open and F is closed in (X, τ) .
2. generalized locally closed (briefly glc) [1] if $S = U \cap F$, where U is g -open and F is g -closed in (X, τ) .
3. generalized locally semi-closed (briefly glsc) [7] if $S = U \cap F$, where U is g -open and F is semi-closed in (X, τ) .
4. locally semi-closed (briefly lsc) [7] if $S = U \cap F$, where U is open and F is semi-closed in (X, τ) .
5. α -locally closed (briefly αlc) [7] if $S = U \cap F$, where U is α -open and F is α -closed in (X, τ) .
6. g^\wedge -locally closed set (briefly $g^\wedge lc$) [21] if $S = U \cap F$, where U is g^\wedge -open and F is g^\wedge -closed in (X, τ) .
7. $g^\#$ -locally closed (briefly $g^\# lc$) [19] if $S = U \cap F$, where U is $g^\#$ -open and F is $g^\#$ -closed in (X, τ) .
8. g^* -locally closed (briefly $g^* lc$) [22] if $S = U \cap F$, where U is g^* -open and F is g^* -closed in (X, τ) .

The class of all locally closed (resp. generalized locally closed, generalized locally semi-closed, locally semi-closed, g^\wedge -locally closed, $g^\#$ -locally closed, g^* -locally closed) sets in X is denoted by $LC(X)$ (resp. $GLC(X)$, $GLSC(X)$, $LSC(X)$, $G^\wedge LC$, GLC , $G^* LC$)

Definition 2.4: [14] A space (X, τ) is called a ${}_\alpha T_{1/2}^{**}$ -space if every $g^*\alpha$ -closed set is closed.

Recall that a subset A of a space (X, τ) is called dense if $cl(A) = X$

Definition 2.5: A subset A of a topological space (X, τ) is called

1. submaximal [5] if every dense subset is open.
2. g -submaximal [1] if every dense subset is g -open.
3. rg -submaximal [13] if every dense subset is rg -open.

Remark 2.6: For a topological space (X, τ) , the following statements hold:

1. Every closed set is $g^*\alpha$ -closed but not conversely [14]
2. Every g -closed set is $g^*\alpha$ -closed but not conversely [14]
3. Every g^* -closed set is $g^*\alpha$ -closed but not conversely [14]
4. Every α -closed set is $g^*\alpha$ -closed but not conversely [14]
5. A subset A of X is $g^*\alpha$ -closed iff $g^*\alpha-cl(A) = A$ [14]
6. A subset A of X is $g^*\alpha-int A$ iff $g^*\alpha-int(A) = A$ [14]
7. Every ${}_\alpha T_{1/2}^{**}$ -space is a $T_{1/2}$ -space [14]

Corollary 2.7: If A is $g^*\alpha$ -closed and F is closed, then $A \cap F$ is a $g^*\alpha$ -closed set.

3. $G^*\alpha$ -Locally closed Sets

We introduce the following definition.

Definition 3.1: A subset A of (X, τ) is called $g^*\alpha$ -locally closed (briefly $g^*\alpha$ -lc) if $A = S \cap G$, where S is $g^*\alpha$ -open and G is $g^*\alpha$ -closed in (X, τ) .

The class of all $g^*\alpha$ -locally closed sets in X is denoted by $G^*\alpha LC(X)$.

Proposition 3.2: Every $g^*\alpha$ -closed (resp $g^*\alpha$ -open) is $g^*\alpha$ lc but not conversely.

Proof: It follows from Definition 3.1

Example 3.3: Let $X = \{1, 2, 3\}$ with the topology $\tau = \{\emptyset, X, \{1\}, \{1, 2\}, \{1, 3\}\}$. Then the set $\{1\}$ is $g^*\alpha$ -lc but not $g^*\alpha$ -closed and the set $\{2\}$ is $g^*\alpha$ -lc but not $g^*\alpha$ -open in (X, τ) .

Proposition 3.4: Every lc set is $g^*\alpha lc$ but not conversely.

Proof: It follows from Remark 2.6(1)

Example 3.5: Let $X = \{1, 2, 3\}$ with the topology $\tau = \{\emptyset, X, \{2\}, \{2, 3\}\}$. Then the set $\{1, 2\}$ is $g^*\alpha$ -lc but not lc.

Proposition 3.6: Every glc^* set is $g^*\alpha lc$ but not conversely

Proof: It follows from Remark 2.6(2)

Example 3.7: Let $X = \{1, 2, 3, 4\}$ with the topology $\tau = \{\emptyset, X, \{1\}, \{1, 4\}, \{1, 2, 4\}\}$. Then the set $\{1, 3\}$ is $g^*\alpha$ -lc but not glc^* set in (X, τ)

Prop 3.8: Every alc is $g^*\alpha$ lc but not conversely

Proof: It follows from Remark 2.6(4)

Example 3.9: Let $X = \{1,2,3,4,5\}$ with the topology $\tau = \{\emptyset, X, \{1,2\}, \{3,4\}, \{1,2,3,4\}\}$. Then the set $\{2, 3, 4, 5\}$ is $g^*\alpha$ -lc but not alc

Prop 3.10: Every lsc and glsc are $g^*\alpha$ -lc but not conversely.

Proof: Proof follows from the Definition 2.3(3 &4)

Example 3.11: Let $X = \{1,2,3,4\}$ with the topology $\tau = \{\emptyset, X, \{1\}, \{2\}, \{1,2\}, \{1,2,3\}\}$. Then the set $\{1, 2, 4\}$ is $g^*\alpha$ -lc but not lsc and glsc

Prop 3.12: Every $g^*\text{lc}$ and $g^\# \text{lc}$ are $g^*\alpha$ -lc but not conversely.

Proof: Proof follows from Remark 2.6(3) and Definition 2.3(7)

Example 3.13: Let $X = \{1,2,3\}$ with the topology $\tau = \{\emptyset, X, \{2\}\}$. Then the set $\{1,2\}$ is $g^*\alpha$ -lc but not $g^*\text{lc}$ and $g^\# \text{lc}$

Prop 3.14: Every $g^\wedge \text{lc}$ is $g^*\alpha$ -lc but not conversely.

Proof: Proof follows from Definition 2.3(6)

Example 3.15: Let $X = \{1,2,3,4\}$ with the topology $\tau = \{\emptyset, X, \{1\}, \{1,2\}, \{1,3,4\}\}$. Then the set $\{1, 2, 4\}$ is $g^*\alpha$ -lc but not $g^\wedge \text{lc}$

Theorem 3.16: If (X, τ) is a ${}_T\alpha$ -space then $G^*\alpha \text{LC}(X) = \text{LC}(X) = \text{GLC}(X)$

Proof:

- (i) Since every $g^*\alpha$ -open set is open and $g^*\alpha$ -closed set is closed in (X, τ) , $G^*\alpha \text{LC}(X) \subseteq \text{LC}(X)$. Also for any topological space (X, τ) $\text{LC}(X) \subseteq G^*\alpha \text{LC}(X)$. Hence $G^*\alpha \text{LC}(X) = \text{LC}(X)$
- (ii) For any topological space (X, τ) $\text{LC}(X) \subseteq \text{GLC}(X)$. Also by Remark 2.6(7) and by hypothesis $\text{GLC}(X) \subseteq \text{LC}(X)$ and hence $\text{GLC}(X) = \text{LC}(X)$

From (i) and (ii) $G^*\alpha \text{LC}(X) = \text{LC}(X) = \text{GLC}(X)$

Definition 3.18: A subset A of (X, τ) is called

- (i) $g^*\alpha \text{lc}^*$ if $A = S \cap G$, where S is $g^*\alpha$ -open and G is closed in (X, τ) .
- (ii) $g^*\alpha \text{lc}^{**}$ if $A = S \cap G$, where S is open and G is $g^*\alpha$ -closed in (X, τ) .

The class of all $g^*\alpha$ -lc* (resp $g^*\alpha$ -lc**) sets in topological spaces (X, τ) is denoted by $G^*\alpha \text{LC}^*(X)$ (resp $G^*\alpha \text{LC}^{**}(X)$)

Prop 3.19: Every lc set is $g^*\alpha \text{lc}^*$ and $g^*\alpha \text{lc}^{**}$ but not conversely

Proof: It follows from Definition 2.3(1) and Definition 3.18

Example 3.20: Let $X = \{1,2,3,4\}$ with the topology $\tau = \{\emptyset, X, \{2\}, \{3,4\}, \{2,3,4\}\}$. Then the set $\{2, 3\}$ is $g^*\alpha \text{lc}^*$ and $g^*\alpha \text{lc}^{**}$ but not lc.

Prop 3.21: Every $g^*\alpha \text{lc}^*$ set is $g^*\alpha \text{lc}$ but not conversely.

Proof: It follows from Definition 3.18

Example 3.22: Let $X = \{1, 2, 3\}$ with the topology $\tau = \{\emptyset, X, \{1, 3\}\}$. Here the set $\{1, 2\}$ is $g^*\alpha \text{lc}$ but not $g^*\alpha \text{lc}^*$

Prop 3.23: Every lsc is $g^*\alpha \text{lc}^*$ and $g^*\alpha \text{lc}^{**}$ but not conversely

Proof: It follows from Definition 3.18 and Definition 2.3(4)

Example 3.24: Let $X = \{1,2,3\}$ with the topology $\tau = \{\emptyset, X, \{2, 3\}\}$. Here the set $\{2\}$ is $g^*\alpha \text{lc}^*$ and $g^*\alpha \text{lc}^{**}$ but not lsc

Prop 3.25: Every αlc is $g^*\alpha lc^*$ and $g^*\alpha lc^{**}$ but not conversely


Proof: It follows from Definition 3.18 and Definition 2.3(5)

Example 3.26: Let $X = \{1, 2, 3\}$ with the topology $\tau = \{\emptyset, X, \{1,2\}\}$. Here the set $\{1\}$ is $g^*\alpha lc^*$ and $g^*\alpha lc^{**}$ but not αlc

Prop 3.27: Every $glsc$ is $g^*\alpha lc^*$ and $g^*\alpha lc^{**}$ but not conversely

Proof: It follows from Definition 3.18 and Definition 2.3(3)

Example 3.28: Let $X = \{1, 2, 3, 4\}$ with the topology $\tau = \{\emptyset, X, \{1\}, \{1,4\}, \{1,2,4\}\}$. Here the set $\{1, 3\}$ is $g^*\alpha lc^*$ and $g^*\alpha lc^{**}$ but not $glsc$



Prop 3.29: If $G^*\alpha O(X) = \tau$ then $G^*\alpha LC(X) = G^*\alpha LC^*(X) = G^*\alpha LC^{**}(X)$

Proof: Since $G^*\alpha O(X) = \tau$, (X, τ) is a $T_{1/2}^{**}$ -space and hence $G^*\alpha LC(X) = G^*\alpha LC^*(X) = G^*\alpha LC^{**}(X)$

Remark 3.30: The converse of the above proposition need not be true

Let $X = \{1, 2, 3\}$ with the topology $\tau = \{\emptyset, X, \{1\}\}$. Here $G^*\alpha LC(X) = G^*\alpha LC^*(X) = G^*\alpha LC^{**}(X)$

However $G^*\alpha O(X) = \{\emptyset, X, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}\} \neq \tau$

Theorem 3.31: Assume that $G^*\alpha C(X)$ is closed under finite intersections. For a subset A of (X, τ) the following statements are equivalent.

1. $A \in G^*\alpha LC(X)$
2. $A = S \cap g^*\alpha cl(A)$ for some $g^*\alpha$ -open set S
3. $g^*\alpha-cl(A) - A$ is $g^*\alpha$ -closed
4. $A \cup (g^*\alpha-cl(A))^c$ is $g^*\alpha$ -open
5. $A \subseteq g^*\alpha-int(A \cup (g^*\alpha-cl(A))^c)$

Proof:

(1) \Rightarrow (2): Let $A \in G^*\alpha LC(X)$. Then $A = S \cap G$ where S is $g^*\alpha$ -open and G is $g^*\alpha$ -closed. Since $A \subseteq G$, $g^*\alpha-cl(A) \subseteq G$ so $S \cap g^*\alpha-cl(A) \subseteq A$. Also $A \subseteq S$ and $A \subseteq g^*\alpha cl(A)$ implies $A \subseteq S \cap g^*\alpha-cl(A)$ and therefore $A = S \cap g^*\alpha-cl(A)$

(2) \Rightarrow (3): $A = S \cap g^*\alpha cl(A)$ implies $g^*\alpha-cl(A) - A = g^*\alpha cl(A) \cap S^c$ which is $g^*\alpha$ -closed since S^c is $g^*\alpha$ -closed and $g^*\alpha-cl(A)$ is $g^*\alpha$ closed.

(3) \Rightarrow (4): $A \cup (g^*\alpha-cl(A))^c = (g^*\alpha-cl(A) - A)^c$ and by assumption, $(g^*\alpha-cl(A) - A)^c$ is $g^*\alpha$ -open and So is $A \cup (g^*\alpha-cl(A))^c$

(4) \Rightarrow (5): By assumption, $A \cup (g^*\alpha-cl(A))^c = g^*\alpha-int(A \cup (g^*\alpha-cl(A))^c)$ and hence $A \subseteq g^*\alpha-int(A \cup (g^*\alpha-cl(A))^c)$

(5) \Rightarrow (1): By assumption and since $A \subseteq g^*\alpha-cl(A)$, $A = g^*\alpha-int(A \cup (g^*\alpha-cl(A))^c) \cap g^*\alpha-cl(A)$. Therefore $A \in G^*\alpha LC(X)$.

Theorem 3.32: For a subset A of (X, τ) the following statements are equivalent

1. $A \in G^*\alpha LC^*(X)$
2. $A = S \cap cl(A)$ for some $g^*\alpha$ -open set S
3. $cl(A) - A$ is $g^*\alpha$ -closed
4. $A \cup (cl(A))^c$ is $g^*\alpha$ -open

Proof:

(1) \Rightarrow (2): Let $A \in G^*\alpha LC^*(X)$. There exists an $g^*\alpha$ -open set S and a closed set G such that $A = S \cap G$. Since $A \subseteq S$ and $A \subseteq cl(A)$, $A \subseteq S \cap cl(A)$. Also, since $cl(A) \subseteq G$, $S \cap cl(A) \subseteq S \cap G = A$. Therefore $A = S \cap cl(A)$

(2) \Rightarrow (1): Since S is $g^*\alpha$ -open and $cl(A)$ is a closed set $A = S \cap cl(A) \in G^*\alpha LC^*(X)$

(2) \Rightarrow (3): since $cl(A) - A = cl(A) \cap S^c$, $cl(A) - A$ is $g^*\alpha$ -closed by corollary 2.7

(3) \Rightarrow (2): Let $S = (cl(A) - A)^c$. Then S is $g^*\alpha$ -open in (X, τ) and $A = S \cap cl(A)$

(3) \Rightarrow (4): Let $G = cl(A) - A$. Then $G^c = AU (cl(A))^c$ and $AU (cl(A))^c$ is $g^*\alpha$ -open.

(4) \Rightarrow (3): Let $S = AU (cl(A))^c$. Then S^c is $g^*\alpha$ -closed and $S^c = cl(A) - A$ and so $cl(A) - A$ is $g^*\alpha$ -closed

Theorem 3.33: Let A be a subset of (X, τ) . Then $A \in G^*\alpha LC^{**}(X)$ iff $A = S \cap g^*\alpha-cl(A)$ for some open set S .

Proof: Let $A \in G^*\alpha LC^{**}(X)$. Then $A = S \cap G$ where S is open and G is $g^*\alpha$ -closed. Since $A \subseteq G$, $g^*\alpha-cl(A) \subseteq G$. We obtain $A = A \cap g^*\alpha-cl(A) = S \cap G \cap g^*\alpha-cl(A) = S \cap g^*\alpha-cl(A)$

Corollary 3.34: Let A be a subset of (X, τ) . If $A \in G^*\alpha LC^{**}(X)$ then $g^*\alpha-cl(A) - A$ is $g^*\alpha$ -closed and $AU(g^*\alpha-cl(A))^c$ is $g^*\alpha$ -open

Proof: Let $A \in G^*\alpha LC^{**}(X)$. Then by theorem 3.33, $A = S \cap g^*\alpha-cl(A)$ for some open set S and $g^*\alpha-cl(A) - A = g^*\alpha-cl(A) \cap S^c$ is $g^*\alpha$ -closed in (X, τ) . If $G = g^*\alpha-cl(A) - A$, then $G^c = AU(g^*\alpha-cl(A))^c$ and G^c is $g^*\alpha$ -open and so is $AU(g^*\alpha-cl(A))^c$

Prop 3.35:

- 1) If $S \in LC(X, \tau)$ then $S \in G^*\alpha LC(X, \tau), G^*\alpha LC^*(X, \tau), G^*\alpha LC^{**}(X, \tau)$
- 2) If $S \in GLC^*(X, \tau)$ then $S \in G^*\alpha LC(X, \tau), G^*\alpha LC^*(X, \tau), G^*\alpha LC^{**}(X, \tau)$
- 3) If $S \in G^\wedge LC(X, \tau)$ then $S \in G^*\alpha LC(X, \tau), G^*\alpha LC^{**}(X, \tau)$
- 4) If $S \in G^\# LC(X, \tau)$ then $S \in G^*\alpha LC(X, \tau), G^*\alpha LC^{**}(X, \tau)$
- 5) If $S \in G^*LC(X, \tau)$ then $S \in G^*\alpha LC(X, \tau), G^*\alpha LC^{**}(X, \tau)$

The proof is obvious from definitions 2.3, 3.1, 3.18

The converses of the prop 3.35 need not be true as seen from the following example

Example 3.36: Let $X = \{1, 2, 3\}$ with the topology $\tau = \{\emptyset, X, \{1\}, \{1, 3\}\}$. $G^*\alpha LC(X) = P(X)$
 $G^*\alpha LC^*(X) = P(X)$ $G^*\alpha LC^{**}(X) = P(X)$. $LC = \{\emptyset, X, \{1\}, \{2\}, \{3\}, \{1, 3\}, \{2, 3\}\}$. Here $A = \{1, 2\}$ and $\{1, 2\} \in G^*\alpha LC, G^*\alpha LC^*, G^*\alpha LC^{**}$, but $\{1, 2\} \notin LC$

Example 3.37: Let $X = \{1, 2, 3, 4\}$ with the topology $\tau = \{\emptyset, X, \{1\}, \{1, 4\}, \{1, 2, 4\}\}$. $G^*\alpha LC(X, \tau) = P(X)$
 $G^*\alpha LC^*(X) = P(X), G^*\alpha LC^{**}(X) = P(X)$. Let $A = \{1, 3\}$. Here $\{1, 3\} \in G^*\alpha LC, G^*\alpha LC^*, G^*\alpha LC^{**}$ but $\{1, 3\} \notin GLC^*$

Example 3.38: Let $X = \{1, 2, 3\}$ with the topology $\tau = \{\emptyset, X, \{2\}\}$. $G^*\alpha LC(X, \tau) = P(X)$
 $G^*\alpha LC^{**}(X) = P(X)$. Let $A = \{1, 2\}$. Here $\{1, 2\} \in G^*\alpha LC^*, G^*\alpha LC^{**}$ but $\{1, 2\} \notin G^\wedge LC, G^\# LC$

Example 3.39: Let $X = \{1, 2, 3\}$ with the topology $\tau = \{\emptyset, X, \{1\}\}$. $G^*\alpha LC(X, \tau) = P(X)$
 $G^*\alpha LC^{**}(X) = P(X)$. Let $A = \{2\}$. Here $\{1, 2\} \in G^*\alpha LC^*, G^*\alpha LC^{**}$ but $\{1, 2\} \notin G^*LC$

4. $g^*\alpha$ -dense sets and $g^*\alpha$ -submaximal spaces

We introduce the following definition:

Definition 4.1: A subset A of a space (X, τ) is called $g^*\alpha$ -dense if $g^*\alpha-cl(A) = X$

Example 4.2: Consider $X = \{1, 2, 3\}$ with $\tau = \{\emptyset, X, \{1\}, \{1, 3\}\}$. Then the set $\{1, 3\}$ is $g^*\alpha$ -dense in (X, τ)

Prop 4.3: Every $g^*\alpha$ -dense set is dense.

Proof: Let A be a $g^*\alpha$ -dense set in (X, τ) . Then $g^*\alpha-cl(A) = X$. since $g^*\alpha-cl(A) \subseteq cl(A)$. We have $cl(A) = X$ and so A is dense.

The converse of prop 4.3 need not be true as seen from the following example.

Example 4.4: Consider $X = \{1,2,3\}$ with $\tau = \{\emptyset, X, \{1\}, \{1,2\}\}$. Then the set $\{1,3\}$ is dense in (X, τ) but it is not $g^*\alpha$ -dense in (X, τ)

Definition 4.5: A topological space (X, τ) is called $g^*\alpha$ -submaximal if every dense subset in it is $g^*\alpha$ -open in (X, τ)

Prop 4.6: Every submaximal space is $g^*\alpha$ -submaximal

Proof: Let (X, τ) be a submaximal space and A be a dense subset of (X, τ) . Then A is open. But every open set is $g^*\alpha$ -open and so A is $g^*\alpha$ -open. Therefore (X, τ) is $g^*\alpha$ -submaximal.

The converse of prop 4.6 need not be true as seen from the following example.

Example 4.7: Consider $X = \{1, 2, 3\}$ with $\tau = \{\emptyset, X\}$. Then $G^*\alpha O(X) = P(X)$. We have every dense subset is $g^*\alpha$ open and hence (X, τ) is $g^*\alpha$ -submaximal. However the set $\{1,2\}$ is dense in (X, τ) but it is not open in (X, τ) . Therefore (X, τ) is not submaximal.

Prop 4.8: Every g -submaximal space is $g^*\alpha$ -submaximal

Proof: Let (X, τ) be a g -submaximal space and A be a dense subset of (X, τ) . Then A is g -open. But every g -open set is $g^*\alpha$ -open and so A is $g^*\alpha$ -open. Therefore (X, τ) is $g^*\alpha$ -submaximal.

The converse of prop 4.8 need not be true as seen from the following example.


Example 4.9: Consider $X = \{1,2,3,4\}$ with $\tau = \{\emptyset, X, \{1\}, \{2\}, \{1,2\}, \{1,2,3\}\}$. Then $G^*\alpha O(X) = \{\emptyset, X, \{1\}, \{2\}, \{3\}, \{1,2\}, \{1,3\}, \{2,3\}, \{1,2,3\}, \{1,2,4\}\}$. Every dense subset is $g^*\alpha$ -open and hence (X, τ) is $g^*\alpha$ -submaximal. However the set $\{1, 2, 4\}$ is dense in (X, τ) but it is not g -open in (X, τ) . Therefore (X, τ) is not g -submaximal.

Prop 4.10: Every $g^*\alpha$ -submaximal space is rg -submaximal

Proof: Let (X, τ) be a $g^*\alpha$ -submaximal space and A be a dense subset of (X, τ) . Then A is $g^*\alpha$ -open. But every $g^*\alpha$ -open set is rg -open and so A is rg -open. Therefore (X, τ) is rg -submaximal.

The converse of prop 4.10 need not be true as seen from the following example

Example 4.11: Consider $X = \{1,2,3\}$ with $\tau = \{\emptyset, X, \{2,3\}\}$. Then $G^*\alpha O(X) = P(X)$. we have every dense subset is rg open and hence (X, τ) is rg -submaximal. But the set $\{1,3\}$ is dense in (X, τ) but it is not $g^*\alpha$ -open in (X, τ) . Therefore (X, τ) is not $g^*\alpha$ -submaximal.



Theorem 4.12: A of a space (X, τ) is $g^*\alpha$ -submaximal iff $P(X) = G^*\alpha LC^*(X)$

Proof: Necessity: Let $A \in P(X)$ and Let $V = AU(\text{cl}(A))^c$. This implies that $\text{cl}(V) = \text{cl}(A) \cup (\text{cl}(A))^c = X$. Hence $\text{cl}(V) = X$. Therefore V is a dense subset of X . Since (X, τ) is $g^*\alpha$ -submaximal, V is $g^*\alpha$ -open.

Thus $AU(\text{cl}(A))^c$ is $g^*\alpha$ -open and by theorem 3.32, we have $A \in G^*\alpha LC^*(X)$

Sufficiency: Let A be a dense subset of (X, τ) . This implies $AU(\text{cl}(A))^c = AU X^c = AU \emptyset = A$. Now $A \in G^*\alpha LC^*(X)$ implies that $A = AU(\text{cl}(A))^c$ is $g^*\alpha$ -open by theorem 3.32. Hence (X, τ) is $g^*\alpha$ -submaximal.

Prop 4.13: Assume that $G^*O(X)$ forms a topology. For subsets A and B in (X, τ) , the following are true.

1. If $A, B \in G^*\alpha LC(X)$ then $A \cap B \in G^*\alpha LC(X)$
2. If $A, B \in G^*\alpha LC^*(X)$ then $A \cap B \in G^*\alpha LC^*(X)$
3. If $A, B \in G^*\alpha LC^{**}(X)$ then $A \cap B \in G^*\alpha LC^{**}(X)$
4. If $A \in G^*\alpha LC(X)$ and B is $g^*\alpha$ -open (resp $g^*\alpha$ -closed) then $A \cap B \in G^*\alpha LC(X)$
5. If $A \in G^*\alpha LC^*(X)$ and B is $g^*\alpha$ -open (resp closed) then $A \cap B \in G^*\alpha LC^*(X)$
6. If $A \in G^*\alpha LC^{**}(X)$ and B is $g^*\alpha$ -closed (resp open) then $A \cap B \in G^*\alpha LC^{**}(X)$
7. If $A \in G^*\alpha LC^*(X)$ and B is $g^*\alpha$ -closed then $A \cap B \in G^*\alpha LC(X)$
8. If $A \in G^*\alpha LC^{**}(X)$ and B is $g^*\alpha$ -open then $A \cap B \in G^*\alpha LC(X)$
9. If $A \in G^*\alpha LC^{**}(X)$ and B is $G^*\alpha LC^*(X)$ then $A \cap B \in G^*\alpha LC(X)$

Proof: By Remark 2.6 and corollary 2.7 (1) to (8) hold (9) Let $A = S \cap G$ where S is open and G is $g^*\alpha$ -closed and $B = P \cap Q$ where P is $g^*\alpha$ - open and Q is closed. Then $A \cap B = (S \cap P) \cap (G \cap Q)$ where $S \cap P$ is $g^*\alpha$ - open and $G \cap Q$ is $g^*\alpha$ -closed by corollary 2.7. Therefore $A \cap B \in G^*\alpha LC(X)$

Remark 4.14: Union of two $g^*\alpha$ -lc* sets need not be an $g^*\alpha$ -lc* set. This can be proved by the following example.

Example 4.15: Consider $X = \{1, 2, 3\}$ with $\tau = \{\emptyset, X, \{1, 2\}\}$. We have $G^*\alpha LC^*(X) = \{\emptyset, X, \{1\}, \{2\}, \{3\}, \{1, 2\}\}$. Then the sets $\{2\}$ and $\{3\}$ are $g^*\alpha$ -lc* sets but their union $\{2, 3\} \notin G^*\alpha LC^*(X)$

5. $G^*\alpha$ -Locally closed functions and some of their properties

In this section, the concept of $g^*\alpha$ -locally closed functions have been introduced and investigated the relation between $g^*\alpha$ -locally closed functions and some other locally closed functions.

Definition 5.1: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called $G^*\alpha LC$ -irresolute (resp $G^*\alpha LC^*$ -irresolute, $G^*\alpha LC^{**}$ -irresolute) if $f^{-1}(V) \in G^*\alpha LC(X)$ (resp $f^{-1}(V) \in G^*\alpha LC^*(X)$, $f^{-1}(V) \in G^*\alpha LC^{**}(X)$) for each $V \in G^*\alpha LC(Y)$ (resp $V \in G^*\alpha LC^*(Y, \sigma)$, $V \in G^*\alpha LC^{**}(Y, \sigma)$)

Definition 5.2: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called $G^*\alpha LC$ -continuous (resp $G^*\alpha LC^*$ -continuous, $G^*\alpha LC^{**}$ -continuous) if $f^{-1}(V) \in G^*\alpha LC(X, \tau)$ (resp $f^{-1}(V) \in G^*\alpha LC^*(X, \tau)$, $f^{-1}(V) \in G^*\alpha LC^{**}(X, \tau)$) for each open set V of (Y, σ)

Prop 5.3:

- (i) If f is LC -continuous then it is $G^*\alpha LC$, $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$ -continuous.
- (ii) If f is GLC^* -continuous then it is $G^*\alpha LC$, $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$ -continuous.
- (iii) If f is $G^\wedge LC$ -continuous then it is $G^*\alpha LC$ and $G^*\alpha LC^{**}$ -continuous.
- (iv) If f is $G^\# LC$ -continuous then it is $G^*\alpha LC$ and $G^*\alpha LC^{**}$ -continuous.
- (v) If f is $G^* LC$ -continuous then it is $G^*\alpha LC$ and $G^*\alpha LC^{**}$ -continuous.

Proof:

(i) Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be LC -continuous.

To prove $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)-continuous.

Let V be an open set of (Y, σ)

Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is LC -continuous, then $f^{-1}(V) \in LC(X, \tau)$ by prop 3.38(i) $f^{-1}(V) \in G^*\alpha LC(X, \tau)$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)

Therefore $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)-continuous.

(ii) Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be GLC^* -continuous

To prove $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)-continuous.

Let V be an open set of (Y, σ)

Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is GLC^* -continuous, then $f^{-1}(V) \in GLC^*(X, \tau)$ by prop 3.38(ii) $f^{-1}(V) \in G^*\alpha LC(X, \tau)$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)

Therefore $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)-continuous.

(iii) Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be $G^\wedge LC$ -continuous

To prove $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)-continuous.

Let V be an open set of (Y, σ)

Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^\wedge LC$ -continuous, then $f^{-1}(V) \in G^\wedge LC(X, \tau)$ by prop 3.38(iii) $f^{-1}(V) \in G^*\alpha LC(X, \tau)$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)

Therefore $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)-continuous.

(iv) Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be $G^\# LC$ -continuous

To prove $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)-continuous.

Let V be an open set of (Y, σ)

Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^\# LC$ -continuous, then $f^{-1}(V) \in G^\# LC(X, \tau)$ by prop 3.38(iv) $f^{-1}(V) \in G^*\alpha LC(X, \tau)$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)

Therefore $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)-continuous.

(v) Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be $G^* LC$ -continuous

To prove $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)-continuous.

Let V be an open set of (Y, σ)

Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^* LC$ -continuous, then $f^{-1}(V) \in G^* LC(X, \tau)$ by prop 3.38(v) $f^{-1}(V) \in G^*\alpha LC(X, \tau)$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)

Therefore $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ (resp $G^*\alpha LC^*$ and $G^*\alpha LC^{**}$)-continuous.

Example 5.4: Let $X = \{1,2,3\} = Y$ with $\tau = \{\emptyset, X, \{2,3\}\}$ and $\sigma = \{\emptyset, Y, \{2\}\}$. Define $f: (X, \tau) \rightarrow (Y, \sigma)$ by $f(1) = 1, f(2) = 2$ and $f(3) = 3$. LC sets of (X, τ) are $\{\{1\}, \{2,3\}, \emptyset, X\}$. $G^*\alpha$ LC sets of $(X, \tau) = P(X)$. $G^*\alpha$ LC* sets of $(X, \tau) = \{\emptyset, X, \{1\}, \{2\}, \{3\}, \{2,3\}\}$. $G^*\alpha$ LC** sets of $(X, \tau) = P(X)$. Here $f^{-1}(\emptyset) = \emptyset \in G^*\alpha$ LC (X, τ) (resp $G^*\alpha$ LC*(X, τ), $G^*\alpha$ LC**(X, τ)), $f^{-1}(Y) = Y \in G^*\alpha$ LC (X, τ) (resp $G^*\alpha$ LC*(X, τ), $G^*\alpha$ LC**(X, τ)) for every open set V of (Y, σ) . Hence f is $G^*\alpha$ LC, $G^*\alpha$ LC* and $G^*\alpha$ LC**-continuous, Since $\{2\}$ is an open set of (Y, σ) . But $f^{-1}(\{2\}) = \{2\} \notin LC(X, \tau)$

Example 5.5: Let $X = \{1,2,3\} = Y$ with $\tau = \{\emptyset, X, \{1\}, \{1,3\}\}$ and $\sigma = \{\emptyset, Y, \{1,2\}\}$. Define $f: (X, \tau) \rightarrow (Y, \sigma)$ by $f(1) = 1, f(2) = 2$ and $f(3) = 3$. GLC* sets of (X, τ) are $\{\{1\}, \{2\}, \{3\}, \{1,3\}, \{2,3\}, \emptyset, X\}$. $G^*\alpha$ LC sets of $(X, \tau) = P(X)$. $G^*\alpha$ LC* sets of $(X, \tau) = P(X)$. $G^*\alpha$ LC** sets of $(X, \tau) = P(X)$. Here $f^{-1}(\emptyset) = \emptyset \in G^*\alpha$ LC (X, τ) (resp $G^*\alpha$ LC*(X, τ), $G^*\alpha$ LC**(X, τ)), $f^{-1}(Y) = Y \in G^*\alpha$ LC (X, τ) (resp $G^*\alpha$ LC*(X, τ), $G^*\alpha$ LC**(X, τ)) for every open set V of (Y, σ) . Hence f is $G^*\alpha$ LC, $G^*\alpha$ LC* and $G^*\alpha$ LC**-continuous, Since $\{1,2\}$ is an open set of (Y, σ) . But $f^{-1}(\{1,2\}) = \{1,2\} \notin GLC^*(X, \tau)$

Example 5.6: Let $X = \{1,2,3,4\} = Y$ with $\tau = \{\emptyset, X, \{1\}, \{1,2\}, \{1,3,4\}\}$ and $\sigma = \{\emptyset, Y, \{1\}, \{1,4\}, \{1,2,4\}\}$. Define $f: (X, \tau) \rightarrow (Y, \sigma)$ by $f(1) = 1, f(2) = 2, f(3) = 3$ and $f(4) = 4$. $G^*\alpha$ LC sets of (X, τ) are $\{\{1\}, \{2\}, \{1,2\}, \{3,4\}, \{1,3,4\}, \{2,3,4\}, \emptyset, X\}$. $G^*\alpha$ LC sets of $(X, \tau) = P(X)$. $G^*\alpha$ LC** sets of $(X, \tau) = P(X)$. $G^*\alpha$ LC** sets of $(X, \tau) = P(X)$. Here $f^{-1}(\emptyset) = \emptyset \in G^*\alpha$ LC (X, τ) (resp $G^*\alpha$ LC**(X, τ)), $f^{-1}(Y) = Y \in G^*\alpha$ LC (X, τ) (resp $G^*\alpha$ LC**(X, τ)) for every open set V of (Y, σ) . Hence f is $G^*\alpha$ LC and $G^*\alpha$ LC**-continuous, Since $\{1,2,4\}$ is an open set of (Y, σ) . But $f^{-1}(\{1,2,4\}) = \{1,2,4\} \notin G^*\alpha$ LC (X, τ)

Example 5.7: Let $X = \{1, 2, 3\} = Y$ with $\tau = \{\emptyset, X, \{1\}\}$ and $\sigma = \{\emptyset, Y, \{1\}, \{1,3\}\}$. Define $f: (X, \tau) \rightarrow (Y, \sigma)$ by $f(1) = 1, f(2) = 2$ and $f(3) = 3$. $G^{\#}$ LC sets of (X, τ) are $\{\{1\}, \{2,3\}, \emptyset, X\}$. $G^*\alpha$ LC sets of $(X, \tau) = P(X)$. $G^*\alpha$ LC** sets of $(X, \tau) = P(X)$. Here $f^{-1}(\emptyset) = \emptyset \in G^*\alpha$ LC (X, τ) (resp $G^*\alpha$ LC**(X, τ)), $f^{-1}(Y) = Y \in G^*\alpha$ LC (X, τ) (resp $G^*\alpha$ LC**(X, τ)) for every open set V of (Y, σ) . Hence f is $G^*\alpha$ LC and $G^*\alpha$ LC**-continuous, Since $\{1,3\}$ is an open set of (Y, σ) . But $f^{-1}(\{1,3\}) = \{1,3\} \notin G^{\#}$ LC (X, τ)

Example 5.8: Let $X = \{1,2,3\} = Y$ with $\tau = \{\emptyset, X, \{1\}\}$ and $\sigma = \{\emptyset, Y, \{2\}, \{2,3\}\}$. Define $f: (X, \tau) \rightarrow (Y, \sigma)$ by $f(1) = 1, f(2) = 2$ and $f(3) = 3$. $G^{\#}$ LC sets of (X, τ) are $\{\{1\}, \{2,3\}, \emptyset, X\}$. $G^*\alpha$ LC sets of $(X, \tau) = P(X)$. $G^*\alpha$ LC** sets of $(X, \tau) = P(X)$. Here $f^{-1}(\emptyset) = \emptyset \in G^*\alpha$ LC (X, τ) (resp $G^*\alpha$ LC**(X, τ)), $f^{-1}(Y) = Y \in G^*\alpha$ LC (X, τ) (resp $G^*\alpha$ LC**(X, τ)) for every open set V of (Y, σ) . Hence f is $G^*\alpha$ LC and $G^*\alpha$ LC**-continuous, Since $\{2\}$ is an open set of (Y, σ) . But $f^{-1}(\{2\}) = \{2\} \notin G^{\#}$ LC (X, τ)

Theorem 4.9: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ and $g: (Y, \sigma) \rightarrow (Z, \eta)$ be any two functions. Then

- (i) $g \circ f$ is $G^*\alpha$ LC-irresolute if f and g are $G^*\alpha$ LC-irresolute
- (ii) $g \circ f$ is $G^*\alpha$ LC*-irresolute if f and g are $G^*\alpha$ LC*-irresolute
- (iii) $g \circ f$ is $G^*\alpha$ LC**-irresolute if f and g are $G^*\alpha$ LC**-irresolute
- (iv) $g \circ f$ is $G^*\alpha$ LC-continuous if f is $G^*\alpha$ LC-continuous and g is continuous.
- (v) $g \circ f$ is $G^*\alpha$ LC*-continuous if f is $G^*\alpha$ LC*-continuous and g is continuous.
- (vi) $g \circ f$ is $G^*\alpha$ LC**-continuous if f is $G^*\alpha$ LC**-continuous and g is continuous.
- (vii) $g \circ f$ is $G^*\alpha$ LC-continuous if f is $G^*\alpha$ LC-irresolute and g is $G^*\alpha$ LC-continuous
- (viii) $g \circ f$ is $G^*\alpha$ LC*-continuous if f is $G^*\alpha$ LC*-irresolute and g is $G^*\alpha$ LC*-continuous
- (ix) $g \circ f$ is $G^*\alpha$ LC**-continuous if f is $G^*\alpha$ LC**-irresolute and g is $G^*\alpha$ LC**-continuous

Proof:

(i) Given $f: (X, \tau) \rightarrow (Y, \sigma)$ and $g: (Y, \sigma) \rightarrow (Z, \eta)$ are $G^*\alpha$ LC-irresolute.

To prove $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha$ LC-irresolute.

Let $V \in G^*\alpha$ LC (z, η)

Since $g: (Y, \sigma) \rightarrow (Z, \eta)$ is $G^*\alpha$ LC-irresolute, then $g^{-1}(V) \in G^*\alpha$ LC (Y, σ)

Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha$ LC-irresolute, then $f^{-1}(g^{-1}(V)) \in G^*\alpha$ LC (X, τ)

(ie) $(g \circ f)^{-1}(V) \in G^*\alpha$ LC (X, τ)

Thus we get $(g \circ f)^{-1}(V) \in G^*\alpha$ LC (X, τ) for every $v \in G^*\alpha$ LC $\in (Z, \eta)$

Hence $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha$ LC-irresolute.

(ii) Given $f: (X, \tau) \rightarrow (Y, \sigma)$ and $g: (Y, \sigma) \rightarrow (Z, \eta)$ are $G^*\alpha$ LC*-irresolute.

To prove $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha$ LC*-irresolute.

Let $V \in G^*\alpha$ LC*(z, η)

Since $g: (Y, \sigma) \rightarrow (Z, \eta)$ is $G^*\alpha$ LC*-irresolute, then $g^{-1}(V) \in G^*\alpha$ LC*(Y, σ)

Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha$ LC*-irresolute, then $f^{-1}(g^{-1}(V)) \in G^*\alpha$ LC*(X, τ)

(ie) $(g \circ f)^{-1}(V) \in G^*\alpha$ LC*(X, τ)

Thus we get $(g \circ f)^{-1}(V) \in G^*\alpha$ LC*(X, τ) for every $v \in G^*\alpha$ LC* $\in (Z, \eta)$

Hence $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha$ LC*-irresolute.

- (iii)** Given $f: (X, \tau) \rightarrow (Y, \sigma)$ and $g: (Y, \sigma) \rightarrow (Z, \eta)$ are $G^*\alpha LC^{**}$ -irresolute.
 To prove $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC^{**}$ -irresolute.
 Let $V \in G^*\alpha LC^{**}(Z, \eta)$
 Since $g: (Y, \sigma) \rightarrow (Z, \eta)$ is $G^*\alpha LC^{**}$ -irresolute, then $g^{-1}(V) \in G^*\alpha LC^{**}(Y, \sigma)$
 Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC^{**}$ -irresolute, then $f^{-1}(g^{-1}(V)) \in G^*\alpha LC^{**}(X, \tau)$
 (ie) $(g \circ f)^{-1}(V) \in G^*\alpha LC^{**}(X, \tau)$
 Thus we get $(g \circ f)^{-1}(V) \in G^*\alpha LC^{**}(X, \tau)$ for every $v \in G^*\alpha LC^{**} \in (Z, \eta)$
 Hence $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC^{**}$ -irresolute
- (iv)** Given $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ -continuous and $g: (Y, \sigma) \rightarrow (Z, \eta)$ is continuous
 To prove $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC$ -continuous
 Let V be an open set of (Z, η)
 Since $g: (Y, \sigma) \rightarrow (Z, \eta)$ is continuous, then $g^{-1}(V)$ is an open set of (Y, σ)
 Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ -continuous, then $f^{-1}(g^{-1}(V)) \in G^*\alpha LC(X, \tau)$
 (ie) $(g \circ f)^{-1}(V) \in G^*\alpha LC(X, \tau)$
 Thus we get $(g \circ f)^{-1}(V) \in G^*\alpha LC(X, \tau)$ for every open set V of (Z, η)
 Hence $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC$ -continuous.
- (v)** Given $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC^*$ -continuous and $g: (Y, \sigma) \rightarrow (Z, \eta)$ is continuous
 To prove $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC^*$ -continuous
 Let V be an open set of (Z, η)
 Since $g: (Y, \sigma) \rightarrow (Z, \eta)$ is continuous, then $g^{-1}(V)$ is an open set of (Y, σ)
 Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC^*$ -continuous, then $f^{-1}(g^{-1}(V)) \in G^*\alpha LC^*(X, \tau)$
 (ie) $(g \circ f)^{-1}(V) \in G^*\alpha LC^*(X, \tau)$
 Thus we get $(g \circ f)^{-1}(V) \in G^*\alpha LC^*(X, \tau)$ for every open set V of (Z, η)
 Hence $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC^*$ -continuous.
- (vi)** Given $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC^{**}$ -continuous and $g: (Y, \sigma) \rightarrow (Z, \eta)$ is continuous
 To prove $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC^{**}$ -continuous
 Let V be an open set of (Z, η)
 Since $g: (Y, \sigma) \rightarrow (Z, \eta)$ is continuous, then $g^{-1}(V)$ is an open set of (Y, σ)
 Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC^{**}$ -continuous, then $f^{-1}(g^{-1}(V)) \in G^*\alpha LC^{**}(X, \tau)$
 (ie) $(g \circ f)^{-1}(V) \in G^*\alpha LC^{**}(X, \tau)$
 Thus we get $(g \circ f)^{-1}(V) \in G^*\alpha LC^{**}(X, \tau)$ for every open set V of (Z, η)
 Hence $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC^{**}$ -continuous
- vii)** Given $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ -irresolute and $g: (Y, \sigma) \rightarrow (Z, \eta)$ is $G^*\alpha LC$ -continuous
 To prove $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC$ -continuous
 Let V be an open set of (Z, η)
 Since $g: (Y, \sigma) \rightarrow (Z, \eta)$ is $G^*\alpha LC$ -continuous, then $g^{-1}(V) \in G^*\alpha LC(Y, \sigma)$
 Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC$ -irresolute, then $f^{-1}(g^{-1}(V)) \in G^*\alpha LC(X, \tau)$
 (ie) $(g \circ f)^{-1}(V) \in G^*\alpha LC(X, \tau)$
 Thus we get $(g \circ f)^{-1}(V) \in G^*\alpha LC(X, \tau)$ for every open set V of (Z, η)
 Hence $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC$ -continuous.
- viii)** Given $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC^*$ -irresolute and $g: (Y, \sigma) \rightarrow (Z, \eta)$ is $G^*\alpha LC^*$ -continuous
 To prove $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC^*$ -continuous
 Let V be an open set of (Z, η)
 Since $g: (Y, \sigma) \rightarrow (Z, \eta)$ is $G^*\alpha LC^*$ -continuous, then $g^{-1}(V) \in G^*\alpha LC^*(Y, \sigma)$
 Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC^*$ -irresolute, then $f^{-1}(g^{-1}(V)) \in G^*\alpha LC^*(X, \tau)$
 (ie) $(g \circ f)^{-1}(V) \in G^*\alpha LC^*(X, \tau)$
 Thus we get $(g \circ f)^{-1}(V) \in G^*\alpha LC^*(X, \tau)$ for every open set V of (Z, η)
 Hence $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC^*$ -continuous.
- ix)** Given $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC^{**}$ -irresolute and $g: (Y, \sigma) \rightarrow (Z, \eta)$ is $G^*\alpha LC^{**}$ -continuous
 To prove $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC^{**}$ -continuous
 Let V be an open set of (Z, η)
 Since $g: (Y, \sigma) \rightarrow (Z, \eta)$ is $G^*\alpha LC^{**}$ -continuous, then $g^{-1}(V) \in G^*\alpha LC^{**}(Y, \sigma)$
 Since $f: (X, \tau) \rightarrow (Y, \sigma)$ is $G^*\alpha LC^{**}$ -irresolute, then $f^{-1}(g^{-1}(V)) \in G^*\alpha LC^{**}(X, \tau)$
 (ie) $(g \circ f)^{-1}(V) \in G^*\alpha LC^{**}(X, \tau)$
 Thus we get $(g \circ f)^{-1}(V) \in G^*\alpha LC^{**}(X, \tau)$ for every open set V of (Z, η)
 Hence $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ is $G^*\alpha LC^{**}$ -continuous.

REFERENCES

1. k.Balachandran, P.Sundaram and H.Maki, Generalized locally closed sets and GLC –continuous functions, Indian J.Pure Appl Math., 27(3)(1996), 235 -244.
2. P.Bhattacharyya and B.K.Lahiri, Semi-generalized closed sets in topology, Indian J.Math., 29(3) (1987), 375 -382.
3. N.Bourbaki, General topology, Part I, Addison – Wesley, Reading, Mass., (1966)
4. S.G.Crossley and S.K.Hildebrand, Semi-closure, Texas J. Sci., 22(1971), 99-112.
5. J.Dontchev, On submaximal spaces, Tamkang J.Math., 26(1995), 253-260.
6. M.Ganster and I.L.Reilly, Locally closed sets and LC-continuous functions, Internat J.Math. Math. Sci., 12(3) (1989), 417-424.
7. Y.Gnanambal, Studies on generalized pre-regular closed sets and generalization of locally closed sets, Ph.D Thesis, Bharathiar University, Coimbatore (1998).
8. N.Levine, Generalized closed sets in Topology, Rend. Circ. Math. Palermo, 19(2) (1970), 89-96.
9. N.Levine, Semi-open sets and semi-continuity in topological spaces, Amer. Math. Monthly. 70(1963), 36-41.
10. H.Maki, R.Devi and K.Balachandran, Associated topologies of generalized α -closed sets and α -generalized closed sets Mem. Fac. Sci. Kochi. Univ. Ser. A. Math., 15(1994), 51-63.
11. A.S.Mashhour, I.A.Hasanein and S.N.El-Deeb, α -continuous and α -open mappings, Acta Math. Hungar., 41 (1983), 213-218.
12. O.Njastad, On some classes of nearly open sets, Pacific J. Math., 15(1965), 961-970.
13. N.Palaniappan and K.C.Rao, R Regular generalized closed sets, Kyungpook Math. J., 33(1993), 211-219.
14. A.Punitha Tharani and T.Delcia, $g^*\alpha$ -closed sets in Topological spaces, IJMA, 8(10), 2017, 71-80.
15. A.H.Stone, Absolutely FG spaces, Proc. Amer. Math. Soc., 80(1980), 515-520.
16. M.H.Stone, Applications of the theory of Boolean rings to general topology, Trans. Amer. Math. Soc., 41 (1937), 374-481.
17. M. K. R. S. Veera Kumar, Between Closed Sets and g -closed sets *Mem. Fac. Sci. Kochi. Univ. Ser. A. Math.* 21 (2000) 1-19.
18. M. K. R. S. Veera Kumar, $g^\#$ -closed sets in topological spaces, *Mem. Fac. Sci. Kochi Univ. Ser. A. Math.* 24 (2003) 1-13.
19. M. K. R. S. Veera Kumar, $g^\#$ -locally closed sets and $G^\#LC$ -functions. *Antarctica J. Math.* 1 (1) (2004) 35-46.
20. M. K. R. S. Veera Kumar, On g^\wedge -closed sets in topological spaces. *Bulletion Allahabad Math. Soc.* 18 (2003) 99-112.
21. M. K. R. S. Veera Kumar, g^\wedge -locally closed sets and $G^\wedge LC$ –functions, *Indian J. Math.* 43(2) (2001) 231-247.
22. M. K. R. S. Veera Kumar, g^* -locally closed sets and G^*LC -functions. *Univ. Beacu. Stud. St. Ser. Mat* 13 (2003) 49-58.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]