

MULTIPLICATIVE CONNECTIVITY
REVAN INDICES OF CERTAIN FAMILIES OF BENZENOID SYSTEMS

V. R. KULLI*

Department of Mathematics, Gulbarga University, Gulbarga - 585106, India.

(Received On: 06-02-18; Revised & Accepted On: 09-03-18)

ABSTRACT

In Chemical Science, the multiplicative connectivity indices are applied to measure the chemical and biological characteristics of chemical compounds. In this paper, we introduce the multiplicative product connectivity Revan index, multiplicative sum connectivity Revan index, first multiplicative atom bond connectivity Revan index and multiplicative geometric-arithmetic Revan index of a molecular graph and compute these multiplicative connectivity Revan indices of some important chemical structures like triangular benzenoid, benzenoid rhombus and benzenoid hourglass.

Mathematics Subject Classification: 05C12, 05C07, 05C35.

Keywords: molecular graph, multiplicative connectivity Revan indices, triangular benzenoid, benzenoid rhombus.

1. INTRODUCTION

In this paper, we consider only a finite, simple connected graph. Let G be a graph with a vertex set $V(G)$ and an edge set $E(G)$. The degree $d_G(v)$ of a vertex v is the number of vertices adjacent to v . Let $\Delta(G)$ ($\delta(G)$) denote the maximum (minimum) degree among the vertices of G . The Revan vertex degree of a vertex v in G is defined as $r_G(v) = \Delta(G) + \delta(G) - d_G(v)$. The Revan edge connecting the Revan vertices u and v will be denoted by uv . For additional definitions and notations, the reader may refer to [1].

A molecular graph is a graph such that its vertices correspond to the atoms and the edges to the bonds. Chemical graph theory is a branch of Mathematical chemistry which has an important effect on the development of the Chemical Sciences. A topological index is a numerical parameter mathematically derived from the graph structure. In organic chemistry, topological indices have been found to be useful in chemical documentation, isomer discrimination, structure property relationships, structure activity relationships and pharmaceutical drug design. There has been considerable interest in the general problem of determining topological indices.

Best known and used topological indices are the multiplicative connectivity indices, introduced by Kulli in [2]. Motivated by the definitions of the multiplicative connectivity indices and their wide applications, we introduce the multiplicative product connectivity Revan index, multiplicative sum connectivity Revan index, multiplicative atom bond connectivity Revan index and multiplicative geometric-arithmetic Revan index of a molecular graph as follows:

The multiplicative product connectivity Revan index of a graph G is defined as

$$PRII(G) = \prod_{uv \in E(G)} \frac{1}{\sqrt{r_G(u)r_G(v)}}$$

The multiplicative sum connectivity Revan index of a graph G is defined as

$$SRII(G) = \prod_{uv \in E(G)} \frac{1}{\sqrt{r_G(u) + r_G(v)}}$$

Corresponding Author: V. R. Kulli*

Department of Mathematics, Gulbarga University, Gulbarga 585106, India.

The first multiplicative atom bond connectivity Revan index of a graph G is defined as

$$ABC_1RII(G) = \prod_{uv \in E(G)} \sqrt{\frac{r_G(u) + r_G(v) - 2}{r_G(u)r_G(v)}}$$

The multiplicative geometric-arithmetic Revan index of a graph G is defined as

$$GARII(G) = \prod_{uv \in E(G)} \frac{2\sqrt{r_G(u)r_G(v)}}{r_G(u) + r_G(v)}$$

Recently many multiplicative topological indices were studied, for example, in [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 14, 15, 16]. Also some connectivity indices were studied, for example, in [17, 18, 19, 20, 21, 22]. In this paper, we compute multiplicative connectivity Revan indices of triangular benzenoids, benzenoid rhombus and benzenoid hourglass. For more information about these benzenoids see [23, 24].

2. RESULTS FOR TRIANGULAR BENZENOID T_p

In this section, we consider the graph of triangular benzenoid T_p which p is the number of hexagons in the base graph. Clearly T_p has $\frac{1}{2}p(p+1)$ hexagons. The graph of triangular benzenoid T_4 is presented in Figure 1.




Figure-1: The graph of triangular benzenoid T_4 .

Let G be the graph of a triangular benzenoid T_p . By algebraic method, we obtain $|V(T_p)| = p^2 + 4p + 1$ and $|E(T_p)| = \frac{3}{2}p(p+3)$. Also by algebraic method, we obtain that the edge set $E(G)$ can be divided into three partitions:

$$\begin{aligned} E_{22} &= \{uv \in E(G) \mid d_G(u) = d_G(v) = 2\} & |E_{22}| &= 6. \\ E_{23} &= \{uv \in E(G) \mid d_G(u) = 2, d_G(v) = 3\} & |E_{23}| &= 6p - 6. \\ E_{33} &= \{uv \in E(G) \mid d_G(u) = d_G(v) = 3\} & |E_{33}| &= \frac{3}{2}p(p-1). \end{aligned}$$

Clearly $\Delta(G) = 3$ and $\delta(G) = 2$. Therefore $r_G(u) = 5 - d_G(u)$. Thus we ensure that there are three types of Revan edges based on the degree of end Revan vertices of each Revan edge as follows:

$$\begin{aligned} RE_{33} &= \{uv \in E(G) \mid r_G(u) = r_G(v) = 3\}, & |RE_{33}| &= 6. \\ RE_{32} &= \{uv \in E(G) \mid r_G(u) = 3, d_G(v) = 2\}, & |RE_{32}| &= 6(p-1). \\ RE_{22} &= \{uv \in E(G) \mid r_G(u) = r_G(v) = 2\} & |RE_{22}| &= \frac{3}{2}p(p-1). \end{aligned}$$

In the following theorem, we compute the multiplicative product connectivity Revan index of T_p .

Theorem 1: The multiplicative product connectivity Revan index of a triangular benzenoid T_p is given by

$$PRII(T_p) = \left(\frac{1}{3}\right)^6 \times \left(\frac{1}{6}\right)^{3(p-1)} \times \left(\frac{1}{2}\right)^{\frac{3}{2}p(p-1)}.$$

Proof: By definition, we have $PRII(G) = \prod_{uv \in E(G)} \frac{1}{\sqrt{r_G(u)r_G(v)}}$.

$$\begin{aligned} \text{Thus } PRII(T_p) &= \left(\frac{1}{\sqrt{3 \times 3}}\right)^6 \times \left(\frac{1}{\sqrt{3 \times 2}}\right)^{6(p-1)} \times \left(\frac{1}{\sqrt{2 \times 2}}\right)^{\frac{3}{2}p(p-1)} \\ &= \left(\frac{1}{3}\right)^6 \times \left(\frac{1}{6}\right)^{3(p-1)} \times \left(\frac{1}{2}\right)^{\frac{3}{2}p(p-1)}. \end{aligned}$$

In the following theorem, we compute the multiplicative sum connectivity Revan index of T_p .

Theorem 2: The multiplicative sum connectivity Revan index of a triangular benzenoid T_p is given by

$$SRII(T_p) = \left(\frac{1}{6}\right)^3 \times \left(\frac{1}{5}\right)^{3(p-1)} \times \left(\frac{1}{2}\right)^{\frac{3}{2}p(p-1)}.$$

Proof: By definition, we have $SRII(G) = \prod_{uv \in E(G)} \frac{1}{\sqrt{r_G(u) + r_G(v)}}$

$$\begin{aligned} \text{Thus } SRII(T_p) &= \left(\frac{1}{\sqrt{3+3}}\right)^6 \times \left(\frac{1}{\sqrt{3+2}}\right)^{6(p-1)} \times \left(\frac{1}{\sqrt{2+2}}\right)^{\frac{3}{2}p(p-1)}. \\ &= \left(\frac{1}{6}\right)^3 \times \left(\frac{1}{5}\right)^{3(p-1)} \times \left(\frac{1}{2}\right)^{\frac{3}{2}p(p-1)}. \end{aligned}$$

In the following theorem, we compute the multiplicative atom bond connectivity Revan index of T_p .

Theorem 3: The first multiplicative atom bond connectivity Revan index of a triangular benzenoid T_p is given by

$$ABC_1II(T_p) = \left(\frac{2}{3}\right)^6 \times \left(\frac{1}{2}\right)^{3(p-1)} \times \left(\frac{1}{\sqrt{2}}\right)^{\frac{3}{2}p(p-1)}.$$

Proof: By definition, we have $ABCRII(T_p) = \prod_{uv \in E(G)} \sqrt{\frac{r_G(u) + r_G(v) - 2}{r_G(u)r_G(v)}}$.

$$\begin{aligned} \text{Thus } ABC_1RII(T_p) &= \left(\sqrt{\frac{3+3-2}{3 \times 3}}\right)^6 \times \left(\sqrt{\frac{3+2-2}{3 \times 2}}\right)^{6(p-1)} \times \left(\sqrt{\frac{2+2-2}{2 \times 2}}\right)^{\frac{3}{2}p(p-1)}. \\ &= \left(\frac{2}{3}\right)^6 \times \left(\frac{1}{2}\right)^{3(p-1)} \times \left(\frac{1}{\sqrt{2}}\right)^{\frac{3}{2}p(p-1)}. \end{aligned}$$

In the following theorem, we compute the multiplicative geometric-arithmetic Revan index of T_p .

Theorem 4: The multiplicative geometric-arithmetic Revan index of a triangular benzenoid T_p is given by

$$GARII(T_p) = \left(\frac{2\sqrt{6}}{5}\right)^{6(p-1)}.$$

Proof: By definition, we have $GARII(T_p) = \prod_{uv \in E(G)} \frac{2\sqrt{r_G(u)r_G(v)}}{r_G(u) + r_G(v)}$.

$$\begin{aligned} \text{Thus } GARII(T_p) &= \left(\frac{2\sqrt{3 \times 3}}{3+3}\right)^6 \times \left(\frac{2\sqrt{3 \times 2}}{3+2}\right)^{6(p-1)} \times \left(\frac{2\sqrt{2 \times 2}}{2+2}\right)^{\frac{3}{2}p(p-1)}. \\ &= \left(\frac{2\sqrt{6}}{5}\right)^{6(p-1)}. \end{aligned}$$

3. RESULTS FOR BENZENOID RHOMBUS R_p .

In this section, we consider the graph of benzenoid rhombus R_p which is obtained from two copies of a triangular benzenoid T_p by identifying hexagons in one of their base rows. The graph of benzenoid rhombus R_4 is presented in Figure 2.




Figure-2: The graph of benzenoid rhombus R_4 .

Let G be the graph of a benzenoid rhombus R_p . By algebraic method, we obtain $|V(T_p)| = 2p^2 + 4p$ and $|E(R_p)| = 3p^2 + 4p - 1$. It is easy to see that the vertices of benzenoid rhombus R_p are either of degree 2 or 3, see Figure 2. Therefore $\Delta(G)=3$ and $\delta(G)=2$. Thus $r_G(u) = 5 - d_G(u)$. By calculation, we obtain that the edge set $E(R_p)$ can be divided into three partitions:

$$\begin{aligned} E_{22} &= \{uv \in E(G) \mid d_G(u) = d_G(v) = 2\} & |E_{22}| &= 6. \\ E_{23} &= \{uv \in E(G) \mid d_G(u) = 2, d_G(v) = 3\} & |E_{23}| &= 8(p - 1). \\ E_{33} &= \{uv \in E(G) \mid d_G(u) = d_G(v) = 3\} & |E_{33}| &= 3p^2 - 4p + 1. \end{aligned}$$

Thus there are three types of Revan edges as follows:

$$\begin{aligned} RE_{33} &= \{uv \in E(G) \mid r_G(u) = r_G(v) = 3\} & |RE_{33}| &= 6. \\ RE_{32} &= \{uv \in E(G) \mid r_G(u) = 3, r_G(v) = 2\} & |RE_{32}| &= 8(p - 1). \\ RE_{22} &= \{uv \in E(G) \mid r_G(u) = r_G(v) = 2\} & |RE_{22}| &= 3p^2 - 4p + 1. \end{aligned}$$

Theorem 5: The multiplicative product connectivity Revan index of a benzenoid rhombus R_p is given by

$$PRII(R_p) = \left(\frac{1}{3}\right)^6 \times \left(\frac{1}{6}\right)^{4(p-1)} \times \left(\frac{1}{2}\right)^{3p^2-4p+1}.$$

Proof: To compute $PRII(R_p)$, we see that

$$\begin{aligned} PRII(R_p) &= \prod_{uv \in E(G)} \frac{1}{\sqrt{r_G(u)r_G(v)}} = \left(\frac{1}{\sqrt{3 \times 3}}\right)^6 \times \left(\frac{1}{\sqrt{3 \times 2}}\right)^{8(p-1)} \times \left(\frac{1}{\sqrt{2 \times 2}}\right)^{3p^2-4p+1} \\ &= \left(\frac{1}{3}\right)^6 \times \left(\frac{1}{6}\right)^{4(p-1)} \times \left(\frac{1}{2}\right)^{3p^2-4p+1}. \end{aligned}$$

Theorem 6: The multiplicative sum connectivity Revan index of a benzenoid rhombus R_p is given by

$$SRII(R_p) = \left(\frac{1}{6}\right)^3 \times \left(\frac{1}{5}\right)^{4(p-1)} \times \left(\frac{1}{2}\right)^{3p^2-4p+1}.$$

Proof: To compute $SRII(R_p)$, we see that

$$\begin{aligned} SRII(R_p) &= \prod_{uv \in E(G)} \frac{1}{\sqrt{r_G(u) + r_G(v)}} = \left(\frac{1}{\sqrt{3+3}}\right)^6 \times \left(\frac{1}{\sqrt{3+2}}\right)^{8(p-1)} \times \left(\frac{1}{\sqrt{2+2}}\right)^{3p^2-4p+1} \\ &= \left(\frac{1}{6}\right)^3 \times \left(\frac{1}{5}\right)^{4(p-1)} \times \left(\frac{1}{2}\right)^{3p^2-4p+1}. \end{aligned}$$

Theorem 7: The first multiplicative atom bond connectivity Revan index of a benzenoid rhombus R_p is given by

$$ABC_1RII(R_p) = \left(\frac{2}{3}\right)^6 \times \left(\frac{1}{2}\right)^{4(p-1)} \times \left(\frac{1}{\sqrt{2}}\right)^{3p^2-4p+1}.$$

Proof: To compute $ABC_1RII(R_p)$, we see that

$$\begin{aligned}
 ABC_1RII(T_p) &= \prod_{uv \in E(G)} \sqrt{\frac{r_G(u) + r_G(v) - 2}{r_G(u)r_G(v)}} \\
 &= \left(\sqrt{\frac{3+3-2}{3 \times 3}}\right)^6 \times \left(\sqrt{\frac{3+2-2}{3 \times 2}}\right)^{8(p-1)} \times \left(\sqrt{\frac{2+2-2}{2 \times 2}}\right)^{3p^2-4p+1} \\
 &= \left(\frac{2}{3}\right)^6 \times \left(\frac{1}{2}\right)^{4(p-1)} \times \left(\frac{1}{\sqrt{2}}\right)^{3p^2-4p+1}
 \end{aligned}$$

In the following theorem, we compute the multiplicative atom bond connectivity Revan index of T_p .

Theorem 8: The multiplicative geometric-arithmetical Revan index of a triangular benzenoid R_p is given by

$$GARI(T_p) = \left(\frac{2\sqrt{6}}{5}\right)^{8(p-1)}$$

Proof: To compute $GARI(R_p)$, we see that

$$\begin{aligned}
 GARI(R_p) &= \prod_{uv \in E(G)} \frac{2\sqrt{r_G(u)r_G(v)}}{r_G(u) + r_G(v)} \\
 &= \left(\frac{2\sqrt{3 \times 3}}{3+3}\right)^6 \times \left(\frac{2\sqrt{3 \times 2}}{3+2}\right)^{8(p-1)} \times \left(\frac{2\sqrt{2 \times 2}}{2+2}\right)^{3p^2-4p+1} \\
 &= \left(\frac{2\sqrt{6}}{5}\right)^{8(p-1)}
 \end{aligned}$$

4. RESULTS FOR BENZENOID HOURGLASS X_p

In this section, we consider the graph of benzenoid hourglass X_p which is obtained from two copies of a triangular benzenoid T_p by overlapping hexagons. The graph of benzenoid hourglass is shown in Figure 3.




Figure-3: The graph of benzenoid hourglass

Let G be the graph of a benzenoid hourglass X_p . By algebraic method, we obtain $|V(X_p)| = 2(p^2+4p - 2)$ and $|E(X_p)| = 3p^2 + 9p - 4$. It is easy to see that the vertices of benzenoid hourglass X_p are either of degree 2 or 3, see Figure 3. Therefore $\Delta(G)=3$ and $\delta(G)=2$. Thus $r_G(u) = 5 - d_G(u)$. By algebraic method, we obtain that the edge set $E(X_p)$ can be divided into three partitions:

$$\begin{aligned}
 E_{22} &= \{uv \in E(G) \mid d_G(u) = d_G(v) = 2\} & |E_{22}| &= 8 \\
 E_{23} &= \{uv \in E(G) \mid d_G(u) = 2, d_G(v) = 3\} & |E_{23}| &= 4(3p - 4). \\
 E_{33} &= \{uv \in E(G) \mid d_G(u) = d_G(v) = 3\} & |E_{33}| &= 3p^2 - 3p + 4.
 \end{aligned}$$

Thus there are three types of Revan edges as follows:

$$\begin{aligned}
 RE_{33} &= \{uv \in E(G) \mid r_G(u) = r_G(v) = 3\} & |RE_{33}| &= 8. \\
 RE_{32} &= \{uv \in E(G) \mid r_G(u) = 3, r_G(v) = 2\} & |RE_{32}| &= 4(3p - 4). \\
 RE_{22} &= \{uv \in E(G) \mid r_G(u) = r_G(v) = 2\} & |RE_{22}| &= 3p^2 - 3p + 4.
 \end{aligned}$$

We compute the multiplicative connectivity Revan indices of a benzenoid hourglass X_p .

Theorem 9: The multiplicative product connectivity Revan index of a benzenoid hourglass X_p is given by

$$PRII(X_p) = \left(\frac{1}{3}\right)^8 \times \left(\frac{1}{6}\right)^{2(3p-4)} \times \left(\frac{1}{2}\right)^{3p^2-3p+4}.$$

Proof: To compute $PRII(X_p)$, we see that

$$\begin{aligned} PRII(R_p) &= \prod_{uv \in E(G)} \frac{1}{\sqrt{r_G(u)r_G(v)}} \\ &= \left(\frac{1}{\sqrt{3 \times 3}}\right)^8 \times \left(\frac{1}{\sqrt{3 \times 2}}\right)^{4(3p-4)} \times \left(\frac{1}{\sqrt{2 \times 2}}\right)^{3p^2-3p+4} \\ &= \left(\frac{1}{3}\right)^8 \times \left(\frac{1}{6}\right)^{2(3p-4)} \times \left(\frac{1}{2}\right)^{3p^2-3p+4}. \end{aligned}$$

Theorem 10: The multiplicative sum connectivity Revan index of a benzenoid hourglass X_p is given by

$$SRII(X_p) = \left(\frac{1}{6}\right)^4 \times \left(\frac{1}{5}\right)^{2(3p-4)} \times \left(\frac{1}{2}\right)^{3p^2-3p+4}.$$

Proof: To compute $SRII(X_p)$, we see that

$$\begin{aligned} SRII(X_p) &= \prod_{uv \in E(G)} \frac{1}{\sqrt{r_G(u) + r_G(v)}} \\ &= \left(\frac{1}{\sqrt{3+3}}\right)^8 \times \left(\frac{1}{\sqrt{3+2}}\right)^{4(3p-4)} \times \left(\frac{1}{\sqrt{2+2}}\right)^{3p^2-3p+4} \\ &= \left(\frac{1}{6}\right)^4 \times \left(\frac{1}{5}\right)^{2(3p-4)} \times \left(\frac{1}{2}\right)^{3p^2-3p+4}. \end{aligned}$$

Theorem 11: The first multiplicative atom bond connectivity Revan index of a benzenoid hourglass X_p is given by

$$ABC_1RII(X_p) = \left(\frac{2}{3}\right)^8 \times \left(\frac{1}{2}\right)^{2(3p-4)} \times \left(\frac{1}{\sqrt{2}}\right)^{3p^2-2p+4}.$$

Proof: To compute $ABC_1RII(X_p)$, we see that

$$\begin{aligned} ABC_1RII(X_p) &= \prod_{uv \in E(G)} \sqrt{\frac{r_G(u) + r_G(v) - 2}{r_G(u)r_G(v)}} \\ &= \left(\sqrt{\frac{3+3-2}{3 \times 3}}\right)^8 \times \left(\sqrt{\frac{3+2-2}{3 \times 2}}\right)^{4(3p-4)} \times \left(\sqrt{\frac{2+2-2}{2 \times 2}}\right)^{3p^2-3p+4} \\ &= \left(\frac{2}{3}\right)^8 \times \left(\frac{1}{2}\right)^{2(3p-4)} \times \left(\frac{1}{\sqrt{2}}\right)^{3p^2-4p+4}. \end{aligned}$$

Theorem 12: The multiplicative geometric-arithmetic Revan index of a benzenoid hourglass X_p is given by

$$GARII(T_p) = \left(\frac{2\sqrt{6}}{5}\right)^{4(3p-4)}.$$

Proof: To compute $GARII(X_p)$, we see that

$$GARII(X_p) = \prod_{uv \in E(G)} \frac{2\sqrt{r_G(u)r_G(v)}}{r_G(u) + r_G(v)}.$$

$$\begin{aligned}
 &= \left(\frac{2\sqrt{3 \times 3}}{3+3} \right)^8 \times \left(\frac{2\sqrt{3 \times 2}}{3+2} \right)^{4(3p-4)} \times \left(\frac{2\sqrt{2 \times 2}}{2+2} \right)^{3p^2-3p+4} \\
 &= \left(\frac{2\sqrt{6}}{5} \right)^{4(3p-4)}.
 \end{aligned}$$

REFERENCES

1. V.R.Kulli, *Collegiate Graph Theory*, Vishwa International Publications, Gulbarga, India (2012).
2. V.R. Kulli, On multiplicative connectivity indices of certain nanotubes, *Annals of Pure and Applied Mathematics*, 12(2) (2016), 169-176.
3. V.R. Kulli, First multiplicative K Banhatti index and coindex of graphs, *Annals of Pure and Applied Mathematics*, 11(2) (2016), 79-82.
4. V.R. Kulli, On multiplicative K -Banhatti and multiplicative K hyper-Banhatti indices of V -Phenylenic nanotubes and nanotorus, *Annals of Pure and Applied Mathematics*, 11(2) (2016), 145-150.
5. V.R.Kulli, Multiplicative hyper-Zagreb indices and coindices of graphs: Computing these indices of some nanostructures, *International Research Journal of Pure Algebra*, 6(7), (2016) 342-347.
6. V.R.Kulli, Multiplicative connectivity indices of nanostructures, *Journal of Ultra Scientist of Physical Sciences*, A 29(1) (2017) 1-10.
7. V.R. Kulli, Multiplicative connectivity indices of $TUC_4C_8[m,n]$ and $TUC_4[m,n]$ nanotubes, *Journal of Computer and Mathematical Sciences*, 7(11) (2016) 599-605.
8. V.R.Kulli, Two new multiplicative atom bond connectivity indices, *Annals of Pure and Applied Mathematics*, 13(1) (2017) 1-7.
9. V.R.Kulli, Some new multiplicative geometric-arithmetic indices, *Journal of Ultra Scientist of Physical Sciences*, A, 29(2) (2017) 52-57.
10. V.R.Kulli, A new multiplicative arithmetic-geometric index, *International Journal of Fuzzy Mathematical Archive*, 12(2) (2017) 49-53.
11. V.R.Kulli, New multiplicative inverse sum indeg index of certain benzenoid systems, *Journal of Global Research in Mathematical Archives*, 4(10) (2017) 15-19.
12. V.R.Kulli, Edge version of multiplicative connectivity indices of some nanotubes and nanotorus, *International Journal of Current Research in Science and Technology*, 3(11) (2017) 7-15.
13. V.R. Kulli, Multiplicative Revan and multiplicative hyper-Revan indices of certain networks, *Journal of Computer and Mathematical Sciences*, 8(12) (2017) 750-757.
14. V.R. Kulli, Multiplicative connectivity Revan indices of polycyclic aromatic hydrocarbons and benzenoid systems, submitted.
15. V.R. Kulli, Multiplicative product connectivity and multiplicative sum connectivity indices of dendrimer nanostars, submitted.
16. V.R. Kulli, General multiplicative Revan indices of polycyclic aromatic hydrocarbons and benzenoid systems, submitted.
17. V.R. Kulli, On the product connectivity reverse index of silicate and hexagonal networks, *International Journal of Mathematics and its Applications*, 5(4-B) (2017) 175-179.
18. V.R. Kulli, On the sum connectivity reverse index of oxide and honeycomb networks, *Journal of Computer and Mathematical Sciences*, 8(9) (2017) 408-413.
19. V.R. Kulli, Atom bond connectivity reverse and product connectivity reverse indices of oxide and honeycomb networks, *International Journal of Fuzzy Mathematical Archive*, 15(1) (2018) 1-5.
20. V.R. Kulli, The sum connectivity Revan index of silicate and hexagonal networks, *Annals of Pure and Applied Mathematics*, 14(3) (2017) 401-406.
21. V.R. Kulli, On the product connectivity Revan index of certain nanotubes, *Journal of Computer and Mathematical Sciences*, 8(10) (2017) 562-567.
22. V.R.Kulli, B.Chaluvvaraju and H.S.Boregowda, Some degree based connectivity indices of Kulli cycle windmill graphs, *South Asian Journal of Mathematics*, 6(6) (2016) 263-268.
23. V.R. Kulli, General topological indices of some dendrimer nanostars, *Journal of Global Research in Mathematical Archives*, 4(11) (2017) 83-90.
24. A.R.Ashrafi and P. Nikzad, Connectivity index of the family of dendrimer nanostars, *Digest Journal of Nanomaterials and Biostructures*, 4(2) (2009) 269-273.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]