

ARITHMETIC LABELING OF $C_m \times P_n$ AND $P_m \times P_n$

STEPHEN JOHN. B^{*1} AND JOVITVINISHMELMA.G²

¹Associate Professor & ²Research Scholar,
 Department of Mathematics; AnnaiVelankanni College;
 Tholayavattam; Kanniyakumari District; Tamilnadu, India, 629 157.

(Received On: 19-02-18; Revised & Accepted On: 16-03-18)

ABSTRACT

Acharya.B.D. and Hedge.S.M. was introduced the concept of arithmetic labeling and many research articles have published in this topic. In this paper, we have proved that the Cartesian product of $C_m \times P_n$ (where m is odd) and $P_m \times P_n$ are arithmetic graphs. Also we established a general formula to find the labeling the vertices of the graph G .

Key words: Cartesian product, Labeling, Arithmetic Labeling.

INTRODUCTION

In 1989 Acharya.B.D. and Hedge.S.M. Introduced a new version of sequential graph known as arithmetic graph and is defined as follows: Let G is a graph with q edges a and d are the positive integers. A labeling f of G is said to be (a, d) – arithmetic if the vertices are labeled by distinct nonnegative integers and the edge labels are induced by $f(x) + f(y)$ for each xy are in the form of $a, a + d, a + 2d, \dots, a + (q - 1)d$. A graph is called arithmetic if it is an (a, d) - arithmetic for some a and d .

Definition 1.1: A graph G is an ordered pair $(V(G), E(G))$ consisting of a non empty set $V(G)$ of vertices and a set $E(G)$, disjoint from $V(G)$, of edges, together with an incidence function ψ_G that associates with each edge of G is an unordered pair of (not necessarily distinct) vertices of G .

Definition 1.2: Walk is an alternating sequence of vertices and edges starting and ending with vertices.

A walk in which all the vertices are distinct is called a **path**.

A closed path is called a **cycle**.

Definition 1.3: The cartesian product of the graphs G and H is denoted by $G \times H$ and defined the vertex set of $G \times H$ is the cartesian product $V(G) \times V(H)$ and any two vertices (u, u') and (v, v') are adjacent in $G \times H$ if and only if either $u = v$ and u' is adjacent with v' in H or $u' = v'$ and u is adjacent with v in G .

Definition 1.4: A labeling or valuation of a graph G is an assignment f of labels to the vertices of G that induces for each edge xy a label depending on the vertex labels $f(x)$ and $f(y)$.

Definition 1.5: A graph is said to be arithmetic if its vertices can be assigned distinct non negative integers in such a way that the value of the edges are obtained as sum of the values assigned in an arithmetic progression.

Lemma 2.1: Let G be the graph consisting of a cycle $C_{2t+1} = (u_1, u_2, \dots, u_{2t+1}, u_1)$ and a path $P_3 = (u_{2t-1}, w, u_{2t+1})$. Then G is an $(t + 2, 1)$, $t \geq 1$ arithmetic graph.

Lemma 2.2: The graph $P_m * P_n$ is an $(2y + 1, 2)$ - arithmetic graph for all m, n and for any non negative integer y . where $y = f(v_{1,1})$

Theorem 2.3: Let the graph G is the cartesian product of C_m and P_n . That is $G = C_m \times P_n$ Where $m, n > 0$ and m is odd is an $(a, 1)$ - arithmetic graph.

Proof: Let $G = C_m \times P_n$ where $m, n > 0$ and m is odd. Then the graph is given in (figure: 1) as below.

Corresponding Author: Stephen John. B^{*1}




Figure: 1

Let V be the vertex set of G and is denoted by $V(G) = \{V_{ij}/1 \leq i \leq n; 1 \leq j \leq m\}$

Define $f: V(G) \rightarrow N$. Now we are giving the label to the vertices of G as below

$$\begin{aligned} f(v_{2r-1,2k-1}) &= m((2r-1)-1) + k \text{ where } 1 \leq k \leq \frac{m+1}{2}; \\ 1 \leq r \leq \frac{n+1}{2} \forall \text{ odd } n \text{ (Or) } 1 \leq r \leq \frac{n}{2} \forall \text{ even } n \end{aligned} \quad (1)$$

$$\begin{aligned} f(v_{2r-1,2k}) &= m((2r-1)-1) + \left(\frac{m+1}{2}\right) + k \text{ where } 1 \leq k \leq \frac{m-1}{2} \\ 1 \leq r \leq \frac{n+1}{2} \forall \text{ odd } n \text{ (Or) } 1 \leq r \leq \frac{n}{2} \forall \text{ even } n \end{aligned} \quad (2)$$

$$\begin{aligned} f(v_{2r,2k-1}) &= m(2r) - \left(\frac{m-1}{2}\right) + (k-1) \text{ where } 1 \leq k \leq \frac{m+1}{2}; \\ 1 \leq r \leq \frac{n-1}{2} \forall \text{ odd } n \text{ (Or) } 1 \leq r \leq \frac{n}{2} \forall \text{ even } n \end{aligned} \quad (3)$$

$$\begin{aligned} f(v_{2r,2k}) &= m(2r) + k \text{ where } 1 \leq k \leq \frac{m-1}{2}; \\ 1 \leq r \leq \frac{n-1}{2} \forall \text{ odd } n \text{ (Or) } 1 \leq r \leq \frac{n}{2} \forall \text{ even } n \end{aligned} \quad (4)$$

Clearly all the vertices of $G = C_m \times P_n$ which are labeled by distinct labels from 1 to mn . The edge labels induced by $f(uv) = f(u) + f(v)$ for each uv are

$$a, a+d, a+2d, \dots, a+(q-1)d.$$

Therefore, the graph $G = C_m \times P_n$ is an $(a, 1)$ - Arithmetic graph.

Example 2.4: Consider the graph $G = C_7 \times P_5$




Figure: 2

Here, $m = 7$; $n = 5$ and $q = 63$

The vertex labels are given below.

$$\text{Equation (1)} \Rightarrow f(v_{2r-1,2k-1}) = m((2r-1)-1) + k$$

$$\text{where } 1 \leq k \leq \frac{m+1}{2}; 1 \leq r \leq \frac{n+1}{2} \forall \text{ odd } n$$

$$\text{When } r = 1,2,3 \text{ and } k = 1,2,3,4 \Rightarrow f(v_{1,1}) = 1; f(v_{1,3}) = 2;$$

$$f(v_{1,5}) = 3; f(v_{1,7}) = 4; f(v_{3,1}) = 15; f(v_{3,3}) = 16; f(v_{3,5}) = 17;$$

$$f(v_{3,7}) = 18; f(v_{5,1}) = 29; f(v_{5,3}) = 30; f(v_{5,5}) = 31; f(v_{5,7}) = 32.$$

$$\text{Equation (2)} \Rightarrow f(v_{2r-1,2k}) = m((2r-1)-1) + \left(\frac{m+1}{2}\right) + k$$

$$\text{where } 1 \leq k \leq \frac{m-1}{2}; 1 \leq r \leq \frac{n+1}{2} \forall \text{ odd } n.$$

$$\text{When } r = 1,2,3 \text{ and } k = 1,2,3 \Rightarrow f(v_{1,2}) = 5; f(v_{1,4}) = 6; f(v_{1,6}) = 7;$$

$$f(v_{3,2}) = 19; f(v_{3,4}) = 20; f(v_{3,6}) = 21; f(v_{5,2}) = 33;$$

$$f(v_{5,4}) = 34; f(v_{5,6}) = 35.$$

$$\text{Equation (3)} \Rightarrow f(v_{2r,2k-1}) = m(2r) - \left(\frac{m-1}{2}\right) + (k-1) \text{ where } 1 \leq k \leq \frac{m+1}{2}; 1 \leq r \leq \frac{n-1}{2} \forall \text{ odd } n$$

$$\text{When } r = 1,2 \text{ and } k = 1,2,3,4 \Rightarrow f(v_{2,1}) = 11; f(v_{2,3}) = 12;$$

$$f(v_{2,5}) = 13; f(v_{2,7}) = 14; f(v_{4,1}) = 25; f(v_{4,3}) = 26; f(v_{4,5}) = 27;$$

$$f(v_{4,7}) = 28.$$

$$\text{Equation (4)} \Rightarrow f(v_{2r,2k}) = m(2r-1) + k \text{ where } 1 \leq k \leq \frac{m-1}{2}; 1 \leq r \leq \frac{n-1}{2} \forall \text{ odd } n$$

$$\text{When } r = 1,2 \text{ and } k = 1,2,3 \Rightarrow f(v_{2,2}) = 8; f(v_{2,4}) = 9;$$

$$f(v_{2,6}) = 10; f(v_{4,2}) = 22; f(v_{4,4}) = 23; f(v_{4,6}) = 24.$$

In this graph, $a = 5$ and $d = 6 - 5 = 1$.

The edge labels are in the arithmetic progression

$$\begin{aligned} a &= 5, a + d = 6, a + 2d = 7, a + 3d = 8, a + 4d = 9, a + 5d = 10, \\ a + 6d &= 11, a + 7d = 12, a + 8d = 13, a + 9d = 14, a + 10d = 15, \\ a + 11d &= 16, a + 12d = 17, a + 13d = 18, a + 14d = 19, \\ a + 15d &= 20, a + 16d = 21, a + 17d = 22, a + 18d = 23, \\ a + 19d &= 24, a + 20d = 25, a + 21d = 26, a + 22d = 27, \\ a + 23d &= 28, a + 24d = 29, a + 25d = 30, a + 26d = 31, \\ a + 27d &= 32, a + 28d = 33, a + 29d = 34, a + 30d = 35, \\ a + 31d &= 36, a + 32d = 37, a + 33d = 38, a + 34d = 39, \\ a + 35d &= 40, a + 36d = 41, a + 37d = 42, a + 38d = 43, \\ a + 39d &= 44, a + 40d = 45, a + 41d = 46, a + 42d = 47, \\ a + 43d &= 48, a + 44d = 49, a + 45d = 50, a + 46d = 51, \\ a + 47d &= 52, a + 48d = 53, a + 49d = 54, a + 50d = 55, \\ a + 51d &= 56, a + 52d = 57, a + 53d = 58, a + 54d = 59, \\ a + 55d &= 60, a + 56d = 61, a + 57d = 62, a + 58d = 63, \\ a + 59d &= 64, a + 60d = 65, a + 61d = 66, \\ a + (q - 1)d &= a + 62d = 67. \end{aligned}$$

Then the graph $G = C_7 \times P_5$ is an $(5,1)$ - arithmetic graph.

Corollary 2.5: The graph $G = C_{2t+1} \times P_n$ is an $(t+2(f(v_{1,1})), 1)$ - arithmetic graph.

Proof: Let the graph G is the cartesian product of C_{2t+1} and $P_n \forall t, n > 0$. Then the vertex labeling of G is given below.

$$\begin{aligned} f(v_{1,1}) &= \text{Any one positive integer.} \\ f(v_{2r-1,2k-1}) &= m((2r-1)-1) + k + (f(v_{1,1}) - 1) \\ &\quad \text{where } 1 \leq k \leq \frac{m+1}{2}; 1 \leq r \leq \frac{n+1}{2} \forall \text{ odd } n. \\ (\text{Or}) 1 \leq r \leq \frac{n}{2} \forall \text{ even } n \end{aligned} \tag{1}$$

$$\begin{aligned} f(v_{2r-1,2k}) &= m((2r-1)-1) + \left(\frac{m+1}{2}\right) + k + (f(v_{1,1}) - 1) \\ &\quad \text{where } 1 \leq k \leq \frac{m-1}{2}; 1 \leq r \leq \frac{n+1}{2} \forall \text{ odd } n. \\ (\text{Or}) 1 \leq r \leq \frac{n}{2} \forall \text{ even } n \end{aligned} \tag{2}$$

$$\begin{aligned} f(v_{2r,2k-1}) &= m(2r) - \left(\frac{m-1}{2}\right) + (k-1) + (f(v_{1,1}) - 1) \\ &\quad \text{where } 1 \leq k \leq \frac{m+1}{2}; 1 \leq r \leq \frac{n-1}{2} \forall \text{ odd } n. \\ (\text{or}) 1 \leq r \leq \frac{n}{2} \forall \text{ even } n \end{aligned} \tag{3}$$

$$\begin{aligned} f(v_{2r,2k}) &= m(2r-1) + k + (f(v_{1,1}) - 1) \\ &\quad \text{where } 1 \leq k \leq \frac{m-1}{2}; 1 \leq r \leq \frac{n-1}{2} \forall \text{ odd } n \\ (\text{Or}) 1 \leq r \leq \frac{n}{2} \forall \text{ even } n \end{aligned} \tag{4}$$

Clearly all the vertices which are denoted by distinct labels from $f(v_{1,1})$ to $(mn + f(v_{1,1}) - 1)$.

The edge labels induced by $f(uv) = f(u) + f(v)$ for each uv are $a, a + d, a + 2d, \dots, a + (q - 1)d$.

Then the graph $G = C_{2t+1} \times P_n$ is an $(t + 2(f(v_{1,1})), 1)$ - arithmetic graph.

In this graph $a = 6$ and $d = 7 - 6 = 1$

The edge labels are in the arithmetic progression

$$\begin{aligned} a &= 6, a + d = 7, a + 2d = 8, a + 3d = 9, a + 4d = 10, a + 5d = 11, \\ a + 6d &= 12, a + 7d = 13, a + 8d = 14, a + 9d = 15, a + 10d = 16, \\ a + 11d &= 17, a + 12d = 18, a + 13d = 19, a + 14d = 20, \\ a + 15d &= 21, a + 16d = 22, a + 17d = 23, a + 18d = 24, \\ a + 19d &= 25, a + 20d = 26, a + 21d = 27, a + 22d = 28, \\ a + 23d &= 29, a + 24d = 30, a + 25d = 31, a + 26d = 32, \\ a + 27d &= 33, a + 28d = 34, a + 29d = 35, a + 30d = 36, \\ a + 31d &= 37, a + 32d = 38, a + 33d = 39, \\ a + (q - 1)d &= a + 34d = 40. \end{aligned}$$

Here $t = 2$, $(t + 2 (f(v_{1,1})), 1) = (6, 1)$.

Therefore, the graph $G = C_5 \times P_4$ is an $(6, 1)$ - arithmetic graph.

Theorem 2.7: Let the graph G is the cartesian product of P_m and P_n . That is $G = P_m \times P_n$ where $m, n > 0$ is an $(a, 2)$ - arithmetic graph.

Proof: Let $G = P_m \times P_n$ where $m, n > 0$. Then the graph is given in the (figure: 4) as below.




Figure: 4

Let V be the vertex set of G and is denoted by $V(G) = \{v_{ij} / 1 \leq i \leq n; 1 \leq j \leq m\}$

Define $f: V(G) \rightarrow N$.

Now we are giving the label to the vertices of G as below:

$$\begin{aligned} f(v_{ij}) &= (2m - 1)(i - 1) + j \\ \text{where } 1 &\leq i \leq n; 1 \leq j \leq m. \end{aligned}$$

Clearly all the vertices of $G = P_m \times P_n$ are labeled by distinct labels from 1 to $q + 1$.

The edge labels induced by $f(uv) = f(u) + f(v)$ are as follows

$$\begin{aligned} f(v_{i,k}v_{i,k+1}) &= (2k + 1) + 2(2m - 1)(i - 1) \\ \text{where } 1 &\leq i \leq n; 1 \leq k \leq m - 1. \end{aligned}$$

$$f(v_{r,j}v_{r+1,j}) = 2m + (2j - 1) + 2(2m - 1)(r - 1) \\ \text{where } 1 \leq r \leq n - 1; 1 \leq j \leq m.$$

Clearly the edges are labeled as $f(E(G)) = \left\{ \begin{matrix} a, a + d, a + 2d, \dots, \\ a + (q - 1)d \end{matrix} \right\}$.

Therefore, f is an arithmetic labeling.

Hence the graph $G = P_m \times P_n$ is an $(a, 2)$ - arithmetic graph.

Example 2.8: Consider the graph $G = P_4 \times P_4$




Figure: 5

Here, $m = 4; n = 4$ and $q = 24$

The vertex labels are given below.

$$f(v_{ij}) = (2m - 1)(i - 1) + j \text{ Where } 1 \leq i \leq n; 1 \leq j \leq m.$$

$$\begin{aligned} \text{When } i = 1, 2, 3, 4 \text{ and } j = 1, 2, 3, 4 \Rightarrow & f(v_{1,1}) = 1; f(v_{1,2}) = 2; f(v_{1,3}) = 3; \\ & f(v_{1,4}) = 4; f(v_{2,1}) = 8; f(v_{2,2}) = 9; f(v_{2,3}) = 10; f(v_{2,4}) = 11; \\ & f(v_{3,1}) = 15; f(v_{3,2}) = 16; f(v_{3,3}) = 17; f(v_{3,4}) = 18; \\ & f(v_{4,1}) = 22; f(v_{4,2}) = 23; f(v_{4,3}) = 24; f(v_{4,4}) = 25. \end{aligned}$$

The edge labels are given below.

$$f(v_{i,k}v_{i,k+1}) = (2k + 1) + 2(2m - 1)(i - 1) \\ \text{where } 1 \leq i \leq n; 1 \leq k \leq m - 1$$

$$\begin{aligned} \text{When } i = 1, 2, 3, 4 \text{ and } k = 1, 2, 3 \Rightarrow & f(v_{1,1}v_{1,2}) = 3; f(v_{1,2}v_{1,3}) = 5; f(v_{1,3}v_{1,4}) = 7; \\ & f(v_{2,1}v_{2,2}) = 17; f(v_{2,2}v_{2,3}) = 19; f(v_{2,3}v_{2,4}) = 21; \\ & f(v_{3,1}v_{3,2}) = 31; f(v_{3,2}v_{3,3}) = 33; f(v_{3,3}v_{3,4}) = 35; \\ & f(v_{4,1}v_{4,2}) = 45; f(v_{4,2}v_{4,3}) = 47; f(v_{4,3}v_{4,4}) = 49. \\ & f(v_{r,j}v_{r+1,j}) = 2m + (2j - 1) + 2(2m - 1)(r - 1) \\ & \text{where } 1 \leq r \leq n - 1; 1 \leq j \leq m \end{aligned}$$

$$\begin{aligned} \text{When } r = 1, 2, 3 \text{ and } j = 1, 2, 3, 4 \Rightarrow & f(v_{1,1}v_{2,1}) = 9; f(v_{2,1}v_{3,1}) = 23; \\ & f(v_{3,1}v_{4,1}) = 37; f(v_{1,2}v_{2,2}) = 11; f(v_{2,2}v_{3,2}) = 25; \\ & f(v_{3,2}v_{4,2}) = 39; f(v_{1,3}v_{2,3}) = 13; f(v_{2,3}v_{3,3}) = 27; \\ & f(v_{3,3}v_{4,3}) = 41; f(v_{1,4}v_{2,4}) = 15; f(v_{2,4}v_{3,4}) = 29; \\ & f(v_{3,4}v_{4,4}) = 43 \end{aligned}$$

In this graph $a = 3$ and $d = 5 - 3 = 2$

The edge labels are in the arithmetic progression

$$\begin{aligned} a &= 3, a + d = 5, a + 2d = 7, a + 3d = 9, a + 4d = 11, \\ a + 5d &= 13, a + 6d = 15, a + 7d = 17, a + 8d = 19, a + 9d = 21, \\ a + 10d &= 23, a + 11d = 25, a + 12d = 27, a + 13d = 29, \\ a + 14d &= 31, a + 15d = 33, a + 16d = 35, a + 17d = 37, \\ a + 18d &= 39, a + 19d = 41, a + 20d = 43, a + 21d = 45, \\ a + 22d &= 47, a + (q - 1)d = a + 23d = 49. \end{aligned}$$

Therefore, the graph $G = P_4 \times P_4$ is an $(3,2)$ - arithmetic graph.

Corollary 2.9: The graph $G = P_m \times P_n$ is an $(2y+1,2)$ - arithmetic graph. Where $y = f(v_{1,1})$.

Proof: Let the graph G is the cartesian product of P_m and $P_n \forall m, n > 0$. Then the vertex labeling of G is given below.

$$\begin{aligned} f(v_{1,1}) &= \text{any one positive integer} \\ f(v_{ij}) &= (2m - 1)(i - 1) + j + (f(v_{1,1}) - 1) \\ &\text{where } 1 \leq i \leq n; 1 \leq j \leq m \end{aligned}$$

Clearly all the vertices which are denoted by distinct labels from

$$f(v_{1,1}) \text{ to } q + f(v_{1,1}).$$

The edge labeling of G is given as below:

$$\begin{aligned} f(v_{i,k}v_{i,k+1}) &= (2k + 1) + 2(2m - 1)(i - 1) + 2(f(v_{1,1}) - 1) \\ &\text{where } 1 \leq i \leq n; 1 \leq k \leq m - 1 \end{aligned}$$

$$\begin{aligned} f(v_{r,j}v_{r+1,j}) &= 2m + (2j - 1) + 2(2m - 1)(r - 1) + 2(f(v_{1,1}) - 1) \\ &\text{where } 1 \leq r \leq n - 1; 1 \leq j \leq m \end{aligned}$$

The induced edge labels are $f(E(G)) = \{a, a + d, a + 2d, a + 3d, \dots, a + (q - 1)d\}$. Then f is an arithmetic labeling.

Hence the graph $G = P_m \times P_n$ is an $(2y + 1, 2)$ - arithmetic graph. Where $y = f(v_{1,1})$.

Example 2.10: Consider the graph $G = P_5 \times P_4$




Figure: 6

Here, $m = 5; n = 4$ and $q = 31$

The vertex labels are given below

Let $f(v_{1,1}) = 3$

$$\begin{aligned} f(v_{ij}) &= (2m - 1)(i - 1) + j + (f(v_{1,1}) - 1) \\ &\text{where } 1 \leq i \leq n; 1 \leq j \leq m. \end{aligned}$$

When $i = 1, 2, 3, 4$ and $j = 1, 2, 3, 4, 5 \Rightarrow f(v_{1,2}) = 4; f(v_{1,3}) = 5; f(v_{1,4}) = 6;$
 $f(v_{1,5}) = 7; f(v_{2,1}) = 12; f(v_{2,2}) = 13; f(v_{2,3}) = 14;$
 $f(v_{2,4}) = 15; f(v_{2,5}) = 16; f(v_{3,1}) = 21; f(v_{3,2}) = 22;$
 $f(v_{3,3}) = 23; f(v_{3,4}) = 24; f(v_{3,5}) = 25; f(v_{4,1}) = 30;$
 $f(v_{4,2}) = 31; f(v_{4,3}) = 32; f(v_{4,4}) = 33; f(v_{4,5}) = 34.$

The edge labels are given below.

$$f(v_{i,k}v_{i,k+1}) = (2k + 1) + 2(2m - 1)(i - 1) + 2(f(v_{1,1}) - 1)$$

where $1 \leq i \leq n; 1 \leq k \leq m - 1$

When $i = 1, 2, 3, 4$ and $k = 1, 2, 3, 4 \Rightarrow f(v_{1,1}v_{1,2}) = 7; f(v_{1,2}v_{1,3}) = 9;$
 $f(v_{1,3}v_{1,4}) = 11; f(v_{1,4}v_{1,5}) = 13; f(v_{2,1}v_{2,2}) = 25; f(v_{2,2}v_{2,3}) = 27;$
 $f(v_{2,3}v_{2,4}) = 29; f(v_{2,4}v_{2,5}) = 31; f(v_{3,1}v_{3,2}) = 43; f(v_{3,2}v_{3,3}) = 45;$
 $f(v_{3,3}v_{3,4}) = 47; f(v_{3,4}v_{3,5}) = 49; f(v_{4,1}v_{4,2}) = 61; f(v_{4,2}v_{4,3}) = 63;$
 $f(v_{4,3}v_{4,4}) = 65; f(v_{4,4}v_{4,5}) = 67.$
 $f(v_{r,j}v_{r+1,j}) = 2m + (2j - 1) + 2(2m - 1)(r - 1) + 2(f(1,1) - 1)$

Where $1 \leq r \leq n - 1; 1 \leq j \leq m$

When $r = 1, 2, 3$ and $j = 1, 2, 3, 4, 5 \Rightarrow f(v_{1,1}v_{2,1}) = 15; f(v_{2,1}v_{3,1}) = 33;$
 $f(v_{3,1}v_{4,1}) = 51; f(v_{1,2}v_{2,2}) = 17; f(v_{2,2}v_{3,2}) = 35; f(v_{3,2}v_{4,2}) = 53;$
 $f(v_{1,3}v_{2,3}) = 19; f(v_{2,3}v_{3,3}) = 37; f(v_{3,3}v_{4,3}) = 55; f(v_{1,4}v_{2,4}) = 21;$
 $f(v_{2,4}v_{3,4}) = 39; f(v_{3,4}v_{4,4}) = 57; f(v_{1,5}v_{2,5}) = 23; f(v_{2,5}v_{3,5}) = 41;$
 $f(v_{3,5}v_{4,5}) = 59.$

In this graph $a = 7$ and $d = 9 - 7 = 2$.

The edge labels are in the arithmetic progression

$$\begin{aligned} a &= 7, a + d = 9, a + 2d = 11, a + 3d = 13, a + 4d = 15, a + 5d = 17, \\ a + 6d &= 19, a + 7d = 21, a + 8d = 23, a + 9d = 25, a + 10d = 27, \\ a + 11d &= 29, a + 12d = 31, a + 13d = 33, a + 14d = 35, \\ a + 15d &= 37, a + 16d = 39, a + 17d = 41, a + 18d = 43, a + 19d = 45, \\ a + 20d &= 47, a + 21d = 49, a + 22d = 51, a + 23d = 53, a + 24d = 55, \\ a + 25d &= 57, a + 26d = 59, a + 27d = 61, a + 28d = 63, a + 29d = 65, \\ a + (q - 1)d &= a + 30d = 67. \end{aligned}$$

Here $y = 3, (2y + 1, 2) = (7, 2)$

Therefore, the graph $G = C_5 \times P_4$ is an $(7, 2)$ - arithmetic graph.

CONCLUSION

In this paper we have studied about arithmetic labeling of $C_m \times P_n \forall$ odd m and $P_m \times P_n$. We assure that it will help to the new researches in this area.

REFERENCE

1. Acharya .B.D. and Hedge .S.M., Arithmetic Graphs, Journ.Graph Theory, 14(3) (1989), 275-299.
2. Augustinovich .S. and Fon-Der-Flaass. D, Cartesian Products of Graphs and Metric Spaces, Europ. J. Combinatorics (2000) 21, 847-851 Article No. 10.1006 leujc. 2000.0401.
3. Grace .T., On sequential labeling of Graphs Jour of Graph Theory, 7-195-201 (1983).
4. Harary .F., Graph Theory, Narosa Publishing House (1969).
5. Joseph .A. Gallian, A Dynamic Survey of Graph Labeling, The Electronic Journal of Combinatorics, (2013).
6. Rosa .A., On Certain Valuations of The Vertices of A Graph Theory of Graphs (Internet, Symposium, Rome, July 1966), Gordon and Dunod Paris (1967)
7. Suresh Singh .G., A note on labeling of Graphs, Graphs And Combinatorics (1998), 201-207.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]