

μ - α -SEMI GENERALIZED CLOSED SETS IN GENERALIZED TOPOLOGICAL SPACES

SARANYA M*

Assistant Professor, Department of Mathematics,
AJK College of Arts and Science, Coimbatore, Tamil Nadu, India.

(Received On: 27-01-18; Revised & Accepted On: 09-03-18)

ABSTRACT

In this paper, I introduce a new class of sets in generalized topological spaces called μ - α -semi generalized closed sets. Also I investigate some of their basic properties and obtained some interesting theorems.

Keywords: Generalized topological spaces, μ - α -semi closed sets, μ - α -semi generalized closed sets.

1. INTRODUCTION

The concept of generalized topological spaces was introduced and investigated by A. Csaszar [1]. Many μ -closed sets like μ -semi closed sets, μ -pre closed sets etc., in generalized topological spaces are introduced by him. In this paper, I introduce a new class of sets in generalized topological spaces called μ - α -semi generalized closed sets. Also I investigate some of their basic properties and produced many interesting theorems.

2. PRELIMINARIES

Definition 2.1: [1] Let X be a nonempty set. A collection μ of subsets of X is a generalized topology (or briefly GT) on X if it satisfies the following:

1. $\emptyset, X \in \mu$ and
2. If $\{M_i : i \in I\} \subseteq \mu$, then $\cup_{i \in I} M_i \in \mu$.

If μ is a GT on X , then (X, μ) is called a generalized topological space (or briefly GTS) and the elements of μ are called μ -open sets and their complement are called μ -closed sets.

Definition 2.2: [1] Let (X, μ) be a GTS and let $A \subseteq X$. Then the μ -closure of A , denoted by $c_\mu(A)$, is the intersection of all μ -closed sets containing A .

Definition 2.3: [1] Let (X, μ) be a GTS and let $A \subseteq X$. Then the μ -interior of A , denoted by $i_\mu(A)$, is the union of all μ -open sets contained in A .

Definition 2.4: [1] Let (X, μ) be a GTS. A subset A of X is said to be

1. μ -semi-closed set if $i_\mu(c_\mu(A)) \subseteq A$
2. μ -pre-closed set if $c_\mu(i_\mu(A)) \subseteq A$
3. μ - α -closed set if $c_\mu(i_\mu(c_\mu(A))) \subseteq A$
4. μ - β -closed set if $i_\mu(c_\mu(i_\mu(A))) \subseteq A$
5. μ -regular-closed set if $A = c_\mu(i_\mu(A))$

Definition 2.5: [3] Let (X, μ) be a GTS. A subset A of X is said to be

1. μ -regular generalized closed set if $c_\mu(A) \subseteq U$ whenever $A \subseteq U$, where U is μ -regular open in X
2. μ -generalized closed set if $c_\mu(A) \subseteq U$ whenever $A \subseteq U$, where U is μ -open in X
3. μ -generalized- α -closed set if $\alpha c_\mu(A) \subseteq U$ whenever $A \subseteq U$, where U is μ -open in X

The complement of μ -semi closed (resp., μ -pre closed, μ - α -closed, μ - β -closed, μ -regular closed, μ -generalized closed) is said to be μ -semi open (resp., μ -pre open, μ - α -open, μ - β -open, μ -regular open, μ -generalized open) in X .

Corresponding Author: Saranya M*
Assistant Professor, Department of Mathematics,
AJK College of Arts and Science, Coimbatore, Tamil Nadu, India.

3. μ - α -SEMI GENERALIZED CLOSED SETS

In this section I introduce μ - α -semi generalized closed sets in generalized topological spaces and studied some of their basic properties. Some interesting and important theorems are also obtained.

Definition 3.1: Let (X, μ) be a GTS. Then a non-empty subset A is said to be a μ - α -semi generalized closed set (briefly μ - α -SGCS) if $sc_\mu(A) \subseteq U$ whenever $A \subseteq U$ and U is μ - α -open in X .

Example 3.2: Let $X = \{a, b, c\}$ and let $\mu = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$. Then (X, μ) is a GTS. Now,
 μ - α O(X) = $\{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$.

Then $A = \{a\}$ is a μ - α -semi generalized closed set in (X, μ) .

Theorem 3.3: Every μ -closed set in (X, μ) is a μ - α -semi generalized closed set in (X, μ) but not conversely in general.

Proof: Let A be a μ -closed set in (X, μ) , then $c_\mu(A) = A$. Now let $A \subseteq U$ and U be μ - α -open in X . Then $sc_\mu(A) = A \cup i_\mu(c_\mu(A)) = A \cup i_\mu(A) \subseteq A \subseteq U$, by hypothesis. Therefore $sc_\mu(A) \subseteq U$. This implies, A is a μ - α -semi generalized closed set in (X, μ) .

Example 3.4: Let $X = \{a, b, c, d\}$ and let $\mu = \{\emptyset, \{b\}, \{d\}, \{b, d\}, X\}$. Then (X, μ) is a GTS. Now,
 μ - α O(X) = $\{\emptyset, \{b\}, \{d\}, \{b, d\}, X\}$.

Then $A = \{b\}$ is a μ - α -semi generalized closed set in (X, μ) . But, as $c_\mu(A) = c_\mu(\{b\}) = \{a, b, c\} \neq A$, A is not a μ -closed set in (X, μ) .

Theorem 3.5: Every μ -semi closed set in (X, μ) is a μ - α -semi generalized closed set in (X, μ) .

Proof: Let A be a μ -semi closed set in (X, μ) . Then $i_\mu(c_\mu(A)) \subseteq A$. Now let $A \subseteq U$ and U be μ - α -open in X . Then $sc_\mu(A) = A \cup i_\mu(c_\mu(A)) \subseteq A \cup A = A \subseteq U$, by hypothesis. Therefore A is a μ - α -semi generalized closed set in (X, μ) .

Remark 3.6: Every μ - α -semi generalized closed sets and μ -pre closed sets are independent in general in (X, μ) .

Example 3.7: Let $X = \{a, b, c\}$ and let $\mu = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$. Then (X, μ) is a GTS. Now,
 μ - α O(X) = $\{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$.

Then $A = \{a\}$ is a μ - α -semi generalized closed set but not a μ -pre closed set as $c_\mu(i_\mu(A)) = c_\mu(i_\mu(\{a\})) = \{a, c\} \not\subseteq A$.

Example 3.8: Let $X = \{a, b, c\}$ and let $\mu = \{\emptyset, \{a, b\}, X\}$. Then (X, μ) is a GTS. Now let $A = \{a\}$. Then $c_\mu(i_\mu(A)) = c_\mu(i_\mu(\{a\})) = \emptyset \subseteq A$. Therefore A is a μ -pre closed set, but A is not a μ - α -semi generalized closed set as $A \subseteq U = \{a, b\}$, where U is a μ - α -open set and now,

$$\mu$$
- α O(X) = $\{\emptyset, \{a, b\}, X\}$ and $sc_\mu(A) = X \not\subseteq \{a\} = U$.

Theorem 3.9: Every μ - α -closed set in (X, μ) is a μ - α -semi generalized closed set in (X, μ) but not conversely in general.

Proof: Let A be a μ - α -closed set in (X, μ) . Then $c_\mu(i_\mu(c_\mu(A))) \subseteq A$. Now let $A \subseteq U$ and U be μ - α -open in X . Then $sc_\mu(A) = A \cup i_\mu(c_\mu(A)) \subseteq A \cup c_\mu(i_\mu(c_\mu(A))) \subseteq A \cup A = A \subseteq U$, by hypothesis. Therefore $sc_\mu(A) \subseteq U$. This implies, A is a μ - α -semi generalized closed set in (X, μ) .

Example 3.10: Let $X = \{a, b, c\}$ and let $\mu = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, c\}, X\}$. Then (X, μ) is a GTS. Now,
 μ - α O(X) = $\{\emptyset, \{a\}, \{b\}, \{c\}, \{a, c\}, X\}$.

Then $A = \{c\}$ is a μ - α -semi generalized closed set in (X, μ) . But, A is not a μ - α -closed set in X , as $c_\mu(i_\mu(c_\mu(A))) = c_\mu(i_\mu(c_\mu(\{c\}))) = \{a, c\} \not\subseteq A$.

Remark 3.11: A μ - β -closed set is not a μ - α -semi generalized closed set in (X, μ) in general.

Example 3.12: Let $X = \{a, b, c\}$ and let $\mu = \{\emptyset, \{a, b\}, X\}$. Then (X, μ) is a GTS. Let $A = \{a\}$. Then, $i_\mu(c_\mu(i_\mu(A))) = i_\mu(c_\mu(i_\mu(\{a\}))) = \emptyset \subseteq \{a\} = A$. Therefore A is a μ - β -closed set in (X, μ) , but not a μ - α -semi generalized closed set as $A \subseteq U = \{a, b\}$, where U is a μ - α -open set and now,

$$\mu$$
- α O(X) = $\{\emptyset, \{a, b\}, X\}$ and $sc_\mu(A) = X \not\subseteq \{a\} = U$.

Theorem 3.13: Every μ -regular closed set in (X, μ) is a μ - α -semi generalized closed set in (X, μ) but not conversely.

Proof: Let A be a μ -regular closed set in (X, μ) . Then $A = c_\mu(i_\mu(A))$. Now let $A \subseteq U$ and U be μ - α -open in X . Then $sc_\mu(A) = A \cup i_\mu(c_\mu(A)) = A \cup i_\mu(c_\mu(i_\mu(A))) = A \cup i_\mu(c_\mu(i_\mu(A))) = A \cup c_\mu(i_\mu(A)) = A \cup A = A \subseteq U$, by hypothesis. Therefore A is a μ - α -semi generalized closed set in X .

Example 3.14: Let $X = \{a, b, c, d\}$ and let $\mu = \{\emptyset, \{a\}, \{c\}, \{b, d\}, X\}$. Then (X, μ) is a GTS.

Now, $\mu\text{-}\alpha O(X) = \{\emptyset, \{a\}, \{c\}, \{b, d\}, X\}$. Then $A = \{c\}$ is a μ - α - semi generalized closed set but not a μ -regular closed set as $c_\mu(i_\mu(A)) = c_\mu(i_\mu(\{c\})) = \{a, c\} \neq \{c\} = A$.

Remark 3.15: Every μ - α -semi generalized closed set and μ -generalized closed set in (X, μ) are independent to each other in general.

Example 3.16: Let $X = \{a, b, c, d\}$ and let $\mu = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}, X\}$. Then (X, μ) is a GTS. Now,

$$\mu\text{-}\alpha O(X) = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}, X\}.$$

Then $A = \{a\}$ is a μ - α - semi generalized closed set in (X, μ) , but not a μ -generalized closed set as $c_\mu(A) = c_\mu(\{a\}) = \{a, d\} \not\subseteq \{a, b, c\} = U$.

Example 3.17: Let $X = \{a, b, c\}$ and let $\mu = \{\emptyset, \{a, b\}, X\}$. Then (X, μ) is a GTS. Let $A = \{a, b\}$ and $U = X$. Then $A \subseteq U$ and $c_\mu(A) = c_\mu(\{a, b\}) = X \subseteq U$. Therefore A is a μ -generalized closed set, but not a μ - α - semi generalized closed set as $A \subseteq U = \{a, b\}$, where U is a μ - α -open set and now,

$$\mu\text{-}\alpha O(X) = \{\emptyset, \{a, b\}, X\} \text{ and } sc_\mu(A) = X \not\subseteq \{a, b\} = U.$$

Remark 3.18: Every μ - α - semi generalized closed set and μ -generalized- α -closed set in (X, μ) are independent to each other in general.


Example 3.19: Let $X = \{a, b, c, d\}$ and let $\mu = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$. Then (X, μ) is a GTS.

Now, $\mu\text{-}\alpha O(X) = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}, X\}$. Then $A = \{b\}$ is a μ - α - semi generalized closed set in (X, μ) , but not a μ -generalized- α -closed set as $ac_\mu(A) = \{b, c, d\} \not\subseteq \{a, b, c\} = U$ and $A \subseteq U$.

Example 3.20: Let $X = \{a, b, c\}$ and let $\mu = \{\emptyset, \{a, b\}, X\}$. Then (X, μ) is a GTS. Let $A = \{a\}$ and $U = X$. Then $A \subseteq U$ and $ac_\mu(A) = A \cup c_\mu(i_\mu(c_\mu(A))) = \{a\} \cup X = X \subseteq U$. Therefore A is a μ -generalized- α -closed set, but not a μ - α - semi generalized closed set as $A \subseteq U = \{a\}$ where U is a μ - α -open set and now,

$$\mu\text{-}\alpha O(X) = \{\emptyset, \{a, b\}, X\} \text{ and } sc_\mu(A) = X \not\subseteq \{a\} = U.$$

In the following diagram, we have provided relations between various types of μ -closed sets.



Remark 3.21: Union of any two μ - α -semi generalized closed sets in (X, μ) need not be a μ - α - semi generalized closed set in X .

Example 3.22: Let $X = \{a, b, c\}$ and let $\mu = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$. Then (X, μ) is a GTS. Now,

$$\mu\text{-}\alpha O(X) = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}.$$

Then $A = \{a\}$ and $B = \{b\}$ are μ - α - semi generalized closed sets in (X, μ) . But, $A \cup B = \{a, b\}$ is not a μ - α - semi generalized closed set as $sc_\mu(\{a, b\}) = \{a, b\} \cup i_\mu(c_\mu(\{a, b\})) = \{a, b\} \cup X = X \not\subseteq \{a, b\} = U$ and $A \cup B \subseteq U$.

Theorem 3.23: If a subset A in X is a μ - α - semi generalized closed set in (X, μ) , then $sc_\mu(A) - A$ contains no non-empty μ - α -closed set in X .

Proof: Let A be a μ - α - semi generalized closed set in (X, μ) and let $sc_\mu(A) - A$ contains a non-empty μ - α -closed set say F . That is, $F \subseteq sc_\mu(A) - A$. This implies, $F \subseteq sc_\mu(A) \cap A^c$ and hence $F \subseteq sc_\mu(A) \& F \subseteq A^c$ (1)

Now $F \subseteq A^c$ implies $A \subseteq F^c$. Since F^c is μ - α -open and A is a μ - α - semi generalized closed set, we have, $sc_\mu(A) \subseteq F^c$. This implies $F \subseteq (sc_\mu(A))^c$ (2)

From (1) & (2), $F \subseteq sc_\mu(A)$ and $F \subseteq (sc_\mu(A))^c$. This implies $F \subseteq sc_\mu(A) \cap (sc_\mu(A))^c = \emptyset$. Therefore $F = \emptyset$. Thus $sc_\mu(A) - A$ contains no non-empty μ - α -closed set in X .

Theorem 3.24: Let (X, μ) be a GTS. Then for every $A \in sc_\mu(X)$ and for every set $B \subseteq X$, $A \subseteq B \subseteq sc_\mu(A)$ implies $B \in sc_\mu(X)$.

Proof: Let $B \subseteq U$ and U be a μ - α -open set in (X, μ) . Then since $A \subseteq B$ and $A \subseteq U$. By hypothesis $B \subseteq sc_\mu(A)$. Therefore $sc_\mu(B) \subseteq sc_\mu(sc_\mu(A)) = sc_\mu(A) \subseteq U$ as A is a μ - α - semi generalized closed set of X . Hence $B \in sc_\mu(X)$.

Theorem 3.25: In a GTS X , for each $x \in X$, $\{x\}$ is a μ - α -closed set or its complement $X - \{x\}$ is a μ - α - semi generalized closed set in (X, μ) .

Proof: Suppose that $\{x\}$ is not a μ - α closed set in (X, μ) . Then $X - \{x\}$ is not a μ - α -open set in (X, μ) . The only μ - α -open set containing $X - \{x\}$ is X . Thus $X - \{x\} \subseteq X$ and so $sc_\mu(X - \{x\}) \subseteq sc_\mu(X) = X$. Therefore $sc_\mu(X - \{x\}) \subseteq X$ and so $X - \{x\}$ is a μ - α - semi generalized closed set in (X, μ) .

Theorem 3.26: If A is both a μ - α -open set and a μ - α - semi generalized closed set in (X, μ) , then A is a μ -semi closed set in (X, μ) .

Proof: Let A be μ - α -open and μ - α - semi generalized closed set in (X, μ) . Then, $sc_\mu(A) \subseteq A$ as $A \subseteq A$. But always $A \subseteq sc_\mu(A)$. Therefore, $A = sc_\mu(A)$. Hence A is a μ -semi closed set in (X, μ) .

Theorem 3.27: If $A \subseteq Y \subseteq X$ and A is a μ - α - semi generalized closed set in X then A is a μ - α - semi generalized closed set relative to Y .

Proof: Given that $A \subseteq Y \subseteq X$ and A is a μ - α - semi generalized closed set in X . Let $A \subseteq Y \cap U$, where U is a μ - α -open set in X . Since A is a μ - α - semi generalized closed set, $A \subseteq U$ implies $sc_\mu(A) \subseteq U$. This implies $Y \cap sc_\mu(A) \subseteq Y \cap U$ and $sc_\mu(A) \subseteq Y \cap U$. That is A is a μ - α - semi generalized closed set relative to Y .

Theorem 3.28: Let A be any μ - α - semi generalized closed set in (X, μ) . Then A is a μ -semi closed set in (X, μ) iff $sc_\mu(A) - A$ is a μ - α closed set in X .

Proof: Necessity: Let A be a μ -semi closed set in (X, μ) . Then $sc_\mu(A) = A$ and so, $sc_\mu(A) \cap A^c = A \cap A^c = \emptyset$. Therefore, $sc_\mu(A) \cap A^c = \emptyset$ and $sc_\mu(A) - A = \emptyset$, Therefore $sc_\mu(A) - A$ is a μ - α -closed set in (X, μ) .

Sufficiency: Let $sc_\mu(A) - A$ be a μ - α - closed set and A be a μ - α -semi generalized closed set in X . Then by Theorem 3.23, $sc_\mu(A) - A$ does not contain any non-empty μ - α -closed set and hence $sc_\mu(A) - A = \emptyset$. That is $sc_\mu(A) = A$. Hence A is a μ -semi closed set in (X, μ) .

Theorem 3.29: Every subset of X is a μ - α - semi generalized closed set in X iff every μ - α -open set is a μ -semi closed set in X .

Proof: Necessity: Let A be μ - α -open in X , and by hypothesis, A is a μ - α - semi generalized closed set in X . Hence by Theorem 3.26, A is a μ -semi closed set in X .

Sufficiency: Let $A \subseteq U$ where U is a μ - α -open set in X . Then by hypothesis, U is a μ -semi closed set. This implies $sc_\mu(U) = U$ and $sc_\mu(A) \subseteq sc_\mu(U) = U$. Hence $sc_\mu(A) \subseteq U$. Thus A is a μ - α - semi generalized closed set in X .

Theorem 3.30: Let A and B be μ - α - semi generalized closed set in (X, μ) such that $c_\mu(A) = sc_\mu(A)$ and $c_\mu(B) = sc_\mu(B)$, then $A \cup B$ is a μ - α - semi generalized closed set in X .

Proof: Let $A \cup B \subseteq U$, where U is a μ - α -open set. Then $A \subseteq U$ and $B \subseteq U$. Since A and B are μ - α - semi generalized closed sets, $sc_\mu(A) \subseteq U$ and $sc_\mu(B) \subseteq U$. Now $sc_\mu(A \cup B) \subseteq c_\mu(A \cup B) = c_\mu(A) \cup c_\mu(B) = sc_\mu(A) \cup sc_\mu(B) \subseteq U \cup U = U$. Hence $A \cup B$ is μ - β - semi generalized closed set in X .

4. REFERENCES

1. Csaszar, A., Generalized open sets in generalized topologies, Acta Mathematica Hungaria 96(2002), 351-357.
2. Csaszar, A., Generalized open sets in generalized topologies, Acta Mathematica Hungaria 106(2005), 53-66.
3. Fritzie Mae Vasquez Valenzuela and Helen Moso Rara., μ -rgb-sets in a generalized topological spaces, International Journal of Mathematical Analysis-Vol.8, 2014, 1791-1797.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]