

**STOLARSKY3- MEAN LABELING
OF ARBITRARY SUPER SUBDIVISIONS OF SOME STANDARD GRAPHS**

S. S. SANDHYA¹, E. EBIN RAJA MERLY² AND S. KAVITHA³

¹Department of Mathematics,
Sree Ayyappa College for Women,
Chunkankadai-629 003, Kanyakumari District, Tamilnadu, India.

²Department of Mathematics,
Nesamony Memorial Christian College,
Marthandam-629 165, Kanyakumari District, Tamilnadu, India.

³Department of Mathematics,
Holy Cross College, Nagercoil -629 004, Kanyakumari District, Tamilnadu, India.

E-mail: sssandhya2009@gmail.com¹, ebinmerly@gmail.com² and kavis.maths@gmail.com³

ABSTRACT

The Concept of Stolarsky-3 Mean labeling was already introduced. In this paper we introduced Arbitrary Super Subdivision in stolarsky-3 Mean graph. Further we investigate Arbitrary Super Subdivisions of Path, Cycle, H-graph ($n \leq 10$ if n is even and $n \leq 11$ if n is odd) and Middle graphs.

Keywords: Graph Labeling, Stolarsky-3 mean labeling, Super subdivision of graphs, Arbitrary Super subdivision of graphs, H graph, Middle graph.

1. INTRODUCTION

The graph $G = (V, E)$ is considered here will be finite, simple and undirected. We follow Gallian[1] for all detailed survey of graph labeling and we refer Harary [2] for all other standard terminologies and notations. The concept of “Mean labeling” has been introduced by S.Somasundaram and R.Ponraj in 2004[7]. The concept of “Harmonic Mean Labeling” has been introduced by S. Somasundaram, R. Ponraj and S.S. Sandhya in [8]. The concept of “Stolarsky-3 Mean Labeling” has introduced by S.S. Sandhya, E.Ebin Raja Merely and S.Kavitha in [6]. G. Sethuraman and P.Selvaraju have introduced a new method of construction called Super subdivision of graphs and prove that arbitrary super subdivisions of paths are graceful in [9]. Motivating these results we investigate Stolarsky-3 Mean labeling of Arbitrary super subdivisions of some standard graphs. Following definitions are useful for the present investigations.

Definition 1.1: A graph G with p vertices and q edges is said to be Stolarsky-3 Mean graph if each vertex $x \in V$ is assigned distinct labels $f(x)$ from $1, 2, \dots, q+1$ and each edge $e=uv$ is assigned the distinct labels

$f(e=uv) = \left\lfloor \sqrt{\frac{f(u)^2 + f(u)f(v) + f(v)^2}{3}} \right\rfloor$ (or) $\left\lfloor \sqrt{\frac{f(u)^2 + f(u)f(v) + f(v)^2}{3}} \right\rfloor$ then the resulting edge labels are distinct. In this case f is called a Stolarsky-3 Mean labeling of G .

Definition 1.2: A walk in which all the vertices u_1, u_2, \dots, u_n are distinct is called a path. It is denoted by P_n .

Definition 1.3: A closed path is called a cycle. A cycle on n vertices is denoted by C_n .

Definition 1.4: The H-graph of a path P_n is the graph obtained from two copies of P_n with vertices $v_1, v_2, v_3, \dots, v_n$ and u_1, u_2, \dots, u_n by joining the vertices $v_{\frac{n+1}{2}}$ and $u_{\frac{n+1}{2}}$ if n is odd and the vertices $v_{\frac{n}{2}+1}$ and $u_{\frac{n}{2}}$ if n is even.

Definition 1.5: The Middle graph $M(G)$ of a graph G is the graph whose vertex set is $V(G) \cup E(G)$ and in which two vertices are adjacent if and only if either they are adjacent edges of G or one is a vertex of G and the other is an edge incident with it.

Definition 1.6: Let G be a graph. A graph H is called a super subdivision of G if H is obtained from G by replacing every edge e_i of G by a complete bipartite graph k_{2,m_i} for some $m_i, 1 \leq i \leq q$ in such a way that the ends of e_i are merged with the two vertices part of k_{2,m_i} after removing the edge e_i from graph G .

Definition 1.7: A Super subdivision H of G is said to be an Arbitrary super subdivision of G if every edge of G is replaced by an arbitrary k_{2,m_i} where m_i may vary for each edge arbitrarily. It is denoted by ASS (G).

2. MAIN RESULTS

Theorem 2.1: Arbitrary Super subdivisions of path are Stolarsky- 3 mean graphs.

Proof: Let P_n be the path with successive vertices $u_1, u_2, u_3, \dots, u_n$ and let e_i denote the edge $u_i u_{i+1}$ of P_n for $1 \leq i \leq n-1$

Let H be an arbitrary super subdivision of a path P_n , where each edge e_i of P_n is replaced by a complete bipartite graph k_{2,m_i} where m_i is any positive integer.

$$V(H) = \{u_1, u_2, u_3, \dots, u_n, v_1, v_2, v_3, \dots, v_{m_1}, v_{m_1+1}, v_{m_1+2}, \dots, v_{m_1+m_2}, \dots, v_{m_1+m_2+m_3}, \dots, v_{m_1+m_2+m_3+\dots+m_{n-1}}\}$$




Figure-1: Arbitrary Super subdivision of P_n

Here we consider two different cases

Case-(i): $m_1=m_2=m_3= \dots =m_{n-1} = 4$




Figure-2: Arbitrary Super subdivision of P_4

Define $\varphi: V(H) \rightarrow \{1, 2, \dots, q+1\}$ by
 $\varphi(u_1) = 1.$
 $\varphi(u_i) = 8(i-1), 2 \leq i \leq n.$
 $\varphi(v_1) = 2.$
 $\varphi(v_i) = 2i-1, 2 \leq i \leq m_1 + m_2 + \dots + m_{n-1}.$

Then the edge labels are distinct.

In this case H is Stolarsky-3 mean graph.

Case-(ii): $m_1=m_2=m_3= \dots =m_{n-1}=5$




Figure-3: Arbitrary Super subdivision of P_4

Define $\varphi: V(H) \rightarrow \{1, 2, \dots, q+1\}$ by

$$\varphi(u_1) = 1.$$

$$\varphi(u_i) = 10(i-1), 2 \leq i \leq n.$$

$$\varphi(v_1) = 2.$$

$$\varphi(v_i) = 2i-1, 2 \leq i \leq m_1 + m_2 + \dots + m_{n-1}.$$

Then the edge labels are distinct.

From case (i) and (ii) we can conclude that Arbitrary super subdivisions of path are Stolarsky-3 mean graphs.

Remark 2.2: The above result is true for all values of n 's and m_i 's with the condition $m_1 \leq 8$ and m_i 's, $2 \leq i \leq n - 1$.

Theorem 2.3: Arbitrary Super subdivisions of cycle are Stolarsky- 3 mean graphs.

Proof: Let C_n be a cycle with consecutive vertices $u_1, u_2, u_3, \dots, u_n$ and let e_i denote the edge $u_{i-1}u_i$ of C_n for $1 \leq i \leq n-1$.

Let H be an arbitrary super subdivision of a cycle C_n , where each edge e_i of C_n is replaced by a complete bipartite graph K_{2,m_i} where m_i is any positive integer.

$$V(H) = \{u_1, u_2, u_3, \dots, u_n, v_1, v_2, v_3, \dots, v_{m_1}, v_{m_1+1}, v_{m_1+2}, \dots, v_{m_1+m_2}, \dots, v_{m_1+m_2+m_3}, \dots, v_{m_1+m_2+m_3+\dots+m_{n-1}}\}$$




Figure-4: Arbitrary Super subdivision of C_n

Case-(i): $m_1=m_2=m_3= \dots =m_{n-1} = 3$




Figure 5: Arbitrary Super subdivision of C_4

Define $\varphi: V(H) \rightarrow \{1, 2, \dots, q+1\}$ by

$$\varphi(u_1) = 1.$$

$$\varphi(u_i) = 6i-5, 2 \leq i \leq n.$$

$$\varphi(v_i) = 2i, 1 \leq i \leq m_1 + m_2 + \dots + m_{n-1}.$$

$$\varphi(v_{m_1+m_2+\dots+m_{n-1}+1}) = \varphi(u_n) + 1.$$

Then the edge labels are distinct.

In this case H is Stolarsky-3 mean graph.

Case-(ii): $m_1=m_2=m_3=\dots=m_{n-1} = 4$




Figure-6: Arbitrary Super subdivision of C_5

Define $\varphi: V(H) \rightarrow \{1, 2, \dots, q+1\}$ by

$$\varphi(u_1) = 1.$$

$$\varphi(u_i) = 8i-7, 2 \leq i \leq n.$$

$$\varphi(v_i) = 2i, \quad 1 \leq i \leq m_1 + m_2 + \dots + m_{n-1}.$$

$$\varphi(v_{m_1+m_2+\dots+m_{n-1}+1}) = \varphi(u_n) + 1.$$

Then the edge labels are distinct.

From case (i) and (ii) we can conclude that Arbitrary Super subdivisions of cycle are Stolarsky-3 mean graphs.

Remark 2.4: The above result is true for all values of n's and m_i's with the condition m₁ ≤ 4 and m_i's, 2 ≤ i ≤ n - 1.

Theorem 2.5: Arbitrary Super subdivision of H-graph is Stolarsky-3 mean graph (n ≤ 11 if n is odd, n ≤ 10 if n is even).

Proof: Let G be a H-graph with vertices v₁, v₂, ..., v_n and u₁, u₂, ..., u_n and let u_iu_{i+1}, v_iv_{i+1}, 1 ≤ i ≤ n - 1, $\frac{v_{n+1}u_{n+1}}{2}$ if n is odd and $\frac{v_{\frac{n}{2}+1}u_{\frac{n}{2}}}{2}$ if n is even be the edges of a H-graph G.

Let H be an Arbitrary super subdivision of a H-graph G, where each edge e_i of G is replaced by a complete bipartite graph k_{2,m_i} where m_i is any positive integer.

$$V(H) = \{v_1, v_2, v_3, \dots, v_n, u_1, u_2, u_3, \dots, u_n, W_1, W_2, \dots, W_{m_1},$$

$$W_{m_1+1}, W_{m_1+2}, \dots, W_{m_1+m_2}, W_{m_1+m_2+m_3+\dots+m_{n-1}}, X_1, X_2, X_3, \dots, X_{m_1},$$

$$X_{m_1+1}, X_{m_1+2}, \dots, X_{m_1+m_2}, X_{m_1+m_2+m_3+\dots+m_{n-1}}, X_{m_k}\}$$




Figure-7: Arbitrary Super Subdivision of H- graph (n is Odd)




Figure-8: Arbitrary Super Subdivision of H- graph (n is Even)

Case-(i): When n is odd ($n \leq 11$) and $m_1=m_2=m_3= \dots =m_{n-1}= 3$




Figure-9: Arbitrary Super subdivision of H- graph with n=5

Define a function $\varphi: V(H) \rightarrow \{1, 2, \dots, q+1\}$ by

$$\varphi(v_1) = 1.$$

$$\varphi(v_i) = 6(i-1), 2 \leq i \leq n.$$

$$\varphi(u_1) = \varphi(v_n) + 1.$$

$$\varphi(u_i) = \varphi(u_{i-1}) + 6, 1 \leq i < \frac{n+1}{2} \& \left(\frac{n+1}{2}\right) + 1 \leq i \leq n.$$

$$\varphi(u_i) = \varphi(u_{i-1}) + 7, i = \frac{n+1}{2}.$$

$$\varphi(w_1) = 2.$$

$$\varphi(w_i) = 2i-1, 2 \leq i \leq m_1+m_2+m_3+\dots+m_{n-1}.$$

$$\varphi(x_1) = \varphi(u_1) + 1.$$

$$\varphi(x_i) = \varphi(x_{i-1}) + 2, 2 \leq i \leq m_1+m_2 \& m_1+m_2+2 \leq i \leq m_1+m_2+m_3+\dots+m_{n-1}$$

$$\varphi(x_{(m_1+m_2)+1}) = \varphi\left(u_{\frac{n+1}{2}}\right) + 1.$$

$$\varphi(x_{(m_k)}) = \varphi(x_{m_1+m_2}) + 1.$$

Then the edge labels are distinct.

Case-(ii): When n is even ($n \leq 10$) and $m_1=m_2=m_3=\dots=m_{n-1}=3$




Figure-10: Arbitrary Super subdivision of H-graph with $n=6$

Define a function $\varphi: V(H) \rightarrow \{1, 2, \dots, q+1\}$ by

$$\varphi(v_1) = 1.$$

$$\varphi(v_i) = 6(i-1), 2 \leq i \leq n.$$

$$\varphi(u_1) = \varphi(v_n) + 1.$$

$$\varphi(u_i) = \varphi(u_{i-1}) + 6, 2 \leq i < \frac{n}{2} \& \left(\frac{n}{2} + 1\right) + 1 \leq i \leq n.$$

$$\varphi(u_{\frac{n}{2}}) = \varphi(u_{\frac{n}{2}-1}) + 7.$$

$$\varphi(w_1) = 2.$$

$$\varphi(w_i) = 2i-1, 2 \leq i \leq m_1+m_2+m_3+\dots+m_{n-1}.$$

$$\varphi(x_1) = \varphi(w_{m_1+m_2+\dots+m_{n-1}}) + 3.$$

$$\varphi(x_i) = \varphi(x_{i-1}) + 2, 2 \leq i \leq m_1+m_2 \& m_1+m_2+2 \leq i \leq m_1+m_2+m_3+\dots+m_{n-1}.$$

$$\varphi(x_{m_1+m_2+1}) = \varphi(u_{\frac{n}{2}}) + 1.$$

$$\varphi(x_{(m_k)}) = \varphi(x_{m_1+m_2}) + 1.$$

Then the edge labels are distinct.

From case (i) and (ii), Arbitrary Super subdivision of H-graph is stolarsky-3 Mean graph.

Remark 2.6: The above result is true for all values of n's if n is odd ($n \leq 11$) and n is even ($n \leq 10$) and m_i 's with the condition $m_1 \leq 8$ and m_i 's, $2 \leq i \leq n-1$.

Theorem 2.7: Arbitrary Super Subdivision of Middle graph $M(P_n)$ is Stolarsky-3 Mean graph.

Proof: Let u_1, u_2, \dots, u_n be the vertices and e_1, e_2, \dots, e_{n-1} be the edges of path P_n and $G=M(P_n)$ be the middle graph of path P_n . $V(G) = V(P_n) \cup E(P_n)$ and let $u_i x_i$ ($1 \leq i \leq n-1$), $u_i x_{i-1}$, $2 \leq i \leq n$, $x_i x_{i+1}$, $1 \leq i \leq n-2$ be the edges of $M(P_n)$.

Let H be an arbitrary super subdivision of $M(P_n)$, where each edge of $M(P_n)$ is replaced by a complete bipartite graph k_{2, m_i} where m_i is any positive integer.

$$V(H) = \{u_1, u_2, \dots, u_n, x_1, x_2, \dots, x_{n-1}, v_1, v_2, \dots, v_{m_1}, v_{m_1+1}, v_{m_1+2}, \dots, v_{m_1+m_2}, v_{m_1+m_2+m_3}, \dots, v_{m_1+m_2+m_3+\dots+m_{2n-2}}, w_1^{(k)}, w_2^{(k)}, \dots, w_{n-2}^{(k)}\}.$$




Figure-11: Arbitrary Super subdivision of $M(P_n)$

When $m_1=m_2=m_3= \dots = m_{2n-2}=3$.




Figure-12: Arbitrary Super subdivision of $M(P_3)$

Define a function $\varphi: V(H) \rightarrow \{1, 2, \dots, q+1\}$ by

$$\begin{aligned} \varphi(u_1) &= 1. \\ \varphi(u_i) &= 14(i-1) - 2, 1 \leq i \leq n. \\ \varphi(x_1) &= 6. \\ \varphi(x_i) &= 14(i-1) + 6, 1 \leq i \leq n-1. \\ \varphi(v_1) &= \varphi(u_1) + 1. \\ \varphi(v_2) &= \varphi(v_1) + 1. \\ \varphi(v_i) &= \varphi(v_{i-1}) + 2, 3 \leq i \leq m_1 + m_2 + m_3 + \dots + m_{2n-2} \\ &\text{But } i \neq (m_1 + m_2) + 1, (m_1 + m_2 + m_3) + 1, \dots \\ &\quad (m_1 + m_2 + m_3 + \dots + m_{2n-3}) + 1 \\ \varphi(w_i^{(k)}) &= 14i + 5, 1 \leq i \leq n-2. \\ \varphi(v_{(\sum_{i=1}^k m_i)+1}) &= \varphi(v_{\sum_{i=1}^k m_i}) + 3, k = 2, 3, 4, \dots, 2n-3. \end{aligned}$$

Then the edge labels are distinct.

Hence Arbitrary Super subdivision of Middle graph $M(P_n)$ is stolarsky-3 Mean graph.

Remark 2.8: The above result is true for all values of n 's and m_i 's with the condition $m_1 \leq 8$ and m_i 's, $2 \leq i \leq n-1$.

CONCLUSION

In this paper we studied the stolarsky-3 Mean labeling behavior of Arbitrary Super subdivision of some standard graphs. The authors are of the opinion that the study of Stolarsky-3 Mean labeling of Arbitrary Super subdivision of some standard graphs shall be quite interesting and also will lead to new results.

REFERENCES

1. J.A. Gallian, "A dynamic survey of graph labeling", The electronic Journal of Combinatorics 17(2017), #DS6.
2. F.Harary, 1988, "Graph Theory" NarosaPuplishing House Reading, New Delhi.
3. K.M. Kathiresan and S. Amutha, Arbitrary Super subdivisions of stars are graceful, Indian J.Pure Appl.Math., 35(1) 81-84(2004).
4. V.Ramachandran and C.Sekar, One Modulo N Gracefulness of Arbitrary Super subdivisions of graphs, International J. Math. Combin. Vol.2 (2014), 36-46.
5. V.Ramachandran and C.Sekar, , Graceful labeling of Arbitrary Super subdivision of disconnected graph, Ultra Scientist of Physical Sciences, Vol. 25(2) A.Aug. 2013,315-318.
6. S.S.Sandhya, E. Ebin Raja Merly and S.Kavitha "Stolarsky-3 Mean Labeling of Graphs" Communicated to Journal of discrete Mathematical Sciences and Cryptography".
7. S.Somasundram, and R.Ponraj 2003 "Mean Labeling of Graphs", National Academy of Science Letters Vol. 26, p.210-213.
8. S.Somasundram, R.Ponraj and S.S.Sandhya, "Harmonic Mean Labeling of Graphs" communicated to Journal of Combinatorial Mathematics and combinational computing.
9. G.Sethuraman and P.Selvaraju, Gracefulness of arbitrary Super subdivisions of graphs, Indian J.Pure Appl. Math., 1059-1064 (2001).

Source of support: Proceedings of National Conference January 11-13, 2018, on Discrete & Computational Mathematics (NCDCM - 2018), Organized by Department of Mathematics, University of Kerala, Kariavattom Thiruvananthapuram-695581.