

**BRIDGE GRAPH RELATED FAMILIES OF E-CORDIAL GRAPHS**

**MUKUND V.BAPAT\***

**Mukund V. Bapat, Hindale, Devgad, Sindhudurg, Maharashtra India: 416630.**

*(Received On: 16-03-18; Revised & Accepted On: 17-04-18)*

**ABSTRACT**

*The two copies of graph  $G(p,q)$  are joined by  $t$  paths on  $n$ -points each. These paths are bridges between the two graphs. We represent the family by  $G(tP_n)$ . The paths are attached at the same fixed point on  $G$ . We discuss E-cordiality of  $C_3(P_n)$ ,  $K_4(P_n)$ ,  $S_4(P_n)$  (shel graph  $S_4$ ). We show that under certain conditions these graphs are E-cordial.*

**Key words:** graph, E-cordial, shel graph,  $C_3$ ,  $C_4$ .

**Subject Classification:** 05C78.

**1. INTRODUCTION**

In 1997 Yilmaz and Cahit introduced weaker version of edge graceful labeling E-cordial labeling [4] Let  $G$  be a  $(p,q)$ .  $f: E \rightarrow \{0,1\}$  Define  $f$  on  $V$  by  $f(v) = \sum \{f(vu)(vu) \in E(G)\} \pmod{2}$ . The function  $f$  is called as E-cordial labeling if  $|v_f(0)-v_f(1)| \leq 1$  and  $|e_f(0)-e_f(1)| \leq 1$  where  $v_f(i)$  is the number of vertices labeled with  $i=0, 1$ . And  $e_f(i)$  is the number of edges labeled with  $i=0, 1$ . We follow the convention that  $v_f(0, 1) = (a, b)$  for  $v_f(0) = a$  and  $v_f(1) = b$  further  $e_f(0,1) = (x, y)$  for  $e_f(0)=x$  and  $e_f(1) = y$ . A graph that admits E-cordial labeling is called as E-cordial graph. Yilmaz and Cahit prove that trees  $T_n$  are E-cordial iff for  $n$  not congruent to  $2 \pmod{4}$ ,  $K_n$  are E-cordial iff  $n$  not congruent to  $2 \pmod{4}$ , Fans  $F_n$  are E-cordial iff for  $n$  not congruent to  $1 \pmod{4}$ . Yilmaz and Cahit observe that A graph on  $n$  vertices cannot be E-cordial if  $n$  is congruent to  $2 \pmod{4}$ . One should refer Dynamic survey of graph labeling by Joe Gallian [2] for more results on E-cordial graphs. The graphs we consider are finite, undirected, simple and connected. For terminology and definitions we refer Harary[3] and Dynamic survey of graph labeling by Joe Gillian [2]. The families we discuss are obtained by taking two copies of graph  $G$  and join them by  $t$  paths of equal length. The paths are attached at the same fixed point on  $G$ . we represent these families by  $G(tP_n)$ . We take  $t = 2$  and  $G = C_3, S_4, K_4, C_4$ .


**2. PRELIMINARIES**

**2.1 Fusion of vertex.** Let  $G$  be a  $(p, q)$  graph. Let  $u \neq v$  be two vertices of  $G$ . We replace them with single vertex  $w$  and all edges incident with  $u$  and that with  $v$  are made incident with  $w$ . If a loop is formed is deleted. The new graph has at least  $p-1$  vertices and  $q-1$  edges. [5].

**3. THEOREMS PROVED**

**3.1 Theorem.**  $G = C_3(2P_n)$  is e-cordial iff  $n$  is odd number.

**Proof:** The ordinary labeling of  $G$  is shown below.



**Figure-3.1:** ordinary labeling of  $C_3(2P_n)$

**Corresponding Author: Mukund V.Bapat\***

**Mukund V. Bapat, Hindale, Devgad, Sindhudurg, Maharashtra India: 416630.**

The graph has  $2n+2$  vertices and  $2n+4$  edges.

Define  $f: E(G) \rightarrow \{0, 1\}$  as follows:

$f(c_i) = 0$  for  $i=1,3,5,\dots$  and  $i \leq n$ .


$f(c_i) = 1$  for  $i=2,4,6,\dots$  and  $i \leq n-1$ .

$f(c_i) = 1$  for  $i = n+1, n+2, \dots, n+\frac{n+3}{2}$ .

$f(c_i) = 0$  for all other  $i > \frac{3n+3}{2}$ . This gives vertex and label number distribution as  $v_f(0,1) = (n+1, n+1)$  and  $e_f(0,1) = (n+2, n+2)$ .

**3.2 Theorem.**  $G = C_4(2P_n)$  is e-cordial iff  $n$  is even number.

**Proof:**  $G$  has  $2n + 4$  vertices and  $2n+6$  edges. Define  $f: V(E) \rightarrow \{0,1\}$  as follows.



**Figure-3.2:** ordinary labeling of  $C_4(2P_n)$

$f(c_i) = 0$  for  $i=1,3,5,\dots$  and  $i \leq n+1$

$f(c_i) = 1$  for  $i=2,4,6,\dots$  and  $i \leq n$

$f(c_i) = 1$  for  $i=n+2, n+3,\dots, n+x+4$ .

$f(c_i) = 0$  for rest of  $i$ . The label number distribution is  $v_f(0,1) = (n+2, n+2)$  and  $e_f(0,1) = (n+3, n+3)$  as required.

**3.3 Theorem.**  $G = S_4(2P_n)$  is e-cordial iff  $n$  is even number.

**Proof:**  $G$  has  $2n + 4$  vertices and  $2n+8$  edges. Define  $f: V(E) \rightarrow \{0,1\}$  as follows.



**Figure-3.3:** ordinary labeling of  $S_4(2P_n)$

$f(c_i) = 1$  for  $i=1,2,3,4,5$


$f(c_{5+i}) = 0$  for  $i=2p+1$  and  $p = 0, 1, \dots, x-1$ .

$f(c_i) = 1$  for  $i=7,9,\dots, n+3$

$f(c_{n+3+i}) = 1$  for  $i=1, 2, 3,\dots, x$

$f(c_i) = 0$  for all rest of  $i$ .

The label number distribution is  $v_f(0, 1) = (n+2, n+2)$  and  $e_f(0, 1) = (n+4, n+4)$  as required. In the above graph two paths  $P_n$  are fused at vertex of degree 2 on  $S_4$ . If we change the vertex of fusion on  $S_4$  from degree two vertex to degree 3 vertex then also the same labeling function as above will do. This is shown in ordinary labeling in diagram below and with example in fig.3.5.



**Figure-3.4:** ordinary labeling of  $S_4(2P_n)$




Figure-3.5: labeling of  $S_4(2P_8)$

**3.4 Theorem.**  $G= K_4(2P_n)$  is e-cordial iff  $n$  is even number.

**Proof:**  $G$  has  $2n + 4$  vertices and  $2n+10$  edges. Define  $f: V(E) \rightarrow \{0,1\}$  as follows.




Figure-3.6: ordinary labeling of  $K_4(2P_n)$

$$f(c_i) = 1 \text{ for } i=1,2,3,4,5,6$$

$$f(c_{6+2i}) = 1 \text{ for } i=1, 2, \dots, \left(\frac{n}{2}-1\right)$$

$$f(c_{6+2i+1}) = 1 \text{ for } i=0, 1, 2, \dots, \left(\frac{n}{2}-1\right)$$

$$f(c_{n+5+i}) = 1 \text{ for } i=1, 2, 3, \dots, x$$

$$f(c_i) = 0 \text{ for all rest of } i.$$

The label number distribution is  $v_f(0,1) = (n+2, n+2)$  and  $e_f(0,1) = (n+5, n+5)$  as required.

**3.5 Theorem.**  $G= H(2P_n)$  is e-cordial iff  $n$  is odd number where  $H$  is house graph on  $C_4$

**Proof:**  $G$  has  $2n + 6$  vertices and  $2n+10$  edges. From figure below it follows that we can fuse the paths at three different points, 'a', 'b' and 'c' on  $H$ .




Figure-3.6: House graph  $H$  with vertices 'a', 'b', and 'c'




Figure-3.7: ordinary labeling of  $H(2P_n)$

Define  $f: V(E) \rightarrow \{0,1\}$  as follows.

$$f(c_1) = 1,$$

$$f(c_2) = 0,$$

$$f(c_3) = 0,$$

$$f(c_4) = 0,$$

$$f(c_5) = 1,$$

$$f(c_6) = 1$$

$$f(c_{6+i})=0 \text{ for } i=1, 3, 5, \dots, \frac{n-1}{2}$$

$$f(c_{7+i}) = 1 \text{ for } i=1, 3, \dots, \frac{n-1}{2}$$

$$f(c_{n+5+i})=1 \text{ for } i=1, 2, \dots, (x+3)$$

$f(c_i)=0$  for all rest of  $i$ . In the above graph we have taken double paths from vertex 'a' on one copy of  $H$  to vertex 'a' of other copy. In fact we need not restrict ourselves to build bridges from same vertex to same vertex. Above labeling will work to build bridges between cross vertices such as 'a' to 'b', 'b' to 'd' etc. Since there are multiple bridges between two vertices by deleting any bridge or edge there of will not add to the components of graph which is contrary to the established sense of bridge. Conclusions: A new family of graphs is obtained when two copies of given graph are joined by two paths of same length at the same vertex. We have shown that for  $G = C_3, C_4, S_4$  (shel graph) and  $K_4$  the graphs  $G(2P_m)$  are E-cordial. We also noticed that for house graph  $H$  the two paths need not be between same point on  $H$ . We can choose cross points still the cordiality will not affect. This is advantageous.

## REFERENCES

1. Bapat M.V. Ph.D. thesis, University Of Mumbai, 2004.
2. Joe Gallian Dynamic survey of graph labeling 2016.
3. Harary, Graph Theory, Narosa publishing, New Delhi.
4. Yilmaz, Cahit, E-cordial graphs, Ars combina, 46, 251-256.
5. Introduction to Graph Theory by D. WEST, Pearson Education Asia.

**Source of support: Nil, Conflict of interest: None Declared.**

***[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]***