

**SOLUTION OF EIGHTH ORDER BOUNDARY VALUE PROBLEMS
BY ONE DIMENSIONAL DIFFERENTIAL TRANSFORM METHOD**

FIDA HUSSAIN* AND MUHAMMAD ARSHAD

School of Electrical and Information Engineering, Jiangsu University, Zhenjiang, China.

(Received On: 15-04-18; Revised & Accepted On: 14-05-18)

ABSTRACT

One dimensional Differential Transformation Method (1-dim DTM) is used to find the solution of the eighth order boundary value problems. The approximate solution of the problem is obtained in the form of a rapid convergent series. To illustrate the ability, efficiency and reliability of the method, some examples are given, revealing its effectiveness and simplicity and the calculated results are compared with exact solution.

Keywords: Eighth order boundary value problem, one dimensional differential transform method, Series solution.

1. INTRODUCTION

Generally, the eighth-order boundary value problems (BVPs) are recognized to occur in Mathematics, Physics and Engineering Sciences [1, 2]. In the book written by Chandrasekhar [3], we find therein that whenever horizontal infinite layers of fluid are heated from below and those are then under the rotation-action, and hence instability develops. When the instability develops as ordinary convection, then that ordinary differential equation (ODE) is a sixth-order ODE. When the said instability sets as over stability, then it is to be modelled by eighth-order ODE. Bishop investigated eighth-order D.E for uniform beams for their torsional vibration. Over the years, some authors used different methods and worked on such types (eighth-order) boundary value problems [4, 5, 6, 7, 8, 9, 10, 11, 12]. Differential quadrature method [9], finite difference method (FDM) [4], spline method and modified decomposition method [10] are applicable to find the solution of eighth-order BVPs. But, these methods involve huge calculations and the solution is found at grid points only.

In 1986, Zhou and Pukhov have developed a so-called differential transformation method (DTM) for electrical circuit's problems [13]. The DTM is a technique that uses Taylor series for the solution of differential equations in the form of a polynomial. The Taylor series method is computationally unexciting for high order equations. The DTM leads to an iterative procedure for obtaining an analytic series solutions of functional equations. In recent years, many papers were devoted to the problem of approximate solution of system of differential equations. The implementation of the Differential Transform Method (DTM) [14, 15, 16] amongst others has shown reliable results to solve ordinary differential equations, partial differential equations, integral-differential equations, the Schrödinger equations, Analytic solution for Telegraph equation, Systems of Volterra Integral Equations of the First Kind, non-linear fractional differential equations and delay differential equations.

2. ONE-DIMENSIONAL DIFFERENTIAL TRANSFORMS METHOD

Definition:

One dimensional Differential transformation of function $y(x)$ is defined as follows [14]

$$Y(K) = \frac{1}{K!} \left[\frac{d^K y(x)}{dx^K} \right]_{x=0} \quad (1)$$

In equation (1), $y(x)$ is the original function and $Y(K)$ is the transformed function. One dimensional Differential inverse transform of $Y(K)$ is defined as

$$y(x) = \sum_{K=0}^{\infty} x^K Y(K) \quad (2)$$

**Corresponding Author: Fida Hussain*,
School of Electrical and Information Engineering, Jiangsu University, Zhenjiang, China.**

In fact, from (1) and (2), we obtain

$$y(x) = \sum_{K=0}^{\infty} \frac{x^K}{K!} \left[\frac{d^K y(x)}{dx^K} \right]_{x=0} \tag{3}$$

Equation (3) implies that the concept of differential transformation is derived from the Taylor series expansion. The differential transform's operators are shown in the table 1.

Table-1: Differential transform operators

Sr. No	Original function	Transform function
1	$u(x) \pm v(x)$	$U(K) \pm V(K)$
2	$\alpha u(x)$ Where α is a constant	$\alpha U(K)$
3	$\frac{du(x)}{dx}$	$(K + 1) U(K + 1)$
4	$\frac{d^n u(x)}{dx^n}$	$(K + 1)(K + 2) \dots (K + n) U(K + n)$
6	x^m	$\delta(K - m)$ where $\delta(K - m) = \begin{cases} 1, & \text{if } K = m \\ 0, & \text{if } K \neq m \end{cases}$
7	$e^{\alpha x}$	$\frac{\alpha^K}{K!}$
8	$\sin(\omega x + \alpha)$	$\frac{\omega^K}{K!} \sin\left(\frac{\pi K}{2} + \alpha\right)$
9	$\cos(\omega x + \alpha)$	$\frac{\omega^K}{K!} \cos\left(\frac{\pi K}{2} + \alpha\right)$

3. NUMERICAL APPLICATION

Example 1: [12]

$$y^{(viii)}(x) = e^{-x}y^2 \quad 0 < x < 1$$

Initial conditions are

$$y^i(0) = 1, \quad i = 0, 1, 2, \dots, 7$$

The exact solution is $y(x) = e^x$

Solution:

Applying 1-dim D.T.M. on initial condition we get

$$\begin{aligned} Y(0) = 1, \quad Y(1) = 1, \quad Y(2) = \frac{1}{2}, \quad Y(3) = \frac{1}{3!} \\ Y(4) = \frac{1}{4!}, \quad Y(5) = \frac{1}{5!}, \quad Y(6) = \frac{1}{6!}, \quad Y(7) = \frac{1}{7!} \end{aligned}$$

Using inverse 1-dim differential transform we have

$$\begin{aligned} y(x) &= \sum_{k=0}^{\infty} Y(k)x^k \\ y(x) &= Y(0) + Y(1)x + Y(2)x^2 + Y(3)x^3 + Y(4)x^4 + \dots \\ y(x) &= 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} + \frac{x^7}{7!} + \dots \end{aligned}$$

Table-2: Exact and approximate values of solution, and error of solution

x	Exact $y(x)$	1-Dim DTM $y^*(x)$	$ y(x) - y^*(x) $
0	1.0000000000000000	1.0000000000000000	0
0.1	1.105170918075648	1.105170918075397	$2.511324000 \times 10^{-13}$
0.2	1.221402758160170	1.221402758095238	$6.49318377 \times 10^{-11}$
0.3	1.349858807576003	1.349858805892857	$1.6831460670 \times 10^{-9}$
0.4	1.491824697641270	1.491824680634921	$1.70063498839 \times 10^{-8}$
0.5	1.648721270700128	1.648721168154762	$1.025453664205 \times 10^{-7}$
0.6	1.822118800390509	1.822118354285714	$4.461047944382 \times 10^{-7}$
0.7	2.013752707470477	2.013751158194445	$1.5492760319091 \times 10^{-6}$
0.8	2.225540928492468	2.225536365714286	$4.5627781819491 \times 10^{-6}$
0.9	2.459603111156950	2.459591262678571	$1.18484783784290 \times 10^{-5}$
1.0	2.718281828459046	2.718253968253968	$2.78602050771681 \times 10^{-5}$

$y^*(x)$ is the approximate solution of $y(x)$




Figure-1: Comparative analysis of exact and 1-dim DTM solution results

Example 2: [11]

$$y^{(viii)} + y^{(vii)} + 2y^{(vi)} + 2y^{(v)} + 2y^{(iv)} + 2y''' + 2y'' + y' + y = 14\cos x - 16\sin x - 4x\sin x$$

$$0 < x < 1$$

Boundary conditions are

$$y(0) = 0, y'(0) = -1, y''(0) = 0, y'''(0) = 7$$

$$y(1) = 0, y'(1) = 2\sin 1, y''(1) = 4\cos 1 + 2\sin 1, y'''(1) = 6\cos 1 - 6\sin 1 \tag{4}$$

The exact solution is $y(x) = (x^2 - 1)\sin x$

Solution:

Applying 1-dim D.T.M. on (3) we get

$$Y(0) = 0, Y(0) = -1, Y(2) = 0, Y(3) = \frac{7}{6}$$

Using inverse 1-dim D.T.M. We have

$$y(x) = \sum_{k=0}^{\infty} Y(k)x^k$$

$$y(x) = Y(0) + Y(1)x + Y(2)x^2 + \dots$$

Let $Y(4) = A, Y(5) = B, Y(6) = C, Y(7) = D$

$$y(x) = -x + \frac{7}{6}x^3 + Ax^4 + Bx^5 + Cx^6 + Dx^7 + \dots \tag{5}$$

As $y(1) = 0$ in (4) we get

$$A + B + C + D = -\frac{1}{6} \tag{6}$$

As $y'(1) = 2\sin 1$ in (4) we get

$$4A + 5B + 6C + 7D = -0.8170580304 \tag{7}$$

As $y''(1) = 4\cos 1 + 2\sin 1$ in (4) we get

$$12A + 20B + 30C + 42D = -3.155848807 \tag{8}$$

As $y'''(1) = 6\cos 1 - 6\sin 1$ in (4) we get

$$24A + 60B + 120C + 210D = -8.84867874 \tag{9}$$

From equations (6) to (9), we get the values

$$A = 0.00075045083330$$

$$B = -0.177827573600001$$

$$C = 0.003795158433337$$

$$D = 0.006615297666666$$

Putting the values of A, B, C and D in (5) we get

$$y(x) = -x + \frac{7}{6}x^3 + 0.00075045083330x^4 - 0.177827573600001x^5 + 0.003795158433337x^6$$

$$+ 0.006615297666666x^7 + \dots$$

Table-3: Exact and approximate values of solution, and error of solution

x	Exact $y(x)$	1-Dim DTM $y^*(x)$	$ y(x) - y^*(x) $
0	0	0	0
0.1	-0.098835082480360	-0.098835032107298	$5.0373062054910 \times 10^{-08}$
0.2	-0.190722557563259	-0.190722043202935	$5.1436032327823 \times 10^{-07}$
0.3	-0.268923388061819	-0.268921828916000	$1.55914581861927 \times 10^{-6}$
0.4	-0.327111407539266	-0.327108692673024	$2.71486624231487 \times 10^{-6}$
0.5	-0.359569153953152	-0.359565893801042	$3.26015211038477 \times 10^{-6}$
0.6	-0.361371182972823	-0.361368360786509	$2.82218631347808 \times 10^{-6}$
0.7	-0.328551020491222	-0.328549335580049	$1.68491117297220 \times 10^{-6}$
0.8	-0.258248192723828	-0.258247614837010	$5.77886818231570 \times 10^{-7}$
0.9	-0.148832112829222	-0.148832053983802	$5.8845419786960 \times 10^{-8}$
1.0	0	-0.000000000000001	$1.267220000000 \times 10^{-15}$

$y^*(x)$ is the approximate solution of $y(x)$




Figure-2: Comparative analysis of exact and 1-dim DTM solution results

Example 3: [11]

$$y^{(viii)}(x) + xy = -(48 + 15x + x^3)e^x \quad 0 < x < 1$$

Boundary conditions are

$$y(0) = 0, \quad y'(0) = 1, \quad y''(0) = 0, \quad y'''(0) = -3 \tag{10}$$

$$y(1) = 0, \quad y'(1) = -e, \quad y''(1) = -4e, \quad y'''(1) = -9e \tag{11}$$

The exact solution is $y(x) = x(1 - x)e^x$

Solution:

Applying 1-dim D.T.M. on (10)

$$Y(0) = 0, \quad Y(1) = 1, \quad Y(2) = 0, \quad Y(3) = -\frac{1}{2}$$

Using Inverse 1-dim Differential Transform we have

$$y(x) = Y(0) + Y(1)x + Y(2)x^2 + Y(3)x^3 + Y(4)x^4 + \dots$$

$$y(x) = x - \frac{x^3}{2} + Y(4)x^4 + Y(5)x^5 + Y(6)x^6 + Y(7)x^7 + \dots$$

Let $Y(4) = A, \quad Y(5) = B, \quad Y(6) = C, \quad Y(7) = D$

$$y(x) = x - \frac{x^3}{2} + Ax^4 + Bx^5 + Cx^6 + Dx^7 + \dots \tag{12}$$

As $y(1) = 0$ gives

$$A + B + C + D = -\frac{1}{2} \tag{13}$$

As $y'(1) = -e$ gives

$$4A + 5B + 6C + 7D = -e + \frac{1}{2} \tag{14}$$

As $y''(1) = -4e$ gives
 $12A + 20B + 30C + 42D = -4e + 3$ (15)

As $y'''(1) = -9e$ gives
 $24A + 60B + 120C + 210D = -9e + 3$ (16)

From equation (13), (14), (15) and (16) we get

$$\begin{aligned} A &= -0.331168115016212 \\ B &= -0.133368341115217 \\ C &= -0.021477144261922 \\ D &= -0.013986399606659 \end{aligned}$$

Putting A, B, C and D into (12) we get

$$y(x) = x - \frac{1}{2}x^3 - 0.331168115016212x^4 - 0.133368341115217x^5 - 0.021477144261922x^6 - 0.013986399606659x^7 + \dots$$

Table-4: Exact and approximate values of solution, and error of solution

x	Exact y(x)	1-Dim DTM y*(x)	y(x) - y*(x)
0	0	0	0
0.1	0.099465382626808	0.099465526629303	1.4400249472024x10 ⁻⁷
0.2	0.195424441305627	0.195425899583669	1.45827804226562x10 ⁻⁶
0.3	0.283470349590961	0.283474737535698	4.38794473711646x10 ⁻⁶
0.4	0.358037927433905	0.358045518742553	7.59130864785851x10 ⁻⁶
0.5	0.412180317675032	0.412189383025617	9.06535058464719x10 ⁻⁶
0.6	0.437308512093722	0.437316322770066	7.81067634431088x10 ⁻⁶
0.7	0.422888068568800	0.422892713798975	4.64523017534857x10 ⁻⁶
0.8	0.356086548558795	0.356088136976538	1.58841774272611x10 ⁻⁶
0.9	0.221364280004125	0.221364441395012	1.61390886932460x10 ⁻⁷
1.0	0	-0.000000000000010	1.0023230000000x10 ⁻¹⁴

$y^*(x)$ is the approximate solution of $y(x)$




Figure-3: Comparative analysis of exact and 1-dim DTM solution results

Example 4: [12]

$$y^{(viii)}(x) + 8e^x - y(x) = 0 \quad 0 < x < 1$$

Boundary conditions are

$$y(0) = 0, y'(0) = 0, y''(0) = -1, y'''(0) = -2, y^{(iv)}(0) = -3, y^{(v)}(0) = -4 \quad (17)$$

$$y'(1) = -e, y''(1) = -2e \quad (18)$$

The exact solution is $y(x) = (1 - x)e^x$

Solution:

Applying 1-dim D.T.M. on (17) we get

$$Y(0) = 1, Y(1) = 0, Y(2) = -\frac{1}{2}, Y(3) = -\frac{1}{3}, Y(4) = -\frac{1}{8}, Y(5) = -\frac{1}{30}$$

Using Inverse 1-dim Differential Transform we have

$$y(x) = Y(0) + Y(1)x + Y(2)x^2 + Y(3)x^3 + Y(4)x^4 + Y(5)x^5 + Y(6)x^6 + Y(7)x^7 + \dots$$

$$y(x) = 1 - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{8} - \frac{x^5}{30} + Y(6)x^6 + Y(7)x^7 + \dots$$

Let $Y(6) = A, Y(7) = B$

$$y(x) = 1 - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{8} - \frac{x^5}{30} + Ax^6 + Bx^7 + \dots \tag{19}$$

As $y'(1) = -e$ gives

$$6A + 7B = -e + \frac{8}{3} \tag{20}$$

As $y''(1) = -2e$ Gives

$$30A + 42B = -2e + \frac{31}{6} \tag{21}$$

From equation (20) and (21) we get

$$A = -0.006632330083808$$

$$B = -0.001688740184218$$

Putting A, and B into (19) we get

$$y(x) = 1 - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{8} - \frac{x^5}{30} - 0.006632330083808x^6 - 0.001688740184218x^7 + \dots$$

Table-5: Exact and approximate values of solution, and error of solution

x	Exact $y(x)$	1-Dim DTM $y^*(x)$	$ y(x) - y^*(x) $
0	1.0000000000000000	1.0000000000000000	0
0.1	0.994653826268083	0.994653826532129	$2.6404622930 \times 10^{-10}$
0.2	0.977122206528136	0.97712220581667	$1.40535308946 \times 10^{-8}$
0.3	0.944901165303202	0.944901295703891	$1.304006884695 \times 10^{-7}$
0.4	0.895094818584762	0.895095400477392	$5.818926300805 \times 10^{-7}$
0.5	0.824360635350064	0.824362343226418	$1.7078763537848 \times 10^{-6}$
0.6	0.728847520156204	0.728851288090389	$3.7679341854391 \times 10^{-6}$
0.7	0.604125812241143	0.604132471315550	$6.6590744072226 \times 10^{-6}$
0.8	0.445108185698493	0.445117886643695	$9.7009452019181 \times 10^{-6}$
0.9	0.245960311115695	0.245972088672914	$1.17775572190670 \times 10^{-5}$
1.0	0	0.000012263065307	$1.22630653073522 \times 10^{-5}$

$y^*(x)$ is the approximate solution of $y(x)$




Figure-4: Comparative analysis of exact and 1-dim DTM solution results

4. CONCLUSION

In this work, one dimensional Differential Transformation Method has been applied to obtain the numerical solution of linear and nonlinear eighth order boundary value problems. The present method has been applied in a direct way without using linearization, discretization, or perturbation. The numerical results obtained by this method are in good agreement with the exact solutions available in the literature. This technique is fast converging to the exact solution and requires much less computational work than other methods. The objective of this paper is to present a simple method to solve an eighth order boundary value problem and its easiness for implementation.

REFERENCES

1. A.R. Davies, A. Karageorghis, T.N. Phillips, Spectral Galerkin methods for the primary two-point boundary-value problem in modelling viscoelastic flow, *International Journal of Numerical Methods*, 26 (1988) 647-662.
2. A. Karageorghis, T.N. Phillips, A.R Davies, Spectral collocation methods for the primary two-point boundary-value problem in modelling viscoelastic flows, *International Journal of Numerical Methods in Engineering*, 26 (1988) 805-813.
3. S Chandrasekhar, *Hydrodynamic and Hydromagnetic Stability*, Clarendon Press, Oxford (Reprinted: Dover Books, New York, 1981) 1961.
4. A. Boutayeb, E.H. Twizell, Finite-difference methods for the solution of eighth-order boundary-value problems, *Applied Mathematics Letters*, 48 (1993) 63-75.
5. E.H. Twizell, S.S. Siddiqi, Spline solution of linear eighth-order boundary value problems, *Comput. Methods Appl. Mech. Eng.*, 131 (1996) 309-325.
6. M. Inc, D.J. Evans, An efficient approach to approximate solutions of eighth order boundary-value problems, *Int. J. Comput. Math.*, 81 (2004) 685-692.
7. Shahid, S. Siddiqi, and Ghazala Akram., Nonic spline solutions of eighth order boundary value problems, *J. Comput. Appl. Math.*, 182 (2006) 829-845.
8. Shahid, S. Siddiqi, and Ghazala Akram, Solutions of eighth order boundary value problems using non-polynomial spline technique, *International Journal of Computer Mathematics*, 84(3) (2007) 347-368.
9. G.R. Liu, T.Y. Wu, Differential quadrature solutions of eighth order boundary value differential equations, *J. Comput. Appl. Math.*, 145 (1973) 223-235.
10. Mladen Meštrović, The modified decomposition method for eighth-order boundary value problems, *Appl. Math. And Comput.* 188 (2007) 1437-1444.
11. K.N.S. Kasi Viswanadham and Y. Showri Raju, Quintic B-spline Collocation Method for eighth-order boundary value problems, *Advances in computational math. and its applications*, vol. 1, No1 (2012) pp. 47-52.
12. Sirajul Haq, M. Idrees, S. Islam, Application of optimal Homotopy asymptotic method to eighth-order initial and boundary value problems, *Int. J. of Appl. Math. And Comput.* Volume 2(4), pp 73-80, 2010.
13. Zhou J. K., 1986. *Differential Transformation and Its Applications for Electrical Circuits*, Huazhong University Press, Wuhan, China (in Chinese).
14. Montri Thongmoon, Sasitorn Pusjuso, The numerical solutions of differential transform method and the Laplace transform method for a system of differential equations, *Nonlinear Analysis: Hybrid Systems* 4 (2010) 425-431.
15. F. Ayas, Solutions of the system of differential equations by differential transform method, *Appl. Math. Comput.* 147 (2004) 547-567.
16. I.H. Abdel-Halim Hassan, Differential transformation technique for solving higher-order initial value problems, *Appl. Math. Comput.* 154 (2004) 299-311.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]