

PRE-rg-CLOSED FUNCTIONS IN TOPOLOGY

¹GOVINDAPPA NAVALAGI AND ^{*2}SUJATA MOOKANAGOUDAR

¹Department of Mathematics, KIT Tiptur-572202, Karnataka, India.

^{*2}Department of Mathematics,
Government First Grade College, Haliyal-581329, Karnataka, India.

(Received On: 01-07-18; Revised & Accepted On: 29-07-18)

ABSTRACT

Levine in 1970, introduced the concept of generalized closed (*g*-closed) sets in topological space and a class of topological spaces called $T_{1/2}$ -spaces. Palaniappan *et al* in 1993, introduced the notions of regular generalized (in brief, *rg*-) closed sets, *rg*-continuous functions and *rg*-irresoluteness and in 1997, Arokia Rani *et al.* introduced and studied the concepts of, *rg*-openness and *rg*-closedness in topology. In 2013, Navalagi *et al* have defined and studied the concepts of pre-*rg* - openness and other allied *rg*-open functions. The purpose of this paper is to investigate the concept of pre-regular generalized (in brief, pre-*rg*-) closed functions and some allied *rg*-closed functions in topological spaces. Also, we study some more properties of *rg*-closed functions.

Mathematics Subject Classification (2010): 54A05, 54B05, 54C08.

Key words and Phrases: regular open sets, regular closed sets, *g*-closed sets and *rg*-closed sets, *rg*-openness, *rg*-closedness, *rg*-continuity, *rg*-irresoluteness, pre-*rg*-openness.

1. INTRODUCTION

Levine [6] generalized the closed set to generalized closed set (in brief, *g*-closed set) in topology for the first time. Since then it is noticed that some of the weaker forms of closed sets have been generalized in the literature. Thereafter, several authors have obtained many interesting results on these generalized closed sets [cf. 2]. Palaniappan and Rao [12] introduced and studied the notions of regular generalized (in brief, *rg*-) closed sets, *rg*-continuous functions, *rg*-irresoluteness. In 1997, Arokia Rani *et al.* [1] have studied the further properties of *rg*-continuity, *rg*-irresoluteness, *rg*-open functions, *rg*-closed functions, *rg*-closure operator, strongly *rg*-continuous functions, perfectly *rg*-continuous functions, T_{rg} -spaces, $T^*_{1/2}$ -spaces and *rg*-normal spaces in topology. The purpose of this paper is to investigate some more properties of regular generalized closed (i.e. in brief, *rg*-closed) functions and other allied *rg*-closed functions in topology and introduce and study pre-*rg*-closed functions.

2. PRELIMINARIES

Throughout this paper (X, τ) , (Y, σ) , (Z, γ) (or simply X , Y and Z) always means topological spaces on which no separation axioms are assumed unless explicitly stated. Let A be a subset of space X . We denote the closure of A and the interior of A by $Cl(A)$ and $Int(A)$ respectively. A subset A of a space X is called regular open (resp. regular closed) if $A = Int Cl(A)$ (resp. $A = Cl Int(A)$).

Definition 2.1 [7]: A subset A of a space X is said to be a pre-open if $A \subset Int(Cl(A))$.

The family of all preopen sets in a space X is denoted by $PO(X)$. The complement of a preopen set of a space X is called preclosed [4].

Definition 2.2 [8]: The union of all preopen sets contained in A is called the preinterior of A and is denoted by $pInt(A)$

Corresponding Author: ^{*2}Sujata Mookanagoudar, ^{*2}Department of Mathematics,
Government First Grade College, Haliyal-581329, Karnataka, India.

Definition 2.3[4]: The intersection of all preclosed sets containing A is called the preclosure of A and is denoted by $pCl(A)$.

Definition 2.4 [5]: A subset A of a space X is called a generalized closed set (g-closed) set if $Cl(A) \subset U$ whenever $A \subset U$ and U is open.

Clearly, every closed set is a g-closed set. The complement of a g-closed set in X is called g-open set. Clearly, every open set is a g-open set.

Definition 2.5 [11]: A subset A of a space X is called a generalized preclosed set (in brief ,gp-closed) set if $pCl(A) \subset U$ whenever $A \subset U$ and U is open. The complement of a gp-closed set in X is called gp-open set.

It is obvious that every preopen set is gp-open set.

Definition 2.6 [12]: A subset A of a space X is called a regular generalized closed (rg-closed) set if $Cl(A) \subset U$ whenever $A \subset U$ and U is regular open. The complement of a rg-closed set of a space is called rg-open. The family of all rg-open sets of a space X is denoted by $RGO(X)$.

Definition 2.7: A function $f: X \rightarrow Y$ is said to be :

- i) preopen [8] if the image of each open set U of X, $f(U)$ is preopen in Y.
- ii) preclosed[4] if the image of each closed set F of X, $f(F)$ is preclosed in Y.
- iii) preirresolute [13] if the inverse image of each preopen set of Y is preopen in X.
- iv) generalized preclosed(gp-closed[10]) if the image of each closed set of X is gp-closed in Y.

Definition 2.8: A function $f: X \rightarrow Y$ is called

- i) rg-open [1] if image of each open set of X is rg-open in Y.
- ii) rg-closed [1] if image of each closed set of X is rg-closed in Y.
- iii) g-closed [6] if the image of each closed set of X is g-closed in Y.
- iv) almost closed [14] if the image of each regular closed set of X is closed in Y.

Definition 2.9: A function $f: X \rightarrow Y$ is said to be :

- i) rg-continuous [12] if the inverse image of each closed set of Y is rg-closed in X.
- ii) rg-irresolute [12] if $f^{-1}(V)$ is rg-open in X for every rg-open set V of Y.
- iii) perfectly rg-continuous [1] if the inverse image of each rg-open set in Y is both open and closed in X.

Definition 2.10[9]: A function $f: X \rightarrow Y$ is said to be M-preclosed if the image of each preclosed set of X is preclosed in Y.

3. Pre-rg – closed functions

In view of g-closedness and rg-closedness definitions, we can observe that every g-closed function is rg-closed function .
We, characterize the rg-closedness in the following.

Theorem 3.1: Let $f: X \rightarrow Y$ be a function. Then the following are equivalent:

- i) f is rg-closed.
- ii) The image of each closed set in X is rg-closed in Y.
- iii) The image of each open set in X is rg-open in Y.

Proof:

(i) \Leftrightarrow (ii): Follows from Definition.

(i) \Leftrightarrow (iii): Let F be any open set in X. Then $X - F$ is closed in X. Since f is rg-closed, $f(X - F)$ is rg-closed in Y. But $f(X - F) = Y - f(F)$ is rg-closed in X. Therefore f(F) is rg-open in Y.

We, define the following.

Definition 3.2: A function $f: X \rightarrow Y$ is called strongly rg-closed if the image of each rg-closed set of X is closed in Y.

Theorem 3.3: If $f: X \rightarrow Y$ is rg-closed and $g: Y \rightarrow Z$ is strongly rg-closed, then the composition $g \circ f: X \rightarrow Z$ is closed .

Proof: Obvious.

Theorem 3.4: If $f: X \rightarrow Y$ is closed and $g: Y \rightarrow Z$ is rg-closed, then the composition $g \circ f: X \rightarrow Z$ is rg-closed.

Easy proof is omitted.

We, recall the following.

Definition 3.5[1]: A topological space (X, τ) is said to be $T^*_{1/2}$ iff every rg-closed set is closed.

Theorem 3.6: Let X and Z be any topological spaces and Y be a $T^*_{1/2}$ space. Then the composition $g \circ f: X \rightarrow Z$ of two rg-closed functions $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ is also rg-closed.

Proof: Easy.

We now define the following.

Definition 3.7: A function $f: X \rightarrow Y$ is said to be pre-rg-closed if the image of each regular closed set of X is rg-closed in Y .

Clearly, every rg-closed function is pre-rg-closed.

Theorem 3.8: If a function $f: X \rightarrow Y$ is strongly rg-closed then it is an almost-closed.

Proof: Let F be any regular closed set in X . Then F is rg-closed in X since every regular open set is rg-open. Again, as f is strongly rg-closed and F is rg-closed in X , $f(F)$ is an closed in Y . So, f is almost closed.

Theorem 3.9: A function $f: X \rightarrow Y$ is pre rg-closed iff the image of every regular open set in X is rg-open in Y .

Proof: Easy.

We, define the following.

Definition 3.10: A function $f: X \rightarrow Y$ is said to be always rg-closed, if the image of each rg-closed set of X is rg-closed in Y .

Definition 3.11: A function $f: X \rightarrow Y$ is said to be completely rg-closed, if the image of each rg-closed set of X is regular closed in Y .

Theorem 3.12: If $f: X \rightarrow Y$ is always -rg-closed and $g: Y \rightarrow Z$ is completely rg-closed, then $g \circ f: X \rightarrow Z$ is completely rg-closed.

Proof: Obvious.

Theorem 3.13: If $f: X \rightarrow Y$ is completely rg-closed and $g: Y \rightarrow Z$ is pre-rg-closed, then $g \circ f: X \rightarrow Z$ is always rg-closed.

Proof: Let F be rg-closed set in X . Since f is completely rg-open, $f(F)$ is regular closed in Y . Hence, $g(f(F)) = g \circ f(F)$ is rg-closed in Z as g is pre-rg-closed function. Therefore, $g \circ f$ is always rg-closed.

We, define the following.

Definition 3.14: A function $f: X \rightarrow Y$ is said to be prg-closed if image of each preclosed set of X is rg-closed in Y .

The following is a characterization of prg-open functions.

Theorem 3.15: Let $f: X \rightarrow Y$ be a function. Then the following are equivalent.

- i) f is prg-closed.
- ii) The image of each preclosed set in X is rg-closed in Y .
- iii) The image of each preopen set in X is rg-open in Y .

Proof: Follows from the definition.

Theorem 3.16: If $f: X \rightarrow Y$ is M-preclosed and $g: Y \rightarrow Z$ is prg-closed then the composition $g \circ f: X \rightarrow Z$ is prg-closed

Proof: Let F be preclosed set in X . Since f is M-preclosed, $f(F)$ is preclosed in Y . Therefore $g(f(F)) = g \circ f(F)$ is rg-closed in Z since g is prg-closed. Hence, the composition function $g \circ f$ is prg-closed.

Theorem 3.17: If $f: X \rightarrow Y$ is preclosed and $g: Y \rightarrow Z$ is prg-closed, then the composition $g \circ f: X \rightarrow Z$ is rg-closed.

Proof: Obvious.

We, define the following.

Definition 3.18: A function $f: X \rightarrow Y$ is called strongly p-closed if the image of each preclosed set of X is closed in Y .

Clearly, every strongly p-closed function is M-preclosed.

Theorem 3.19: If $f: X \rightarrow Y$ is prg-closed function and $g: Y \rightarrow Z$ is strongly rg-closed then the composition $g \circ f: X \rightarrow Z$ is strongly p-closed .

Proof: Let F be preclosed set in X . Since f is prg-closed, $f(F)$ is rg-closed in Y . Therefore $g(f(F)) = g \circ f(F)$ is closed in Z . Hence, $g \circ f$ is strongly p-closed function.

Theorem 3.20: If $f: X \rightarrow Y$ is strongly p-closed and $g: Y \rightarrow Z$ is rg-closed then the composition is $g \circ f: X \rightarrow Z$ is prg-closed .

Proof: Obvious.

Theorem 3.21: If $f: X \rightarrow Y$ is rg-closed and $g: Y \rightarrow Z$ is always rg-closed then the composition $g \circ f: X \rightarrow Z$ is rg-closed.

Proof: Let F be a closed set in X . Since f is rg-closed, $f(F)$ is rg-closed. Therefore, $g(f(F)) = g \circ f(F)$ is rg-closed set in Z since g is always rg-closed function. Hence, $g \circ f$ is rg-closed .

Now, we define the following.

Definition 3.22: A function $f: X \rightarrow Y$ is said to be contra strongly rg-closed if the image of each rg-closed set of X is open in Y .

It is easy to see that a function $f: X \rightarrow Y$ is contra strongly rg-closed iff image of each rg-open set of X is closed in Y .

Theorem 3.23: If $f: X \rightarrow Y$ is contra-strongly rg-closed and $g: Y \rightarrow Z$ is contra-open then the composition $g \circ f: X \rightarrow Z$ is contra-strongly rg-closed .

Proof: Obvious.

Definition 3.24: A function $f: X \rightarrow Y$ is said to be contra rg-closed if the image of each closed set of X is rg-open in Y . Routine proof of the following Theorem is omitted.

Theorem 3.25: Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be two functions. Then, the following statements are valid:

- (i) If f is preclosed and g is prg-closed, then $g \circ f$ is rg-closed.
- (ii) If f is M-preclosed and g is strongly p-closed, then $g \circ f$ is strongly p-closed .
- (iii) If f is prg-closed and g is always rg-closed, then $g \circ f$ is prg-closed.
- (iv) If f is pre-rg-closed and g is strongly rg-closed, then $g \circ f$ is almost - closed.
- (v) If $g \circ f$ is always rg-closed and f is rg-irresolute surjection, then g is always rg-closed
- (vi) If $g \circ f$ is pre-rg-closed and f is completely rg-continuous, then g is always rg-closed .
- (vii) If $g \circ f$ is strongly rg-closed and f is rg-continuous surjection, then g is closed .
- (viii) If $g \circ f$ is prg-closed and f is precontinuous surjective, then g is rg-closed.
- (ix) If $g \circ f$ is strongly rg-closed and f is rg-continuous surjection, then g is closed .
- (x) If $g \circ f$ is M-preclosed and g is prg-continuous injection, then f is prg-closed .
- (xi) If $g \circ f$ is always rg-closed and g is rg-irresolute injection, then f is always rg-closed.
- (xii) If $g \circ f$ is pre-rg-closed and g is strongly rg-continuous injection, then f is almost oprn.
- (xiii) If $g \circ f$ is contra rg-closed and g is rg-irresolute injection, then f is rg-closed.
- (xiv) If $g \circ f$ is prg-closed and g is rg-irresolute injection, then f is prg-closed.
- (xv) If $g \circ f$ is rg-closed and g is strongly rg-continuous injective, then f is open.

REFERENCES

1. Arokia Rani and K.Balachandran, On regular generalized continuous maps in topological spaces, Kyungpook Math. J., 37(1997), 305-314.
2. K. Balachandran, P. Sundaram and H. Maki, On generalized continuous maps in topological spaces, Memoris Fac. Sci. Kochi. Univ., 12 (1991), 5-13.
3. J. Dontchev, Contra-continuous functions and strongly S-closed spaces, Internat. J. Math. Math. Sci., 19 (1996), 303-310.
4. S.N. El-Deeb, I. A. Hasanein, A. S. Mashhour and T.Noiri, On p-regular spaces, Bull. Math. Soc.Sci.Math. R.S.Roumanie, 27(1983).
5. N.Levine, Generalized closed sets in topology, Rend. Circ. Math Palermo, 19(1970), 89-96.
6. S.R.Malghan, Generalized closed maps, J.Karnatak Univ. Sci., 27(1982), 82-88.
7. A. S. Mashhour, M. E. Abd El-Monsef and S. N. El-Deeb, On pre-continuous and weak pre-continuous mappings, Proc. Math.Phys.Soc. Egypt, 53(1982), 47-53.
8. A. S. Mashhour, M. E. Abdel Monsef and I. A. Hasanein, On pre-topological spaces, Bull. Math. Soc. Sci. Math. R.S. Roumanie, 28(76) (1984), 34-45.
9. A. S. Mashhour, M. E. Abdel Monsef and I. A. Hasanein and T.Noiri, Delta J. Sci., 8 (1984), 30-46.
10. G.Navalagi, M.Savita, Supriya Kadam and Sujatha S.P , Pre-rg-open functions in topology, American Journal of Mathematics and Science, Vol.2,No.1, (Jan.2013), 201-207.
11. T. Noiri, H.Maki And J. Umehara, Generalized preclosed functions. Mem. Fac. Sci. Kochi Univ. Ser. A (Math.), Vol.19 (1998), 13-20.
12. N.Palaniappan and K.C.Rao, Regular generalized closed sets, Kyungpook Math. J., 33(1993), 211-219.
13. I.L.Reilly and M.K.Vamanmurthy, On α -continuity in topological spaces, Acta. Math. Hungar., 45(1-2), (1985), 27-32.
14. M.K.Singal and Asha Rani Singal, Almost continuous mappings, Yokohama Math. J., 16 (1968), 63-73.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]