

ENTIRE RATE SEQUENCE SPACE OF INTERVAL NUMBERS

Dr. D. CHRISTIAJEB A KUMARI

**Assistant professor, Department of Mathematics,
 Pope's College, Sawyerpuram-628251, Thoothukudi District,
 Affiliated to Manonmaniam Sundaranar University,
 Tirunelveli-627012, Tamilnadu, India.**

(Received On: 17-04-18; Revised & Accepted On: 13-08-18)

ABSTRACT

In this paper we introduced the new concept of interval valued sequence space $\Gamma_{\pi}(IR)$ where (π_k) is a sequence of positive numbers. We present the different properties like completeness, solidness, AB property etc.

2000 Mathematics subject classification: 46A 45.

Keywords and Phrases: Banach space, AB space, AK property, interval numbers.

1. INTRODUCTION

The power of interval arithmetic lies in its implementation on computers. In particular, outwardly rounded interval arithmetic allows rigorous enclosures for the ranges of operations and functions. This makes a qualitative difference in scientific computations, since the results are now intervals in which the exact result must lie. It also enables the use of computations for proving automated theorem.

Interval arithmetic is a tool in numerical computing where the rules for the arithmetic of intervals are explicitly stated. It was first suggested by P.S.Dwyer [10] in 1951. Development of interval arithmetic as a formal system and evidence of its value as a computational device was provided by R.E.Moore [34], [35] in 1959 and 1962.

A set consisting of a closed interval of real numbers x such that $a \leq x \leq b$ is called an interval number. A real interval can also be considered as a set. We denote the set of all real valued closed intervals by $I\mathfrak{R}$. Any element of $I\mathfrak{R}$ may be called closed interval and denoted by \hat{x} . That is, $\hat{x} = [x_l, x_r] = \{x \in \mathfrak{R} : x_l \leq x \leq x_r\}$. An interval number \hat{x} is a closed subset of real numbers. Let x_l and x_r be respectively referred to as the infimum (lower bound) and supremum (upper bound) of the interval number \hat{x} . If $\hat{x} = [0,0]$, then \hat{x} is said to be a zero interval. It is denoted by $\hat{0}$.

For $\bar{x}_1, \bar{x}_2 \in I\mathfrak{R}$, we define $\bar{x}_1 = \bar{x}_2$ if and only if $x_{1l} = x_{2l}$ and $x_{1r} = x_{2r}$

$$\bar{x}_1 + \bar{x}_2 = \{x \in \mathfrak{R} : x_{1l} + x_{2l} \leq x \leq x_{1r} + x_{2r}\}$$

$$\bar{x}_1 \times \bar{x}_2 = \{x \in \mathfrak{R} : \min(x_{1l}x_{2l}, x_{1l}x_{2r}, x_{1r}x_{2l}, x_{1r}x_{2r}) \leq x \leq \max(x_{1l}x_{2l}, x_{1l}x_{2r}, x_{1r}x_{2l}, x_{1r}x_{2r})\}$$

The set of all interval numbers $I\mathfrak{R}$ is a complete metric space defined by

$$d(\bar{x}_1, \bar{x}_2) = \max\{|\bar{x}_{1l} - \bar{x}_{2l}|, |\bar{x}_{1r} - \bar{x}_{2r}|\}$$

In the special case $\bar{x}_1 = [a, a]$ and $\bar{x}_2 = [b, b]$, we obtain usual metric of \mathfrak{R} .

Corresponding Author: Dr. D. ChristiaJebakumari

Assistant professor, Department of Mathematics,

Pope's College, Sawyerpuram-628251, Thoothukudi District,

Affiliated to Manonmaniam Sundaranar University, Tirunelveli-627012, Tamilnadu, India.

Let us define transformation $f: \mathbb{N} \rightarrow \mathfrak{R}$, $k \rightarrow f(k) = \bar{x}_k$, then $\bar{x} = (\bar{x}_k)$ is called sequence of interval numbers. \bar{x}_k is called k^{th} term of sequence $\bar{x} = (\bar{x}_k)$, ω^i denotes the set of all interval numbers with real terms and the algebraic properties of ω^i are in[7].

A sequence $\bar{x} = (\bar{x}_k)$ of interval numbers is said to be convergent to the interval number \bar{x}_0 if for each $\varepsilon > 0$ there exists a positive integer k_0 such that $d(\bar{x}_k, \bar{x}_0) < \varepsilon$ for all $k \geq k_0$ and we denote it by $\lim_k \bar{x}_k = \bar{x}_0$. Equivalently $\lim_k \bar{x}_k = \bar{x}_0$ iff $\lim_k x_{kl} = x_{0l}$ and $\lim_k x_{kr} = x_{0r}$.

2. MAIN RESULTS

The entire sequence space of symmetric interval numbers is denoted by $\Gamma_\pi(IR)$ where (π_k) is a sequence of positive numbers and the set of functional of interval numbers is denoted by $\Lambda_\pi(IR)$.

$$\Gamma_\pi(IR) = \left\{ \tilde{x} = \left(\frac{\tilde{x}_k}{\pi_k} \right) \in w(IR) : \lim_k D \left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) = \tilde{0} \right\}$$

$$\Lambda_\pi(IR) = \left\{ \tilde{x} = \left(\frac{\tilde{x}_k}{\pi_k} \right) \in w(IR) : \sup_k D \left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) < \infty \right\}$$

where $D \left(\frac{\tilde{x}_k}{\pi_k}, \frac{\tilde{y}_k}{\pi_k} \right) = \max \left\{ \left| \frac{x_k - y_k}{\pi_k} \right|^{1/k}, \left| \frac{\bar{x}_k - \bar{y}_k}{\pi_k} \right|^{1/k} \right\}$

The metric \tilde{d} is defined by

$$\tilde{d} \left(\frac{\tilde{x}_k}{\pi_k}, \frac{\tilde{y}_k}{\pi_k} \right) = \sup_k \max \left\{ \left| \frac{x_k - y_k}{\pi_k} \right|^{1/k}, \left| \frac{\bar{x}_k - \bar{y}_k}{\pi_k} \right|^{1/k} \right\} = \sup_k D \left(\frac{\tilde{x}_k}{\pi_k}, \frac{\tilde{y}_k}{\pi_k} \right) \tag{2.1}$$

which satisfies the metric space axioms .

Theorem 2.1: The sequence space of interval numbers $\Gamma_\pi(IR)$ is a complete metric space with respect to the metric defined by (2.1).

Proof: Let $\left(\frac{\tilde{x}^{(n)}}{\pi^{(n)}} \right)$ be a interval number fundamental sequence in $\Gamma_\pi(IR)$. Then for a given $\varepsilon > 0$ there exists a positive integer n_0 such that

$$\tilde{d} \left(\frac{\tilde{x}_k^{(n)}}{\pi_k^{(n)}}, \frac{\tilde{x}_k^{(m)}}{\pi_k^{(m)}} \right) = \sup_k D \left(\frac{\tilde{x}_k^{(n)}}{\pi_k^{(n)}}, \frac{\tilde{x}_k^{(m)}}{\pi_k^{(m)}} \right) < \varepsilon \text{ for all } n, m \geq n_0$$

For each k , we have $D \left(\frac{\tilde{x}_k^{(n)}}{\pi_k^{(n)}}, \frac{\tilde{x}_k^{(m)}}{\pi_k^{(m)}} \right) < \varepsilon$ for all $n, m \geq n_0$

$$\max \left\{ \left| \frac{x_k^{(n)} - x_k^{(m)}}{\pi_k} \right|^{1/k}, \left| \frac{\bar{x}_k^{(n)} - \bar{x}_k^{(m)}}{\pi_k} \right|^{1/k} \right\} < \varepsilon \text{ for all } n, m \geq n_0$$

$$\left| \frac{x_k^{(n)} - x_k^{(m)}}{\pi_k} \right|^{1/k} < \varepsilon^k \text{ and } \left| \frac{\bar{x}_k^{(n)} - \bar{x}_k^{(m)}}{\pi_k} \right|^{1/k} < \varepsilon^k \text{ for all } n, m \geq n_0$$

This leads to the fact $\frac{\tilde{x}_k^{(n)}}{\pi_k^{(n)}}$ is a interval number fundamental sequence in IR . Since IR is a complete metric space,

$$\frac{\tilde{x}_k^{(n)}}{\pi_k^{(n)}} \text{ is convergent } \lim_n \frac{\tilde{x}_k^{(n)}}{\pi_k^{(n)}} = \frac{\tilde{x}_k}{\pi_k} \text{ for each } k \in \mathbb{N}.$$

Hence $\sup_k D\left(\frac{\tilde{x}_k^{(n)}}{\pi_k^{(n)}}, \frac{\tilde{x}_k}{\pi_k}\right) < \varepsilon$

So $\frac{\tilde{x}_k^{(n)}}{\pi_k^{(n)}} \rightarrow \frac{\tilde{x}_k}{\pi_k}$ as $n \rightarrow \infty$ in $\Gamma_\pi(IR)$

We have to show that $\tilde{x}_k = \left(\frac{\tilde{x}_k}{\pi_k}\right) \in \Gamma_\pi(IR)$

Since $\tilde{x}_k \in \Gamma_\pi(IR)$, we have $\tilde{d}\left(\frac{\tilde{x}_k^{(n)}}{\pi_k^{(n)}}, \tilde{0}\right) < \varepsilon$

Consider

$$\begin{aligned} \tilde{d}\left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0}\right) &= \sup_k D\left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0}\right) \\ &\leq \sup_k D\left(\frac{\tilde{x}_k^{(n)}}{\pi_k^{(n)}}, \frac{\tilde{x}_k}{\pi_k}\right) + \sup_k D\left(\frac{\tilde{x}_k^{(n)}}{\pi_k^{(n)}}, \tilde{0}\right) \\ &< \varepsilon + \varepsilon \\ &= 2\varepsilon \end{aligned}$$

Hence $\left(\frac{\tilde{x}_k}{\pi_k}\right) \in \Gamma_\pi(IR)$. This completes the proof.

Theorem 2.2: A necessary and sufficient condition that $D\left(\sum \frac{\tilde{x}_k \tilde{y}_k}{\pi_k \pi_k}, \tilde{0}\right)$ should be convergent for every $\left(\frac{\tilde{x}_k}{\pi_k}\right)$ in

which $\lim_k \left(D\left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0}\right)\right) = 0$ is that $D\left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0}\right)$ should be bounded.

Proof: Suppose $D\left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0}\right)$ is bounded. Then there is an M so that $D\left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0}\right) \leq M$ for $k \geq 1$.

Since $D\left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0}\right) \rightarrow \tilde{0}$ as $k \rightarrow \infty$, there exists a positive integer k_0 so that

$$\begin{aligned} D\left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0}\right) &\leq \frac{1}{2M}, k \geq k_0 \\ [D\left(\frac{\tilde{x}_k \tilde{y}_k}{\pi_k \pi_k}, \tilde{0}\right)]^k &\leq [D\left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0}\right)]^k [D\left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0}\right)]^k \\ &< \left(\frac{1}{2M}\right)^k M^k = \frac{1}{2^k} \end{aligned}$$

Then $\sum \left[D\left(\frac{\tilde{x}_k \tilde{y}_k}{\pi_k \pi_k}, \tilde{0}\right)\right]^k$ converges

Conversely,

Suppose $D\left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0}\right)$ is not bounded. Then there is an increasing sequence $\{k_p\}$ of integers such that

$$D\left(\frac{\tilde{y}_{k_p}}{\pi_{k_p}}, \tilde{0}\right) \geq p, p = 1, 2, \dots$$

That is, $\left[D \left(\frac{\tilde{y}_{k_p}}{\pi_{k_p}}, \tilde{0} \right) \right]^{k_p} \geq p^{k_p}, p = 1, 2, \dots$

Take $\frac{\tilde{x}_k}{\pi_k} = \begin{cases} \left[0, \frac{1}{p^k} \right] & \text{if } k = k_p \\ [0, 0] & \text{if } k \neq k_p \end{cases}$

Then $\lim_k \left(D \left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) \right) = 0$

But $\left[D \left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) \right]^{k_p} \left[D \left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0} \right) \right]^{k_p} = \left[\left(\frac{1}{p^{k_p}} \right)^{1/k_p} \right]^{k_p} \left[\left(p^{k_p} \right)^{1/k_p} \right]^{k_p} = \frac{1}{p^{k_p}} \cdot p^{k_p} = 1 \text{ for } k = k_p$

So that $\sum \left[D \left(\frac{\tilde{x}_k}{\pi_k}, \frac{\tilde{y}_k}{\pi_k}, \tilde{0} \right) \right]^k$ does not converge.

Hence $D \left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0} \right)$ is bounded.

Theorem 2.3: The sequence spaces of interval numbers $\Gamma_\pi(IR)$ and $\Lambda_\pi(IR)$ are solid.

Proof: Consider, $\Gamma_\pi(IR)$. Now let $\tilde{d} \left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0} \right) \leq \tilde{d} \left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right)$ for all $k \in N$ and for some $\tilde{x} \in \Gamma_\pi(IR)$.

Then, $\max \left\{ \left| \underline{y}_k \right|^{1/k}, \left| \bar{y}_k \right|^{1/k} \right\} \leq \max \left\{ \left| \underline{x}_k \right|^{1/k}, \left| \bar{x}_k \right|^{1/k} \right\}, \underline{y}_k \leq \underline{x}_k \text{ and } \bar{y}_k \leq \bar{x}_k$

$$D \left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0} \right) \leq D \left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right)$$

$$\lim_{k \rightarrow \infty} D \left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0} \right) \leq \lim_{k \rightarrow \infty} D \left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) = 0$$

It is clear that $\tilde{y}_k \in \Gamma_\pi(IR)$

Therefore $\Gamma_\pi(IR)$ is solid.

Similarly, it can be proved that $\Lambda_\pi(IR)$ is solid.

Theorem 2.4: The sequence of interval numbers $(\tilde{e}_1, \tilde{e}_2, \dots, \tilde{e}_k, \dots)$ is schauder base for $\Gamma_\pi(IR)$,

where $\tilde{e}_k = \{ \tilde{0}, \tilde{0}, \dots, [1, 1], \tilde{0}, \dots \}$.

Proof: Let $\tilde{x}_k = \left(\frac{\tilde{x}_k}{\pi_k} \right) \in \Gamma_\pi(IR)$. Therefore for every $\varepsilon > 0$ there exists a positive integer $n \in N$ such that $k \geq n$,

$$\tilde{d} \left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) = \sup_k D \left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) < \varepsilon$$

It is enough to show that $\lim_{k \rightarrow \infty} \tilde{d} \left[\frac{\tilde{x}_k}{\pi_k} - \sum \left(\tilde{e}_k \frac{\tilde{x}_k}{\pi_k} \right), \tilde{0} \right] = 0$

Now,

$$\begin{aligned} \tilde{d} \left[\frac{\tilde{x}_k}{\pi_k} - \sum_{k=1}^n \left(\tilde{e}_k \frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) \right] &= \tilde{d} \left[([x_1, \bar{x}_1], [x_2, \bar{x}_2], \dots, [x_k, \bar{x}_k], \dots) - ([x_1, \bar{x}_1], [x_2, \bar{x}_2], \dots, [x_n, \bar{x}_n]), \tilde{0} \right] \\ &= \tilde{d} \left[([\tilde{0}, \tilde{0}], \dots, [x_{n+1}, \bar{x}_{n+1}], [x_{n+2}, \bar{x}_{n+2}], \dots), \tilde{0} \right] \\ &= \sup_{k \geq n+1} \max \left\{ |x_k|^{1/k}, |\bar{x}_k|^{1/k} \right\} \rightarrow \tilde{0} \text{ as } n \rightarrow \infty \end{aligned}$$

We have, $\frac{\tilde{x}_k}{\pi_k} = \sum_{k=1}^{\infty} \tilde{e}_k \frac{\tilde{x}_k}{\pi_k}$ (2.2)

Let us show the uniqueness of the representation given by (5.2) for $\tilde{x} = \left(\frac{\tilde{x}_k}{\pi_k} \right) \in \Gamma_{\pi} (IR)$

Suppose that there exists a representation $\tilde{x} \frac{\tilde{x}_k}{\pi_k} = \sum_{k=1}^{\infty} \tilde{e}_k \frac{\tilde{y}_k}{\pi_k}$, then,

$$\begin{aligned} \tilde{d} \left(\sum_{k=1}^n \left(\frac{\tilde{x}_k}{\pi_k} - \frac{\tilde{y}_k}{\pi_k} \right) \tilde{e}_k, \tilde{0} \right) &= \sum_{k=1}^n \tilde{d} \left(\left(\frac{\tilde{x}_k}{\pi_k} - \frac{\tilde{y}_k}{\pi_k} \right) \tilde{e}_k, \tilde{0} \right) \\ &= \sup_{k \geq n+1} \max \left\{ \left| \frac{y_k - x_k}{\pi_k} \right|^{1/k}, \left| \frac{\bar{y}_k - \bar{x}_k}{\pi_k} \right|^{1/k} \right\} \rightarrow 0 \text{ as } n \rightarrow \infty \\ &\quad \left| \frac{y_k - x_k}{\pi_k} \right|^{1/k} \rightarrow 0 \text{ and } \left| \frac{\bar{y}_k - \bar{x}_k}{\pi_k} \right|^{1/k} \rightarrow 0 \text{ as } n \rightarrow \infty \end{aligned}$$

Therefore $\underline{y}_k = \underline{x}_k$ and $\bar{y}_k = \bar{x}_k$. That is $\frac{\tilde{y}}{\pi} = \frac{\tilde{x}}{\pi}$.

Theorem 2.5: The β dual of the sequence space of interval numbers $\Gamma_{\pi} (IR)$ is $\Lambda_{\pi} (IR)$.

Proof: Let us suppose that $\tilde{y} = \left(\frac{\tilde{y}_k}{\pi_k} \right) \in \Lambda_{\pi} (IR)$ for every $\tilde{x} = \left(\frac{\tilde{x}_k}{\pi_k} \right) \in \Gamma_{\pi} (IR)$, then

$$\begin{aligned} \lim_n D \left(\sum_{k=1}^n \frac{\tilde{x}_k}{\pi_k} \frac{\tilde{y}_k}{\pi_k}, \tilde{0} \right) &= \lim_n D \left(\sum_{k=1}^n [\underline{y}_k, \bar{y}_k], [\underline{x}_k, \bar{x}_k], \tilde{0} \right) \\ &= \lim_n D \left(\sum_{k=1}^n \left[\min(\underline{y}_k \underline{x}_k, \underline{y}_k \bar{x}_k, \bar{y}_k \underline{x}_k, \bar{y}_k \bar{x}_k), \max(\underline{y}_k \underline{x}_k, \underline{y}_k \bar{x}_k, \bar{y}_k \underline{x}_k, \bar{y}_k \bar{x}_k) \right], \tilde{0} \right) \\ &= \lim_n \max \left\{ \left| \sum_{k=1}^n \underline{y}_k \underline{x}_k \right|^{1/k}, \left| \sum_{k=1}^n \underline{y}_k \bar{x}_k \right|^{1/k}, \left| \sum_{k=1}^n \bar{y}_k \underline{x}_k \right|^{1/k}, \left| \sum_{k=1}^n \bar{y}_k \bar{x}_k \right|^{1/k} \right\} \\ &\leq \lim_n \max \left\{ \sum_{k=1}^n |\underline{y}_k \underline{x}_k|^{1/k}, \sum_{k=1}^n |\underline{y}_k \bar{x}_k|^{1/k}, \sum_{k=1}^n |\bar{y}_k \underline{x}_k|^{1/k}, \sum_{k=1}^n |\bar{y}_k \bar{x}_k|^{1/k} \right\} \\ &= \lim_n M \max \left\{ \sum_{k=1}^n |\underline{x}_k|^{1/k}, \sum_{k=1}^n |\bar{x}_k|^{1/k} \right\} \end{aligned}$$

Where $M = \max\{M_1, M_2\}$ $M_1 = \sup_k |\underline{y}_k|^{1/k}$, $M_2 = \sup_k |\bar{y}_k|^{1/k}$

$$\begin{aligned} \lim_n D \left(\sum_{k=1}^n \frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) &\leq M \lim_n D \left(\sum_{k=1}^n \frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) \\ &= MD \left(\sum_{k=1}^{\infty} \frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) = M \sum_{k=1}^{\infty} D \left(\frac{\tilde{x}_k}{\pi_k}, \tilde{0} \right) < \infty \end{aligned}$$

Therefore, we get $\tilde{x}_k, \tilde{y}_k \in cs(IR)$.

Hence $(\tilde{x}_k) \in [\Gamma_{\pi}(IR)]^{\beta}$

$$\Lambda_{\pi}(IR) \subset [\Gamma_{\pi}(IR)]^{\beta} \tag{2.3}$$

Let $\tilde{y} = \left(\frac{\tilde{y}_k}{\pi_k} \right) \in [\Gamma_{\pi}(IR)]^{\beta}$, then $\sum D \left(\frac{\tilde{x}_k}{\pi_k}, \frac{\tilde{y}_k}{\pi_k}, \tilde{0} \right)$ converges for every $\tilde{x} = \left(\frac{\tilde{x}_k}{\pi_k} \right) \in \Gamma_{\pi}(IR)$

By theorem 2.2, $D \left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0} \right)$ is bounded.

$\sup_k D \left(\frac{\tilde{y}_k}{\pi_k}, \tilde{0} \right)$ exists, then $\tilde{y} = \left(\frac{\tilde{y}_k}{\pi_k} \right) \in \Lambda_{\pi}(IR)$

$$[\Gamma_{\pi}(IR)]^{\beta} \subset \Lambda_{\pi}(IR) \tag{2.4}$$

From equations (2.3) and (2.4), $[\Gamma_{\pi}(IR)]^{\beta} = \Lambda_{\pi}(IR)$

REFERENCES

1. Brown H.I., Entire methods of summation, *Compositio Mathematica*, to me 21, n^o1(1969), 35-42.
2. Dwyer P.S., *Linear Computation*, New York, Wiley, (1951).
3. Dwyer P.S., Error of matrix computation, simultaneous equations and eigenvalues, *National Bureau of Standards, Applied Mathematics Series*, 29(1953), 49-58.
4. Esi A., Strongly almost λ -convergence and statistically almost λ -convergence of interval numbers, *Scientia Magna*, 7(2)(2011), 117-122.
5. Esi A., A new class of interval numbers, *Journal of Qafqaz University, Mathematics and Computer science*, 31(2011), 98-102.
6. Kuo-Ping, Chiao, Fundamental properties of interval vector max-norm, *Tamsui Oxford Journal of Mathematics*, 18(2) (2002), 219-233.
7. Markov S., Quasilinear spaces and their relation to vector spaces, *Electronic Journal on Mathematics of Computation*, 2(1)(2005).
8. Moore R.E., *Automatic Error Analysis in Digital Computation*, LSMD-48421, Lockheed Missiles and space Company, (1959).
9. Moore R.E. and Yang C.T., *Interval Analysis I*, LMSD-285875, Lockheed Missiles and space Company, (1962).
10. Moore R.E. and Yang C.T., *Theory of an interval algebra and its application to numeric analysis*, RAAG Memories II, Gankutsu Bunken Fukeyu-kai, Tokyo, (1958).
11. Sengönül M. and Eryılmaz A., On the sequence space of interval numbers, *Thai Journal of Mathematics*, 8(3) (2010), 503-510.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]