

ARITHMETIC OPERATIONS
ON SYMMETRIC OCTAGONAL INTUITIONISTIC FUZZY NUMBER

G. UTHRA¹, K. THANGAVELU² AND B. AMUTHA^{*3}

¹P.G. & Research Department of Mathematics,
Pachaiyappa's College, Chennai-600 030, India.

²Department of Mathematics,
C. Kandasamy Naidu College for Men, Anna Nagar, Chennai-600 102, India.

³Department of Mathematics, Jeppiaar Engineering College, Chennai - 600 119, India.

(Received On: 03-05-18; Revised & Accepted On: 08-06-18)

ABSTRACT

This paper introduces Symmetric Octagonal Intuitionistic Fuzzy Number (SOIFN) and its arithmetic operations. The arithmetic operations addition, subtraction, scalar multiplication and additive inverse are performed using (α, β) cuts. An example illustrates the proposed arithmetic operations of Symmetric Octagonal Intuitionistic Fuzzy Numbers(SOIFN).

Keywords: Fuzzy set, Intuitionistic Fuzzy Set, Octagonal Intuitionistic Fuzzy Number (OIFN), Symmetric Octagonal Intuitionistic Fuzzy Number (SOIFN).

1. INTRODUCTION

In real life, there are many situations, where it is impossible to get precise data for the cost parameters, due to complexity and uncertainty of information. Therefore, it is desirable to apply fuzzy sets to eliminate the impreciseness and vagueness..

The concept of fuzzy set was introduced by Zadeh [15] in 1965 and it dealt with imprecision, vagueness in real life situations. In 1970, Bellman & Zadeh [4] proposed the concept of decision making problems involving uncertainty and imprecision. K.T. Attanasov [1], [2], [3] introduced Intuitionistic fuzzy sets and their applications to deal with vagueness. W.L. Gau and D.J. Buehrer [7] discussed the concept of vague sets. Bustine and Burillo [5] insisted that vague sets are intuitionistic fuzzy sets. Dhanalakshmi .V, Felbin .C Kennedy, [6] proposed the ranking algorithm for symmetric octagonal fuzzy numbers. G. Menaka[8] and Rajarajeswari, G.Menaka [12] proposed a new approach for ranking of octagonal intuitionistic fuzzy numbers. G. Sahaya sudha, K.R. Vijayalakshmi [13] presented a value and ambiguity based ranking of a symmetric hexagonal intuitionistic fuzzy numbers in decision making. R. Parvathi, C. Malathi [10] listed the Arithmetic Operations on Symmetric Trapezoidal Intuitionistic Fuzzy Numbers using (α, β) cuts. A. Sahaya Sudha and R.Gokilamani [14] proposed an Arithmetic Operations on Hexadecagonal Fuzzy Numbers. K. Ponnivalavan and T. Pathinathan, [11] defined Intuitionistic Pentagonal Fuzzy Numbers and the Arithmetic Operations on Intuitionistic Pentagonal Fuzzy numbers. A. Mohammed Shapique [9] introduced the Heptagonal Fuzzy numbers and the Arithmetic Operations on Heptagonal Fuzzy Numbers using α cut method.

In this paper, SOIFN have been introduced. Arithmetic operations of SOIFN have been proposed using (α, β) cuts. A numerical example for each arithmetic operation is provided.

The rest of this paper organized as follows. In Section 2, definitions of fuzzy set, intuitionistic fuzzy set and OIFN are given. Section 3, presents introduction of SOIFN. In Section 4, arithmetic operations of SOIFN based on (α, β) cuts are proposed. A numerical example showing each arithmetic operation is illustrates in section5, followed by conclusion in section-6.

Corresponding Author: B. Amutha^{*3}

³Department of Mathematics, Jeppiaar Engineering College, Chennai - 600 119, India.

2. DEFINITIONS

Definition 2.1: Fuzzy Set

Let A be a classical set, $\mu_A(x)$ be a function from A to $[0, 1]$. A fuzzy set A with the membership function $\mu_A(x)$ is defined as $A = \{x, \mu_A(x); x \in A \text{ and } \mu_A(x) \in [0, 1]\}$.

Definition 2.2: Intuitionistic Fuzzy Set

Let X be a given set. An Intuitionistic Fuzzy Set A in X is given by, where $A = \{(x, \mu_A(x), \vartheta_A(x)) \mid x \in X\}$ $X \rightarrow [0, 1]$, where $\mu_A(x)$ is the degree of membership of the element x in A and, $\vartheta_A(x)$ is the degree of non membership of x in A and $0 \leq \mu_A(x) + \vartheta_A(x) \leq 1$.

Definition 2.3: Octagonal Intuitionistic Fuzzy Number (OIFN)

An OIFN is specified by $A = [(a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8); (a'_1, a'_2, a'_3, a'_4, a'_5, a'_6, a'_7, a'_8)]$ where $a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a'_1, a'_2, a'_3, a'_4, a'_5, a'_6, a'_7, a'_8$ are real numbers and its membership and non-membership functions are given below.

$$\mu_A(x) = \begin{cases} 0 & \text{if } x < a_1 \\ k \left(\frac{x - a_1}{a_2 - a_1} \right) & \text{if } a_1 \leq x \leq a_2 \\ k & \text{if } a_2 \leq x \leq a_3 \\ k + (1 - k) \left(\frac{x - a_3}{a_4 - a_3} \right) & \text{if } a_3 \leq x \leq a_4 \\ 1 & \text{if } a_4 \leq x \leq a_5 \\ k + (1 - k) \left(\frac{a_6 - x}{a_6 - a_5} \right) & \text{if } a_5 \leq x \leq a_6 \\ k & \text{if } a_6 \leq x \leq a_7 \\ k \left(\frac{a_8 - x}{a_8 - a_7} \right) & \text{if } a_7 \leq x \leq a_8 \end{cases}$$

$$\vartheta_A(x) = \begin{cases} 1 & \text{if } a'_1 \leq x \\ k + (1 - k) \left(\frac{a'_2 - x}{a'_2 - a'_1} \right) & \text{if } a'_1 \leq x \leq a'_2 \\ k & \text{if } a'_2 \leq x \leq a'_3 \\ k \left(\frac{a_4 - x}{a_4 - a'_3} \right) & \text{if } a'_3 \leq x \leq a_4 \\ 0 & \text{if } a_4 \leq x \leq a_5 \\ k \left(\frac{x - a_5}{a'_6 - a_5} \right) & \text{if } a_5 \leq x \leq a'_6 \\ k & \text{if } a'_6 \leq x \leq a'_7 \\ k + (1 - k) \left(\frac{x - a'_7}{a'_8 - a'_7} \right) & \text{if } a'_7 \leq x \leq a'_8 \end{cases}$$

Where $k = 1/2$




Figure-1: Diagrammatic Representation of OIFN

3. SYMMETRIC OCTAGONAL INTUITIONISTIC FUZZY NUMBER (SOIFN)

A **SOIFN** is given by $A_o = [(a_L - r - s - t, a_L - r - s, a_L - r, a_L, a_U, a_U + r, a_U + r + s, a_U + r + s + t); (a'_L - r' - s' - t', a'_L - r' - s', a'_L - r', a_L, a_U, a_U' + r', a_U' + r' + s', a_U' + r' + s' + t')]$ where $a_L - r - s - t, a_L - r - s, a_L - r, a_L, a_U, a_U + r, a_U + r + s, a_U + r + s + t, a'_L - r' - s' - t', a'_L - r' - s', a'_L - r', a_L, a_U, a_U' + r', a_U' + r' + s', a_U' + r' + s' + t'$ are real numbers such that $(a'_L - r' - s' - t' \leq a_L - r - s - t \leq a'_L - r' - s' \leq a_L - r - s \leq a'_L - r' \leq a_L - r \leq a_L \leq a_U \leq a_U + r \leq a_U' + r' \leq a_U + r + s \leq a_U' + r' + s' \leq a_U + r + s + t \leq a_U' + r' + s' + t')$ and its membership and non-membership functions are given below

$$\mu_{A_o}(x) = \begin{cases} \frac{x - (a_L - r - s - t)}{2t}, & a_L - r - s - t \leq x \leq a_L - r - s \\ \frac{1}{2}, & a_L - r - s \leq x \leq a_L - r \\ \frac{1}{2} + \frac{(x - (a_L - r))}{2r}, & a_L - r \leq x \leq a_L \\ 1, & a_L \leq x \leq a_U \\ \frac{1}{2} + \frac{((a_U + r) - x)}{2r}, & a_U \leq x \leq a_U + r \\ \frac{1}{2}, & a_U + r \leq x \leq a_U + r + s \\ \frac{((a_U + r + s + t) - x)}{2t}, & a_U + r + s \leq x \leq a_U + r + s + t \end{cases}$$

$$\vartheta_{A_o}(x) = \begin{cases} \frac{1}{2} + \frac{(a'_L - r' - s') - x}{2t'}, & a'_L - r' - s' - t' \leq x \leq a'_L - r' - s' \\ \frac{1}{2}, & a'_L - r' - s' \leq x \leq a'_L - r' \\ \frac{1}{2} \left(\frac{a_L - x}{a_L - (a'_L - r')} \right), & a'_L - r' \leq x \leq a_L \\ 0, & a_L \leq x \leq a_U \\ \frac{1}{2} \left(\frac{x - a_U}{(a'_U + r') - a_U} \right), & a_U \leq x \leq a'_U + r' \\ \frac{1}{2}, & a'_U + r' \leq x \leq a'_U + r' + s' \\ \frac{1}{2} + \left(\frac{(x - (a'_U + r' + s'))}{2t'} \right), & a'_U + r' + s' \leq x \leq a'_U + r' + s' + t' \end{cases}$$




Figure-2: Diagrammatic representation of SOIFN

Where $a_{L1} = a_L - r - s - t, a_{L2} = a_L - r - s, a_{L3} = a_L - r, a_{U1} = a_U + r, a_{U2} = a_U + r + s, a_{U3} = a_U + r + s + t, a'_{L1} = a'_L - r' - s' - t', a'_{L2} = a'_L - r' - s', a'_{L3} = a'_L - r', a'_{U1} = a'_U + r', a'_{U2} = a'_U + r' + s', a'_{U3} = a'_U + r' + s' + t'$

4. ARITHMETIC OPERATIONS ON SOIFN BY (α, β) CUTS

1. Addition:

If $A_o = [(a_L - r_1 - s_1 - t_1, a_L - r_1 - s_1, a_L - r_1, a_L, a_U, a_U + r_1, a_U + r_1 + s_1, a_U + r_1 + s_1 + t_1); (a'_L - r'_1 - s'_1 - t'_1, a'_L - r'_1 - s'_1, a'_L - r'_1, a'_L, a'_U, a'_U + r'_1, a'_U + r'_1 + s'_1, a'_U + r'_1 + s'_1 + t'_1)]$
 $B_o = [(b_L - r_2 - s_2 - t_2, b_L - r_2 - s_2, b_L - r_2, b_L, b_U, b_U + r_2, b_U + r_2 + s_2, b_U + r_2 + s_2 + t_2); (b'_L - r'_2 - s'_2 - t'_2, b'_L - r'_2 - s'_2, b'_L - r'_2, b'_L, b'_U, b'_U + r'_2, b'_U + r'_2 + s'_2, b'_U + r'_2 + s'_2 + t'_2)]$ are two SOIFNs then $C_o = A_o + B_o$ is also a SOIFNs and is given by $C_o = (((a_L + b_L) - (r + s + t), (a_L + b_L) - (r + s), (a_L + b_L) - r, a_L + b_L, a_U + b_U, (a_U + b_U) + r, (a_U + b_U) + (r + s), (a_U + b_U) + (r + s + t)); ((a'_L + b'_L)(r' + s' + t'), (a'_L + b'_L) - (r' + s'), (a'_L + b'_L) - r', a'_L + b'_L, a'_U + b'_U, (a'_U + b'_U) + r', (a'_U + b'_U) + (r' + s'), (a'_U + b'_U) + (r' + s' + t')))$
 Where $r_1 + r_2 = r, s_1 + s_2 = s, t_1 + t_2 = t, t'_1 + t'_2 = t', s'_1 + s'_2 = s', r'_1 + r'_2 = r'$

Proof: The membership and non-membership function of IFS $C_o = A_o + B_o$ can be obtained by (α, β) cuts with the transformation $z = x + y$. α -cut for membership function of A_o is $((a_L - r_1 - s_1 - t_1) + 2\alpha t_1, 2\alpha r_1 - r_1 + (a_L - r_1), r_1 - 2\alpha r_1 + (a_U + r_1), (a_U + r_1 + s_1 + t_1) - 2\alpha t_1)$ for every $\alpha \in [0, 1]$. That is $x \in ((a_L - r_1 - s_1 - t_1) + 2\alpha t_1, 2\alpha r_1 - r_1 + (a_L - r_1), r_1 - 2\alpha r_1 + (a_U + r_1), (a_U + r_1 + s_1 + t_1) - 2\alpha t_1)$. α -cut for membership function of B_o is $((b_L - r_2 - s_2 - t_2) + 2\alpha t_2, 2\alpha r_2 - r_2 + (b_L - r_2), r_2 - 2\alpha r_2 + (b_U + r_2), (b_U + r_2 + s_2 + t_2) - 2\alpha t_2)$ for every $\alpha \in [0, 1]$. That is $y \in ((b_L - r_2 - s_2 - t_2) + 2\alpha t_2, 2\alpha r_2 - r_2 + (b_L - r_2), r_2 - 2\alpha r_2 + (b_U + r_2), (b_U + r_2 + s_2 + t_2) - 2\alpha t_2)$. So $z = (x + y) \in ((a_L + b_L) - (r + s + t) + 2\alpha t, r(2\alpha - 1) + [(a_L + b_L) - r], r(1 - 2\alpha) + [(a_U + b_U) + r], [(a_U + b_U) + (r + s + t)] - 2\alpha t)$.

Where $r_1 + r_2 = r, s_1 + s_2 = s, t_1 + t_2 = t$

Therefore the membership function of $C_o = A_o + B_o$ is given by

$$\mu_{C_o}(z) = \begin{cases} \frac{z - [(a_L + b_L) - (r + s + t)]}{2t}, & (a_L + b_L) - (r + s + t) \leq z \leq (a_L + b_L) - (r + s) \\ \frac{1}{2}, & (a_L + b_L) - (r + s) \leq z \leq (a_L + b_L) - r \\ \frac{1}{2} + \frac{(z - ((a_L + b_L) - r))}{2r}, & (a_L + b_L) - r \leq z \leq a_L + b_L \\ \frac{1}{2}, & a_L + b_L \leq z \leq a_U + b_U \\ \frac{1}{2} + \frac{(((a_U + b_U) + r) - z)}{2r}, & a_U + b_U \leq z \leq (a_U + b_U) + r \\ \frac{1}{2}, & (a_U + b_U) + r \leq z \leq (a_U + b_U) + r + s \\ \frac{((a_U + b_U) + r + s + t) - z}{2t}, & (a_U + b_U) + r + s \leq z \leq (a_U + b_U) + r + s + t \end{cases}$$

For non-membership function β -cut for A'_o is $((t'_1 - 2\beta t'_1) + (a'_L - r'_1 - s'_1), a_L - 2\beta(a_L - (a'_L - r'_1)), a_U + 2\beta((a'_U + r'_1) - a_U), 2\beta t'_1 - t'_1 + (a'_U + r'_1 + s'_1))$ for every $\beta \in [0, 1]$. That is $x \in ((t'_1 - 2\beta t'_1) + (a'_L - r'_1 - s'_1), a_L - 2\beta(a_L - (a'_L - r'_1)), a_U + 2\beta((a'_U + r'_1) - a_U), 2\beta t'_1 - t'_1 + (a'_U + r'_1 + s'_1))$ β -cut for non membership function of B'_o is $((t'_2 - 2\beta t'_2) + (b'_L - r'_2 - s'_2), b_L - 2\beta(b_L - (b'_L - r'_2)), b_U + 2\beta((b'_U + r'_2) - b_U), 2\beta t'_2 - t'_2 + (b'_U + r'_2 + s'_2))$ for every $\beta \in [0, 1]$. That is $y \in ((t'_2 - 2\beta t'_2) + (b'_L - r'_2 - s'_2), b_L - 2\beta(b_L - (b'_L - r'_2)), b_U + 2\beta((b'_U + r'_2) - b_U), 2\beta t'_2 - t'_2 + (b'_U + r'_2 + s'_2))$. so $z = (x + y) \in ((1 - 2\beta)t'(a'_L + b'_L) - (r' + s'), (a_L + b_L) - 2\beta((a_L + b_L) - (a'_L + b'_L) - r'), (a_U + b_U) + 2\beta((a'_U + b'_U) + r' - (a_U + b_U)), (2\beta - 1)t' + ((a'_U + b'_U) + (r' + s')))$
 Where $t'_1 + t'_2 = t', s'_1 + s'_2 = s', r'_1 + r'_2 = r$.

Therefore the non-membership function of $C'_o = A'_o + B'_o$ is given by

$$\vartheta_{C_0}(z) = \left\{ \begin{array}{ll} \frac{1}{2} + \frac{((a'_L + b'_L) - (r' + s')) - z}{2t'}, & (a'_L + b'_L) - (r' + s' + t') \leq z \leq (a'_L + b'_L) - (r' + s') \\ \frac{1}{2}, & (a'_L + b'_L) - (r' + s') \leq z \leq (a'_L + b'_L) - r' \\ \frac{1}{2} \left(\frac{(a_L + b_L) - z}{(a_L + b_L) - ((a'_L + b'_L) - r')} \right) & (a'_L + b'_L) - r' \leq z \leq a_L + b_L \\ 0 & a_L + b_L \leq z \leq a_U + b_U \\ \frac{1}{2} \left(\frac{z - (a_U + b_U)}{((a'_U + b'_U) + r') - (a_U + b_U)} \right) & a_U + b_U \leq z \leq (a'_U + b'_U) + r' \\ \frac{1}{2}, & (a'_U + b'_U) + r' \leq z \leq (a'_U + b'_U) + (r' + s') \\ \frac{1}{2} + \left(\frac{(x - [(a'_U + b'_U) + (r' + s')])}{2t'} \right), & (a'_U + b'_U) + (r' + s') \leq z \leq (a'_U + b'_U) + (r' + s' + t') \end{array} \right.$$

2. Subtraction

If $A_o = [(a_L - r_1 - s_1 - t_1, a_L - r_1 - s_1, a_L - r_1, a_L, a_U, a_U + r_1, a_U + r_1 + s_1, a_U + r_1 + s_1 + t_1); (a'_L - r'_1 - s'_1 - t'_1, a'_L - r'_1 - s'_1, a'_L - r'_1, a'_L, a'_U, a'_U + r'_1, a'_U + r'_1 + s'_1, a'_U + r'_1 + s'_1 + t'_1)]$ $B_o = [(b_L - r_2 - s_2 - t_2, b_L - r_2 - s_2, b_L - r_2, b_L, b_U, b_U + r_2, b_U + r_2 + s_2, b_U + r_2 + s_2 + t_2); (b'_L - r'_2 - s'_2 - t'_2, b'_L - r'_2 - s'_2, b'_L - r'_2, b'_L, b'_U, b'_U + r'_2, b'_U + r'_2 + s'_2, b'_U + r'_2 + s'_2 + t'_2)]$ are two SOIFNs then $C_o = A_o - B_o$ is also a SOIFNs and is given by

$$C_o = (((a_L - b_U) - (r + s + t), (a_L - b_U) - (r + s), (a_L - b_U) - r, a_L - b_L, a_U - b_U, (a_U - b_L) + r, (a_U - b_L) + (r + s), (a_U - b_L) + (r + s + t)); ((a'_L - b'_U) - (r' + s' + t'), (a'_L - b'_U) - (r' + s'), (a'_L - b'_U) - r', a'_L - b'_L, a'_U - b'_U, (a'_U - b'_L) + r', (a'_U - b'_L) + (r' + s'), (a'_U - b'_L) + (r' + s' + t')))$$

Where $r_1 + r_2 = r, s_1 + s_2 = s, t_1 + t_2 = t, t'_1 + t'_2 = t', s'_1 + s'_2 = s', r'_1 + r'_2 = r'$.

Proof: The membership and non-membership function of IFS $C_o = A_o - B_o$ can be obtained by (α, β) cuts with the transformation $z = x - y$. α -cut for membership function of A_o is $[(a_L - r_1 - s_1 - t_1) + 2\alpha t_1, 2\alpha r_1 - r_1 + (a_L - r_1), r_1 - 2\alpha r_1 + (a_U + r_1), (a_U + r_1 + s_1 + t_1) - 2\alpha t_1]$ for every $\alpha \in [0, 1]$. That is $x \in [(a_L - r_1 - s_1 - t_1) + 2\alpha t_1, 2\alpha r_1 - r_1 + (a_L - r_1), r_1 - 2\alpha r_1 + (a_U + r_1), (a_U + r_1 + s_1 + t_1) - 2\alpha t_1]$. α -cut for membership function of B_o is $[(b_L - r_2 - s_2 - t_2) + 2\alpha t_2, 2\alpha r_2 - r_2 + (b_L - r_2), r_2 - 2\alpha r_2 + (b_U + r_2), (b_U + r_2 + s_2 + t_2) - 2\alpha t_2]$ for every $\alpha \in [0, 1]$. That is $y \in [(b_L - r_2 - s_2 - t_2) + 2\alpha t_2, 2\alpha r_2 - r_2 + (b_L - r_2), r_2 - 2\alpha r_2 + (b_U + r_2), (b_U + r_2 + s_2 + t_2) - 2\alpha t_2]$. So $z = (x - y) \in [(a_L - b_U) - (r + s + t) + 2\alpha t, r(2\alpha - 1) + (a_L - b_U) - r, r(1 - 2\alpha) + (a_U - b_L) + r, (a_U - b_L) + (r + s + t) - 2\alpha t]$ Where $r_1 + r_2 = r, s_1 + s_2 = s, t_1 + t_2 = t$.

Therefore the membership function of $C_o = A_o - B_o$ is given by

$$\mu_{C_o}(z) = \left\{ \begin{array}{ll} \frac{z - [(a_L - b_U) - (r + s + t)]}{2t}, & (a_L - b_U) - (r + s + t) \leq z \leq (a_L - b_U) - (r + s) \\ \frac{1}{2}, & (a_L - b_U) - (r + s) \leq z \leq (a_L - b_U) - r \\ \frac{1}{2} + \frac{(z - ((a_L - b_U) - r))}{2r} & (a_L - b_U) - r \leq z \leq a_L - b_L \\ \frac{1}{2} & a_L - b_L \leq z \leq a_U - b_U \\ \frac{1}{2} + \frac{((a_U - b_L) + r) - z}{2r} & a_U - b_U \leq z \leq (a_U - b_L) + r \\ \frac{1}{2}, & (a_U - b_L) + r \leq z \leq (a_U - b_L) + (r + s) \\ \frac{((a_U - b_L) + (r + s + t)) - z}{2t}, & (a_U - b_L) + r + s \leq z \leq (a_U - b_L) + r + s + t \end{array} \right.$$

For non-membership function β -cut for A_o' is $((t'_1 - 2\beta t'_1) + (a'_L - r'_1 - s'_1), a_L - 2\beta(a_L - (a'_L - r'_1)), a_U + 2\beta((a'_U + r'_1) - a_U), 2\beta t'_1 - t'_1 + (a'_U + r'_1 + s'_1))$ for every $\beta \in [0, 1]$. That is $x \in ((t'_1 - 2\beta t'_1) + (a'_L - r'_1 - s'_1), a_L - 2\beta(a_L - (a'_L - r'_1)), a_U + 2\beta((a'_U + r'_1) - a_U), 2\beta t'_1 - t'_1 + (a'_U + r'_1 + s'_1))$. β -cut for non membership function of B_o' is $((t'_2 - 2\beta t'_2) + (b'_L - r'_2 - s'_2), b_L - 2\beta(b_L - (b'_L - r'_2)), b_U + 2\beta((b'_U + r'_2) - b_U), 2\beta t'_2 - t'_2 + (b'_U + r'_2 + s'_2))$ for every $\beta \in [0, 1]$. That is $y \in ((t'_2 - 2\beta t'_2) + (b'_L - r'_2 - s'_2), b_L - 2\beta(b_L - (b'_L - r'_2)), b_U + 2\beta((b'_U + r'_2) - b_U), 2\beta t'_2 - t'_2 + (b'_U + r'_2 + s'_2))$. So $z = (x - y) \in [(1 - 2\beta)t' + (a'_L - b'_U) - (r' + s'), (a_L - b_L) - 2\beta((a_L - b_L) - (a'_L - b'_U) - r'), (a_U - b_U) + 2\beta((a'_U - b'_L) + r') - (a_U - b_U), (2\beta - 1)t' + ((a'_U - b'_L) + (r' + s'))]$

Where $t'_1 + t'_2 = t', s'_1 + s'_2 = s', r'_1 + r'_2 = r'$

Therefore the non-membership function of $C_o' = A_o' - B_o'$ is given by

$$\vartheta_{C_o'}(z) = \begin{cases} \frac{1}{2} + \frac{((a_L' - b_U') - (r' + s')) - z}{2t'}, & (a_L' - b_U') - (r' + s' + t') \leq z \leq (a_L' - b_U') - (r' + s') \\ \frac{1}{2}, & (a_L' - b_U') - (r' + s') \leq z \leq (a_L' - b_U') - r' \\ \frac{1}{2} \left(\frac{(a_L - b_L) - z}{(a_L - b_L) - ((a_L' - b_U') - r')} \right), & (a_L' - b_U') - r' \leq z \leq a_L - b_L \\ 0, & a_L - b_L \leq z \leq a_U - b_U \\ \frac{1}{2} \left(\frac{z - (a_U - b_U)}{((a_U' - b_L') + r') - (a_U - b_U)} \right), & a_U - b_U \leq z \leq (a_U' - b_L') + r' \\ \frac{1}{2}, & (a_U' - b_L') + r' \leq z \leq (a_U' - b_L') + (r' + s') \\ \frac{1}{2} + \left(\frac{(z - (a_U' - b_L') + (r' + s'))}{2t'} \right), & (a_U' - b_L') + (r' + s') \leq z \leq (a_U' - b_L') + (r' + s' + t') \end{cases}$$

3. Scalar Multiplication

Let $k \in \mathbb{R}$ if $A_o = [(a_L - r - s - t, a_L - r - s, a_L - r, a_L, a_U, a_U + r, a_U + r + s, a_U + r + s + t); (a_L' - r' - s' - t', a_L' - r' - s', a_L - r', a_L, a_U, a_U' + r', a_U' + r' + s', a_U' + r' + s' + t')]$ is a SOIFN then $C_o = kA_o$ is also a SOIFN and is given by

$$C_o = \begin{cases} [k(a_L - r - s - t), k(a_L - r - s), k(a_L - r), ka_L, ka_U, k(a_U + r), k(a_U + r + s), k(a_U + r + s + t); k(a_L' - r' - s' - t'), k(a_L' - r' - s'), k(a_L' - r'), ka_L, ka_U, k(a_U' + r'), k(a_U' + r' + s'), k(a_U' + r' + s' + t')]; & k > 0 \\ [-k(a_U + r + s + t), -k(a_U + r + s), -k(a_U + r), -ka_U, -ka_L, -k(a_U - r), -k(a_U - r - s), -k(a_U - r - s - t); -k(a_U' + r' + s' + t'), -k(a_U' + r' + s'), -k(a_L' + r'), -ka_U, -ka_L, -k(a_U' - r'), -k(a_U' - r' - s'), -k(a_U' - r' - s' - t')]; & k < 0 \end{cases}$$

Proof:

Case-(i): $k > 0$

The membership and non-membership function of IFS $C_o = kA_o$ can be obtained by (α, β) cuts with the transformation $z = kx$. α -cut for membership function of A_o is $[(a_L - r - s - t) + 2\alpha t, 2\alpha r - r + (a_L - r), r - 2\alpha r + (a_U + r), (a_U + r + s + t) - 2\alpha t]$ for every $\alpha \in [0, 1]$. That is $x \in [(a_L - r - s - t) + 2\alpha t, 2\alpha r - r + (a_L - r), r - 2\alpha r + (a_U + r), (a_U + r + s + t) - 2\alpha t]$. So $z = kx \in [k(a_L - r - s - t) + 2\alpha kt, 2\alpha kr - kr + k(a_L - r), rk - 2\alpha kr + k(a_U + r), k(a_U + r + s + t) - 2\alpha kt]$ Where $r_1 + r_2 = r, s_1 + s_2 = s, t_1 + t_2 = t$.

Therefore the membership function of $C_o = kA_o$ is given by

$$\mu_{C_o}(z) = \begin{cases} \frac{z - k(a_L - r - s - t)}{2kt}, & k(a_L - r - s - t) \leq z \leq k(a_L - r - s) \\ \frac{1}{2}, & k(a_L - r - s) \leq z \leq k(a_L - r) \\ \frac{1}{2} + \frac{(z - k(a_L - r))}{2kr}, & k(a_L - r) \leq z \leq ka_L \\ 1, & ka_L \leq z \leq ka_U \\ \frac{1}{2} + \frac{(k(a_U + r) - z)}{2kr}, & ka_U \leq z \leq k(a_U + r) \\ \frac{1}{2}, & k(a_U + r) \leq z \leq k(a_U + r + s) \\ \frac{(k(a_U + r + s + t) - z)}{2kt}, & k(a_U + r + s) \leq z \leq k(a_U + r + s + t) \end{cases}$$

For non-membership function β -cut for A_o is $((t' - 2\beta t') + (a_L' - r' - s'), a_L - 2\beta(a_L - (a_L' - r'))), a_U + 2\beta((a_U' + r') - a_U), 2\beta t' - t' + (a_U' + r' + s')$ for every $\beta \in [0, 1]$. That is $x \in ((t' - 2\beta t') + (a_L' - r' - s'), a_L - 2\beta(a_L - (a_L' - r'))), a_U + 2\beta((a_U' + r') - a_U), 2\beta t' - t' + (a_U' + r' + s'))$. So $z = kx \in (k(t' - 2\beta t') + k(a_L' - r' - s'), ka_L - 2k\beta(a_L - (a_L' - r')), ka_U + 2k\beta((a_U' + r') - a_U), k(2\beta - 1)t' + k(a_U' + r' + s'))$

Therefore the non membership function of $C_o = kA_0$ is given by

$$\vartheta_{C_o}(z) = \left\{ \begin{array}{ll} \frac{1}{2} + \frac{k(a'_L - r' - s') - z}{2kt'}, & k(a'_L - r' - s' - t') \leq z \leq k(a'_L - r' - s') \\ \frac{1}{2}, & k(a'_L - r' - s') \leq z \leq k(a'_L - r') \\ \frac{1}{2} \left(\frac{ka'_L - z}{k(a'_L - (a'_L - r'))} \right) & k(a'_L - r') \leq z \leq ka'_L \\ 0 & ka'_L \leq z \leq ka_U \\ \frac{1}{2} \left(\frac{z - ka_U}{k((a'_U + r') - a_U)} \right) & ka_U \leq z \leq k(a'_U + r') \\ \frac{1}{2}, & k(a'_U + r') \leq z \leq k(a'_U + r' + s') \\ \frac{1}{2} + \left(\frac{(z - (a'_U + r' + s'))}{2kt'} \right), & k(a'_U + r' + s') \leq z \leq k(a'_U + r' + s' + t') \end{array} \right.$$

Case-(ii): $k < 0$

The membership and non-membership function of IFS $C_o = -kA_0$ can be obtained by (α, β) cuts with the transformation $z = -kx$. α -cut for membership function of A_0 is $[(a_L - r - s - t) + 2\alpha t, 2\alpha r - r + (a_L - r), r - 2\alpha r + (a_U + r), (a_U + r + s + t) - 2\alpha t]$ for every $\alpha \in [0, 1]$. That is $x \in ((a_L - r - s - t) + 2\alpha t, 2\alpha r - r + (a_L - r), r - 2\alpha r + (a_U + r), (a_U + r + s + t) - 2\alpha t)$. So $z = -kx \in (-k(a_L - r - s - t) - 2kat, -k(2\alpha r - r) - k(a_L - r), -k(r - 2\alpha r) - k(a_U + r), -k(a_U + r + s + t) + 2kat)$.

Therefore the membership function of $C_o = -kA_0$ is given by

$$\mu_{C_o}(z) = \left\{ \begin{array}{ll} \frac{z - (-k(a_U + r + s + t))}{2kt}, & -k(a_U + r + s + t) \leq z \leq -k(a_U + r + s) \\ \frac{1}{2}, & -k(a_U + r + s) \leq z \leq -k(a_U + r) \\ \frac{1}{2} + \frac{z - (-k(a_U + r))}{2kr} & -k(a_U + r) \leq z \leq -ka_U \\ \frac{1}{2} & -ka_U \leq z \leq -ka_L \\ \frac{1}{2} + \frac{(-k(a_L - r) - z)}{2kr} & -ka_L \leq z \leq -k(a_L - r) \\ \frac{1}{2}, & -k(a_L - r) \leq z \leq -k(a_L - r - s) \\ \frac{-k(a_L - r - s - t) - z}{2kt}, & -k(a_L - r - s) \leq z \leq -k(a_L - r - s - t) \end{array} \right.$$

For non-membership function β -cut for A_0 is $(t' - 2\beta t') + (a'_L - r' - s')$, $a_L - 2\beta(a_L - (a'_L - r'))$, $a_U + 2\beta((a'_U + r') - a_U)$, $2\beta t' - t' + (a'_U + r' + s')$ for every $\beta \in [0, 1]$. That is $x \in ((t' - 2\beta t') + (a'_L - r' - s'), a_L - 2\beta(a_L - (a'_L - r')), a_U + 2\beta((a'_U + r') - a_U), 2\beta t' - t' + (a'_U + r' + s'))$. So $z = -kx \in (-k(t' - 2\beta t') - k(a'_L - r' - s'), -ka_L + 2k\beta(a_L - (a'_L - r')), -ka_U - 2k\beta((a'_U + r') - a_U), -k(2\beta t' - t') - k(a'_U + r' + s'))$.

Therefore the non membership function of $C_o = -kA_0$ is given by

$$\vartheta_{C_0}(z) = \left\{ \begin{array}{ll} \frac{1}{2} + \frac{(-k)(a'_U + r' + s') - z}{2kt'}, & -k(a'_U + r' + s' + t') \leq z \leq -k(a'_U + r' + s') \\ \frac{1}{2}, & -k(a'_U + r' + s') \leq z \leq -k(a'_U + r') \\ \frac{1}{2} \left(\frac{(-k)a_U - z}{k((a'_U + r') - a_U)} \right) & -k(a'_U + r') \leq z \leq -ka_U \\ 0 & -ka_U \leq z \leq -ka_L \\ \frac{1}{2} \left(\frac{z - (-k)a_L}{k(a_L - (a'_L - r'))} \right) & -ka_L \leq z \leq -k(a'_L - r') \\ \frac{1}{2}, & -k(a'_L - r') \leq z \leq -k(a'_L - r' - s') \\ \frac{1}{2} + \left(\frac{(z - (-k)(a'_L - r' - s'))}{2kt'} \right), & -k(a'_L - r' - s') \leq z \leq -k(a'_L - r' - s' - t') \end{array} \right.$$

4. Additive Image

If $A_o = [(a_L - r - s - t, a_L - r - s, a_L - r, a_L, a_U, a_U + r, a_U + r + s, a_U + r + s + t); (a'_L - r' - s' - t', a'_L - r' - s', a'_L - r', a'_L, a'_U, a'_U + r', a'_U + r' + s', a'_U + r' + s' + t')]$ additive image of A_o' is also a SOIFN and is given by $A_o' = (- (a_U + r + s + t), -(a_U + r + s), -(a_U + r), -a_U, -a_L, -(a_L - r), -(a_L - r - s), -(a_L - r - s - t) - (a'_U + r' + s' + t'), -(a'_U + r' + s'), -(a'_U + r'), -a_U, -a_L, -(a'_L - r'), -(a'_L - r' - s'), -(a'_L - r' - s' - t'))$

Proof: The membership and non-membership function of IFS $A'_o = -A_o$ can be obtained by (α, β) cuts with the transformation $z = -x$. α -cut for membership function of A_o is $[(a_L - r - s - t) + 2\alpha t, 2\alpha r - r + (a_L - r), r - 2\alpha r + (a_U + r), (a_U + r + s + t) - 2\alpha t]$ for every $\alpha \in [0, 1]$. That is $x \in ((a_L - r - s - t) + 2\alpha t, 2\alpha r - r + (a_L - r), r - 2\alpha r + (a_U + r), (a_U + r + s + t) - 2\alpha t)$. So $z = -x \in [-(a_U + r + s + t) + 2\alpha t, r(2\alpha - 1) - (a_U + r), r(1 - 2\alpha) - (a_L - r), (a_L - r - s - t) - 2\alpha t]$.

Therefore the membership function of $A'_o = -A_o$ is given by

$$\mu_{A'_o}(z) = \left\{ \begin{array}{ll} \frac{z - (-(a_U + r + s + t))}{2t}, & -(a_U + r + s + t) \leq z \leq -(a_U + r + s) \\ \frac{1}{2}, & -(a_U + r + s) \leq z \leq -(a_U + r) \\ \frac{1}{2} + \frac{z - (-(a_U + r))}{2r}, & -(a_U + r) \leq z \leq -a_U \\ \frac{1}{2}, & -a_U \leq z \leq -a_L \\ \frac{1}{2} + \frac{(-(a_L - r) - z)}{2r}, & -a_L \leq z \leq -(a_L - r) \\ \frac{1}{2}, & -(a_L - r) \leq z \leq -(a_L - r - s) \\ \frac{(-(a_L - r - s - t) - z)}{2t}, & -(a_L - r - s) \leq z \leq -(a_L - r - s - t) \end{array} \right.$$

For non-membership function β -cut for A_o is $((t' - 2\beta t') + (a'_L - r' - s'), a_L - 2\beta(a_L - (a'_L - r')), a_U + 2\beta((a'_U + r') - a_U), 2\beta t' - t' + (a'_U + r' + s'))$ for every $\beta \in [0, 1]$. That is $x \in ((t' - 2\beta t') + (a'_L - r' - s'), a_L - 2\beta(a_L - (a'_L - r')), a_U + 2\beta((a'_U + r') - a_U), 2\beta t' - t' + (a'_U + r' + s'))$. So $z = -x \in ((t' - 2\beta t') - (a'_U + r' + s'), -a_U - 2\beta((a'_U + r') - a_U), -a_L + 2\beta(a_L - (a'_L - r')), (2\beta t' - t') - (a'_L - r' - s'))$

Therefore the non membership function of $A'_o = -A_o$ is given by

$$\vartheta_{A_0'}(z) = \begin{cases} \frac{1}{2} + \frac{-(a'_U + r' + s') - z}{2t'}, & -(a'_U + r' + s' + t') \leq z \leq -(a'_U + r' + s') \\ \frac{1}{2}, & -(a'_U + r' + s') \leq z \leq -(a'_U + r') \\ \frac{1}{2} \left(\frac{-a_U - z}{(a'_U + r') - a_U} \right), & -(a'_U + r') \leq z \leq -a_U \\ 0, & -a_U \leq z \leq -a_L \\ \frac{1}{2} \left(\frac{z - (-a_L)}{a_L - (a'_L - r')} \right), & -a_L \leq z \leq -(a'_L - r') \\ \frac{1}{2}, & -(a'_L - r') \leq z \leq -(a'_L - r' - s') \\ \frac{1}{2} + \left(\frac{(z - (-a_L - r' - s'))}{2t'} \right), & -(a'_L - r' - s') \leq z \leq -(a'_L - r' - s' - t') \end{cases}$$

5. NUMERICAL EXAMPLE

Consider two SOIFNS $A = [(7, 9, 11, 13, 15, 17, 19, 21); (4, 7, 10, 13, 15, 18, 21, 24)]$ and

$B = [(8, 10, 12, 14, 16, 18, 20, 22); (5, 8, 11, 14, 16, 19, 22, 25)]$ then

(i) Addition

$A + B = [(7+8, 9+10, 11+12, 13+14, 15+16, 17+18, 19+20, 21+22); (4+5, 7+8, 10+11, 13+14, 15+16, 18+19, 21+22, 24+25)]$
 $= [(15, 19, 23, 27, 31, 35, 39, 43); (9, 15, 21, 27, 31, 37, 43, 49)]$

(ii) Subtraction

$A - B = [(7 - 22, 9 - 20, 11 - 18, 13 - 14, 15 - 16, 17 - 12, 19 - 10, 21 - 8); (4 - 25, 7 - 22, 10 - 19, 13 - 14, 15 - 16, 18 - 11, 21 - 8, 24 - 5)]$
 $= [(-15, -11, -7, -1, -1, 5, 9, 13); (-21, -15, -9, -1, -1, 7, 13, 19)]$

(iii) Scalar Multiplication

Case-(i): If $k > 0$. Let $k = 2$

$kA = [(14, 18, 22, 26, 30, 34, 38, 42); (8, 14, 20, 26, 30, 36, 42, 48)]$

Case-(ii): If $k < 0$. Let $k = -2$

$-kA = [(-42, -38, -34, -30, -26, -22, -18, -14); (-48, -42, -36, -30, -26, -20, -14, -8)]$

(iv) Additive Image

$-A = [(-21, -19, -17, -15, -13, -11, -9, -7); (-24, -21, -18, -15, -13, -10, -7, -4)]$

6. CONCLUSION

In this paper we have introduced a new type of fuzzy number called SOIFN. The four arithmetic operations of SOIFN are obtained using (α, β) cuts. The SOIFN are used to solve fuzzy optimization problems and industrial applications concerning decision making in an uncertain environment.

REFERENCES

1. K.T. Atanassov, "Intuitionistic Fuzzy Sets Theory and Applications", Physics - Verlag, Heidelberg, New York, 1999.
2. K.T. Atanassov, "Intuitionistic Fuzzy Sets", Fuzzy Sets and Systems, 1986, 20, 87-96.
3. K.T. Atanassov, "More on Intuitionistic Fuzzy Sets", Fuzzy Sets and Systems, 1989, 33, 37-46.
4. R.E. Bellman and L.A. Zadeh, "Decision making in a fuzzy environment", Management Science. 1970, 17, 141-164
5. H. Bustine and P. Burillo, "Vague sets are Intuitionistic fuzzy sets", Fuzzy Sets and Systems, 1996, 79, 403-405.
6. Dhanalakshmi .V, Felbin .C Kennedy, "Ranking algorithm for symmetric octagonal Fuzzy Numbers", Journal of computer and mathematical sciences, 2014, 5, 3, 310-318.
7. W.L. Gau and D.J. Buehrer, "Vague sets", IEEE Transactions on Systems, Man and Cybernetics, 1993, 23, 2, 610-614.
8. G. Menaka, "Ranking of octagonal intuitionistic fuzzy numbers", IOSR journal of mathematics, 2017, 13, 63-71.

9. A. Mohammed Shapique “Arithmetic operations on Heptagonal Fuzzy Numbers”, Asian Research Journal of Mathematics, 2017, 2, 5, 1-25.
10. R. Parvathi, C. Malathi , “Arithmetic Operations on Symmetric Trapezoidal Intuitionistic Fuzzy Numbers”, International journal of soft computing and engineering, 2012, 2, 2, 168-173.
11. K. Ponnivalavan and T. Pathinathan, “Intuitionistic Pentagonal Fuzzy Numbers”, ARPN Journal of Engineering and Applied Sciences, 2015, 10,12, 5446- 5450.
12. P. Rajarajeswari, G.Menaka, “A new approach for ranking of octagonal intuitionistic fuzzy numbers”, International journal of fuzzy logic systems, 2017, 7, 2, 1-11.
13. G. Sahaya sudha, K.R. Vijayalakshmi, “A value and ambiguity based ranking of a symmetric hexagonal intuitionistic fuzzy numbers in decision making”, Advances in fuzzy mathematics, 2017, 12, 4, 867-879.
14. A. Sahaya Sudha and R.Gokilamani, “An Arithmetic Operations on Hexadecagonal Fuzzy Numbers”, International Journal of fuzzy logic systems, 2017, 7, 1, 17-26.
15. L.A. Zadeh, Fuzzy sets, Information and computation, 1965, 8, 338-353.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]