

RAINBOW TOTAL CONNECTION NUMBER OF SOME WHEEL RELATED GRAPHS

JONAH GAY V. PEDRAZA\*<sup>1</sup> AND MICHAEL P. BALDADO JR.<sup>2</sup>

<sup>1</sup>College of Education, Samar State University, Catbalogan Samar, Philippines.

<sup>2</sup>College of Art and Sciences, Negros Oriental State University, Dumaguete City, Philippines.

(Received On: 30-12-18; Revised & Accepted On: 07-02-19)

ABSTRACT

A path  $P$  connecting two vertices  $u$  and  $v$  in a totally colored graph  $G$  is called a rainbow total-path between  $u$  and  $v$  if all elements in  $V(P) \cup E(P)$ , except for  $u$  and  $v$ , are assigned distinct colors. A total-colored graph is rainbow total-connected if it has a rainbow total-path between every two vertices. The rainbow total-connection number of a graph  $G$  is the minimum colors such that  $G$  is rainbow total-connected. In this paper, we gave the rainbow total-connection number of sunflower graph and lotus inside circle graph.

**Keywords:** total-colored graph; rainbow total-connection number; sunflower graph; lotus inside circle.

I. INTRODUCTION

Chartrand *et al.* [2008] introduced the concept rainbow coloring. They determined rainbow connection number of the cycle, path, tree and wheel graphs. Since then many are studying the concept. Please see Li *et al.* [2013] and Sun *et al.* [2012]. Li *et al.* [2013] studied the rainbow connection numbers of line graphs in the light of particular properties of line graphs and gave two sharp upper bounds for rainbow connection number of a line graph. While, Sun *et al.* [2012] investigated the rainbow connection number of the line graph, middle graph and total graph of a connected triangle-free graph and obtained three (near) sharp upper bounds in terms of the number of vertex-disjoint cycles of the original graph. Continuing the study of rainbow coloring, Uchizawa *et al.* [2011] introduced and studied the rainbow total-connection number of graphs. Later, Sun [2013, 2015] also studied rainbow total-connection number. They characterized the rainbow total-connection number of trees, and gave the rainbow total-connection number of cycles, path and wheels. In this paper, we gave the rainbow total-connection number of some wheel related graphs. In particular, we gave the rainbow total-connection number of sunflower graphs, lotus inside circle and helms.

A graph  $G$  is an ordered pair  $(V, E)$  where  $V$  is a non-empty finite set and  $E$  is a family of two element subsets of  $V$ . The elements of  $V$  are called *vertices* and the elements of  $E$  are called *edges*. If  $\{u, v\}$  is an edge, then we say that vertices  $u$  and  $v$  are *adjacent*, and that  $u$  and  $v$  are *incident* to  $\{u, v\}$ . We write edge  $\{u, v\}$  concisely as  $uv$ . The path  $P_n = (v_1, v_2, \dots, v_n)$  is the graph with vertices  $v_1, v_2, \dots, v_n$  and edges  $v_1v_2, v_2v_3, \dots, v_{n-1}v_n$ . The cycle  $C_n = [v_1, v_2, \dots, v_n]$  is the graph with distinct vertices  $v_1, v_2, \dots, v_n$  and edges  $v_1v_2, v_2v_3, \dots, v_{n-1}v_n, v_nv_1$ . A *complete* graph  $K_n$  is the graph with  $n$  vertices and any two vertices is connected by an edge. The *complement* of a graph  $G$ , denoted by  $\overline{G}$ , is the graph with the same vertices as  $G$  and two vertices in  $\overline{G}$  are adjacent if they are not adjacent in  $G$ .

A total coloring of a graph  $G = (V, E)$  is a function  $f$  from  $V \cup E$  to a set  $C$  whose elements are called *colors*. In this case, we say that  $G(f)$  is totally colored. A path  $P$  connecting two vertices  $u$  and  $v$  in a totally colored graph  $G$  is called a rainbow total-path between  $u$  and  $v$  if all the elements in  $[V(P) \cup E(P)] \setminus \{u, v\}$  are assigned distinct colors. The total-colored graph is *rainbow total-connected* if it has a rainbow total-path in between every two vertices. The *rainbow total-connection number* of a graph  $G$  is the minimum colors such that  $G$  is rainbow total-connected.

**Corresponding Author: Jonah Gay V. Pedraza\*<sup>1</sup>,**

**<sup>1</sup>College of Education, Samar State University, Catbalogan Samar, Philippines.**

The following classes of graphs are found Ponraj *et al.* [2015]. The *graph lotus inside circle*, denoted by  $LC_n$ , is the graph of order  $2n+1$  obtained by joining each vertex  $u_i$  of the star  $K_{1,n} = (\{u\}, \emptyset) + (\{u_1, u_2, \dots, u_n\}, \emptyset)$  to vertices  $w_i$  and  $w_{i+1(\text{mod } n)}$  of the cycle  $C_n = [w_1, w_2, \dots, w_n]$ . The *helm*  $H_n$  of order  $2n+1$  is the graph obtained from  $W_n = (\{u\}, \emptyset) + [w_1, w_2, \dots, w_n]$  by attaching pendant edges  $v_i w_i$  for every  $i = 1, 2, \dots, n$ . The *sunflower* graph  $SF_n$  of order  $2n+1$  obtained by adding vertices  $w_i$  joined by edges to vertices  $v_i$  and  $v_{i+1(\text{mod } n)}$  of the  $W_n = (\{v\}, \emptyset) + [v_1, v_2, \dots, v_n]$  for every  $i = 1, 2, \dots, n$ .

Hereafter, please refer to Yellen *et al.* [2000] for concepts that are used but were not discussed in this paper.

In this study, we determined the rainbow total-connection number of lotus inside circle and sunflower graphs.

## II. RESULTS

This section presents the results of this study.

### A. Total Rainbow Connection Number of Lotus Inside a Circle

This subsection gives the total rainbow connection number of lotus inside a circle graph. Remark 1 states that the rainbow connection number of a graph  $G$  is greater than or equal twice its diameter.

**Remark 1:** Let  $G = (V, E)$  be a graph with diameter  $d$ . Then  $rtc(G) \geq 2d - 1$ .

To see this, let  $u, v \in V$  such that the distance in between  $u$  and  $v$ ,  $d(u, v)$ , is equal to  $d$ . Note that any rainbow path connecting  $u$  and  $v$  requires  $2d - 1$  colors. Hence,  $rtc(G) = 2d - 1$ .

**Theorem 2.2:** Let  $C_n = [w_1, w_2, \dots, w_n]$  be a cycle of order  $n$ , and  $K_{1,n} = (\{u_0\}, \emptyset) + (\{u_1, u_2, \dots, u_n\}, \emptyset)$  be a star of order  $n+1$ . Let  $LC_n$  be the graph lotus inside circle obtained by joining each vertex  $u_i$  to vertices  $w_i$  and  $w_{i+1(\text{mod } n)}$ . Then

$$rtc(LC_n) = \begin{cases} 4 & , \text{ if } n = 3, 4 \\ 6 & , \text{ if } n = 5, 6 \\ 7 & , \text{ if } n \geq 7 . \end{cases}$$

**Proof:**

For  $n = 3$ , define  $f_3 : V(LC_3) \cup E(LC_3) \rightarrow \{1, 2, 3, 4\}$  as follows:

**Table-1:** Images of the elements of  $V(LC_3) \cup E(LC_3)$

$v$	$f(v)$	$e$	$f(e)$	$e$	$f(e)$
$u_0$	1	$u_0 u_1$	4	$u_2 w_3$	4
$u_1$	3	$u_0 u_2$	4	$u_3 w_3$	4
$u_2$	3	$w_3 w_1$	2	$u_3 w_1$	2
$u_3$	3	$u_0 u_3$	2	$w_1 w_2$	2
$w_1$	1	$u_1 w_1$	2	$w_2 w_3$	2
$w_2$	1	$u_1 w_2$	4		
$w_3$	1	$u_2 w_2$	2		

Then  $f_3$  is a total rainbow 4-coloring of  $LC_3$ . Hence,  $rtc(LC_3) \leq 4$ . Since the diameter of  $LC_3$  is equal to 2, by Remark 1, we must have,  $rtc(LC_3) = 4$ . For  $n = 4$ , define  $f_4 : V(LC_4) \cup E(LC_4) \rightarrow \{1, 2, 3, 4\}$  as follows:

**Table-2:** Images of the elements of  $V(LC_4) \cup E(LC_4)$

$v$	$f(v)$	$e$	$f(e)$	$e$	$f(e)$
$u_0$	3	$u_0u_1$	2	$u_3w_4$	4
$u_1$	3	$u_0u_2$	2	$u_4w_4$	3
$u_2$	4	$u_0u_3$	4	$u_4w_1$	3
$u_3$	3	$u_0u_4$	1	$w_1w_2$	4
$u_4$	4	$u_1w_1$	2	$w_2w_3$	3
$w_1$	1	$u_1w_2$	4	$w_3w_4$	4
$w_2$	1	$u_2w_2$	2	$w_4w_1$	3
$w_3$	1	$u_2w_3$	3		
$w_4$	1	$u_3w_3$	2		

Then  $f_4$  is a total rainbow 4-coloring of  $LC_4$ . Hence,  $rtc(LC_3) \leq 4$ . Since the diameter of  $LC_4$  is equal to 2, by Remark 1, we must have,  $rtc(LC_4) = 4$ . For  $n = 5$ , define  $f_5 : V(LC_5) \cup E(LC_5) \rightarrow \{1, 2, 3, 4, 5, 6\}$  as follows:

**Table-3:** Images of the elements of  $V(LC_5) \cup E(LC_5)$

$v$	$f(v)$	$e$	$f(e)$	$e$	$f(e)$
$u_0$	6	$u_0u_1$	1	$u_4w_4$	1
$u_1$	5	$u_0u_2$	1	$u_4w_5$	3
$u_2$	5	$u_0u_3$	2	$u_5w_5$	1
$u_3$	6	$u_0u_4$	2	$u_5w_1$	2
$u_4$	5	$u_0u_5$	3	$w_1w_2$	2
$u_5$	5	$u_1w_1$	3	$w_2w_3$	1
$w_1$	6	$u_1w_2$	2	$w_3w_4$	3
$w_2$	6	$u_2w_2$	3	$w_4w_5$	2
$w_3$	6	$u_2w_3$	2	$w_5w_1$	1
$w_4$	4	$u_3w_3$	3		
$w_5$	4	$u_3w_4$	3		

Then  $f_5$  is a total rainbow 6-coloring of  $LC_5$ . Hence,  $rtc(LC_5) \leq 6$ . Since the diameter of  $LC_5$  is equal to 3, by Remark 1, we must have,  $rtc(LC_5) = 6$ . For  $n = 6$ , define  $f_6 : V(LC_6) \cup E(LC_6) \rightarrow \{1, 2, 3, 4, 5, 6\}$  as follows:

**Table-4:** Images of the elements of  $V(LC_6) \cup E(LC_6)$

$v$	$f(v)$	$e$	$f(e)$	$e$	$f(e)$
$u_0$	4	$u_0u_1$	2	$u_4w_5$	1
$u_1$	4	$u_0u_2$	2	$u_5w_5$	3
$u_2$	6	$u_0u_3$	3	$u_5w_6$	3
$u_3$	5	$u_0u_4$	3	$u_6w_6$	2
$u_4$	4	$u_0u_5$	1	$u_6w_1$	2
$u_5$	6	$u_0u_6$	1	$w_1w_2$	1
$u_6$	5	$u_1w_1$	1	$w_2w_3$	3
$w_1$	5	$u_1w_2$	1	$w_3w_4$	2
$w_2$	4	$u_2w_2$	3	$w_4w_5$	1
$w_3$	6	$u_2w_3$	3	$w_5w_6$	3
$w_4$	5	$u_3w_3$	2	$w_6w_1$	2
$w_5$	4	$u_3w_4$	2		
$w_6$	6	$u_4w_4$	1		

Then  $f_6$  is a total rainbow 6-coloring of  $LC_6$ . Hence,  $rtc(LC_6) \leq 6$ . Since the diameter of  $LC_6$  is equal to 3, by Remark 1, we must have,  $rtc(LC_6) = 6$ . For  $n = 7$ , define  $f_7 : V(LC_7) \cup E(LC_7) \rightarrow \{1, 2, \dots, 8\}$  as follows:

**Table-5:** Images of the elements of  $V(LC_7) \cup E(LC_7)$

$v$	$f(v)$	$e$	$f(e)$	$e$	$f(e)$
$u_0$	5	$u_0u_1$	1	$u_5w_5$	4
$u_1$	8	$u_0u_2$	2	$u_5w_6$	4
$u_2$	7	$u_0u_3$	1	$u_6w_6$	3
$u_3$	8	$u_0u_4$	2	$u_6w_7$	3
$u_4$	7	$u_0u_5$	1	$u_7w_7$	4
$u_5$	8	$u_0u_6$	2	$u_7w_1$	4
$u_6$	7	$u_0u_7$	3	$w_1w_2$	1
$u_7$	8	$u_1w_1$	4	$w_2w_3$	1
$w_1$	8	$u_1w_2$	4	$w_3w_4$	1
$w_2$	8	$u_2w_2$	3	$w_4w_5$	1
$w_3$	7	$u_2w_3$	3	$w_5w_6$	1
$w_4$	8	$u_3w_3$	4	$w_6w_7$	1
$w_5$	7	$u_3w_4$	4	$w_7w_1$	1
$w_6$	8	$u_4w_4$	3		
$w_7$	7	$u_4w_5$	3		

Then  $f_7$  is a total rainbow 7-coloring of  $LC_7$ . It can be shown that a total rainbow coloring of  $LC_7$  can not have a fewer than 7 colors. Hence,  $rtc(LC_7) = 7$ .

For  $n \geq 8$  and  $n$  is even, Then we define  $f_n : V(LC_n) \cup E(LC_n) \rightarrow \{1, 2, \dots, 7\}$  as follows:

$$f_n(x) = \begin{cases} 1 & , \text{ if } x = w_iw_{i+1(\text{mod } n)} \text{ or } x = u_0u_i \text{ with } i \text{ odd} \\ 2 & , \text{ if } x = u_0u_i \text{ with } i \text{ even} \\ 3 & , \text{ if } x = u_iw_i \text{ or } x = u_iw_{i+1(\text{mod } n)}, \text{ with } i \text{ odd} \\ 4 & , \text{ if } x = u_iw_i \text{ or } x = u_iw_{i+1(\text{mod } n)}, \text{ with } i \text{ even} \\ 5 & , \text{ if } x = w_i \text{ or } x = u_{i+1(\text{mod } n)}, \text{ with } i \text{ odd} \\ 6 & , \text{ if } x = u_0 \\ 7 & , \text{ if } x = w_i \text{ or } x = u_{i+1}, \text{ with } i \text{ even} \end{cases}$$

Let  $w, v \in V(LC_n)$  and consider the following cases:

**Case-1:**  $\deg(w) = 4$

If  $\deg(w) = 4$ , then  $w = w_i$  for some  $i = 1, 2, \dots, n$ . Consider the following subcases:

**Subcase-1:**  $\deg(v) = 4$

If  $\deg(v) = 4$ , then  $v = w_j$  for some  $j = 1, 2, \dots, n$  with  $j \neq i$ . Note that if  $i$  and  $j$  have the same parity, then  $(w_i, u_i, u_0, u_{j-1}, w_j)$  is a rainbow path connecting  $w$  and  $v$ , and if  $i$  and  $j$  have the different parity, then  $(w_i, u_i, u_0, u_j, w_j)$  is a rainbow path connecting  $w$  and  $v$ .

**Subcase-2:**  $\deg(v) = 3$

If  $\deg(v) = 3$ , then  $v = u_j$  for some  $j = 1, 2, \dots, n$ . If  $i = j - 1$ , then  $(u_i, w_j)$  is a rainbow path connecting  $w$  and  $v$ , and if  $i = j$ , then  $(u_i, w_j)$  is a rainbow path connecting  $w$  and  $v$ . If  $i$  and  $j$  have the same parity with  $j \neq j - 1, j$ , then  $(w_i, u_{i-1}, u_0, u_j)$  is a rainbow path connecting  $w$  and  $v$ , and if  $i$  and  $j$  have the different parity with  $j \neq j - 1, j$ , then  $(w_i, u_i, u_0, u_j)$  is a rainbow path connecting  $w$  and  $v$ .

**Subcase-3:**  $\deg(v) = n$

If  $\deg(v) = n$ , then  $v = u_0$ . Note that  $(w_i, u_i, u_0)$  is a rainbow path connecting  $w$  and  $v$ .

**Case-2:**  $\deg(w) = 3$

If  $\deg(w) = 3$ , then  $w = u_i$  for some  $i = 1, 2, \dots, n$ . Consider the following subcases:

**Subcase-1:**  $\deg(v) = 3$

If  $\deg(v) = 3$ , then  $v = u_j$  for some  $j = 1, 2, \dots, n$ . If  $i$  and  $j$  have the same parity, then  $(u_i, u_0, u_{j+1}, w_{j+1}, u_j)$  is a rainbow path connecting  $w$  and  $v$ , and if  $i$  and  $j$  have the different parity, then  $(u_i, u_0, u_j)$  is a rainbow path connecting  $w$  and  $v$ .

**Subcase-3.:**  $\deg(v) = n$

If  $\deg(v) = n$ , then  $v = u_0$ . Note that  $(u_i, u_0)$  is a rainbow path connecting  $w$  and  $v$ .

Hence,  $f_n$  is a total rainbow 7-coloring of  $LC_n$ . Hence,  $rtc(LC_n) \leq 7$ . Since the diameter of  $LC_n$  is equal to 4, by

Remark 1, we must have,  $rtc(LC_n) = 7$  if  $n \geq 8$  and  $n$  is even.

For  $n \geq 9$  and  $n$  is odd, Then we define  $f_n : V(LC_n) \cup E(LC_n) \rightarrow \{1, 2, \dots, 7\}$  as follows:

$$f_n(x) = \begin{cases} 1 & , \text{ if } x = w_i w_{i+1(\text{mod } n)} \text{ or } x = u_0 u_i \text{ with } i \text{ odd} \\ 2 & , \text{ if } x = u_0 u_i \text{ with } i \text{ even} \\ 3 & , \text{ if } x = u_i w_i \text{ or } x = u_i w_{i+1(\text{mod } n)}, \text{ with } i \text{ odd} \\ 4 & , \text{ if } x = u_i w_i \text{ or } x = u_i w_{i+1(\text{mod } n)}, \text{ with } i \text{ even} \\ 5 & , \text{ if } x = w_i \text{ or } x = u_{i+1(\text{mod } n)}, \text{ with } i \text{ odd} \\ 6 & , \text{ if } x = u_0 \\ 7 & , \text{ if } x = w_i \text{ or } x = u_{i+1}, \text{ with } i \text{ even} \end{cases}$$

It can also be shown that  $f_n$  is a total rainbow 7-coloring of  $LC_n$ . Hence,  $rtc(LC_n) \leq 7$ . Since the diameter of  $LC_n$  is equal to 4, by Remark 1, we must have,  $rtc(LC_n) = 7$  if  $n \geq 9$  and  $n$  is odd.

## B. Total Rainbow Connection Number of Sunflower Graphs

This subsection gives the total rainbow connection number of sunflower graph. Theorem 3 is due to Sun (2013).

**Theorem 3:** If  $G$  is a connected graph, then

1.  $rtc(G) = 1$  if and only if  $G$  is a complete graph;
2.  $rtc(G) \neq 2$ ;
3.  $rtc(G) = m + n_2$  if and only if  $G$  is a tree.

**Theorem 4:** Let  $W_n = (\{u_0\}, \emptyset) + [u_1, u_2, \dots, u_n]$  be a wheel of order  $n+1$  and  $SF_n$  be the sunflower graph obtained by adding a vertex  $w_i$  joined by an edge to vertices  $u_i$  and  $u_{i+1(\text{mod } n)}$ . Then

$$rtc(SF_n) = \begin{cases} 3 & , \text{ if } n = 3 \\ 6 & , \text{ if } n = 4, 5 \\ 7 & , \text{ if } n \geq 6. \end{cases}$$

**Proof:**

For  $n = 3$ , define  $f_3 : V(SF_3) \cup E(SF_3) \rightarrow \{1, 2, 3\}$  as follows:

**Table-1:** Images of the elements of  $V(SF_3) \cup E(SF_3)$

$v$	$f(v)$	$e$	$f(e)$	$e$	$f(e)$
$u_0$	3	$u_0u_1$	2	$u_3w_2$	2
$u_1$	3	$u_0u_2$	2	$u_3w_3$	1
$u_2$	3	$u_3u_1$	2	$u_1w_3$	2
$u_3$	3	$u_0u_3$	2	$u_1u_2$	2
$w_1$	3	$u_1w_1$	1	$u_2u_3$	2
$w_2$	3	$u_2w_1$	2		
$w_3$	3	$u_2w_2$	1		

Then  $f_3$  is a total rainbow 4-coloring of  $SF_3$ . Hence,  $rtc(SF_3) \leq 3$ . By Theorem 2, we must have,  $rtc(LC_3) = 3$ . For  $n = 4$ , define  $f_4 : V(SF_4) \cup E(SF_4) \rightarrow \{1, 2, 3\}$  as follows:

**Table-2:** Images of the elements of  $V(SF_4) \cup E(SF_4)$

$v$	$f(v)$	$e$	$f(e)$	$e$	$f(e)$
$u_0$	5	$u_0u_1$	1	$u_4w_3$	4
$u_1$	5	$u_0u_2$	1	$u_4w_4$	2
$u_2$	4	$u_0u_3$	2	$u_1w_4$	3
$u_3$	5	$u_0u_4$	3	$u_1u_2$	2
$u_4$	4	$u_1w_1$	1	$u_2u_3$	3
$w_1$	5	$u_2w_1$	2	$u_3u_4$	2
$w_2$	4	$u_2w_2$	1	$u_4u_1$	2
$w_3$	5	$u_3w_2$	3		
$w_4$	4	$u_3w_3$	1		

Then  $f_4$  is a total rainbow 5-coloring of  $SF_4$ . Hence,  $rtc(SF_4) \leq 5$ . Since the diameter of  $SF_4$  is equal to 3, by Remark 1, we must have,  $rtc(SF_4) = 5$ .

For  $n \geq 5$  and  $n$  is even, Then we define  $f_n : V(SF_n) \cup E(SF_n) \rightarrow \{1, 2, \dots, 7\}$  as follows:

$$f_n(x) = \begin{cases} 1 & , \text{ if } x = w_iw_{i+1(\text{mod } n)} \text{ or } x = u_0u_i \text{ with } i \text{ odd} \\ 2 & , \text{ if } x = u_0u_i \text{ with } i \text{ even} \\ 3 & , \text{ if } x = u_iw_i \text{ or } x = u_iw_{i+1(\text{mod } n)}, \text{ with } i \text{ odd} \\ 4 & , \text{ if } x = u_iw_i \text{ or } x = u_iw_{i+1(\text{mod } n)}, \text{ with } i \text{ even} \\ 5 & , \text{ if } x = w_i \text{ or } x = u_{i+1(\text{mod } n)}, \text{ with } i \text{ odd} \\ 6 & , \text{ if } x = u_0 \\ 7 & , \text{ if } x = w_i \text{ or } x = u_{i+1}, \text{ with } i \text{ even} \end{cases}$$

For  $n \geq 9$  and  $n$  is odd, Then we define  $f_n : V(LC_n) \cup E(LC_n) \rightarrow \{1, 2, \dots, 7\}$  as follows:

$$f_n(x) = \begin{cases} 1 & , \text{ if } x = w_iw_{i+1(\text{mod } n)} \text{ or } x = u_0u_i \text{ with } i \text{ odd} \\ 2 & , \text{ if } x = u_0u_i \text{ with } i \text{ even} \\ 3 & , \text{ if } x = u_iw_i \text{ or } x = u_iw_{i+1(\text{mod } n)}, \text{ with } i \text{ odd} \\ 4 & , \text{ if } x = u_iw_i \text{ or } x = u_iw_{i+1(\text{mod } n)}, \text{ with } i \text{ even} \\ 5 & , \text{ if } x = w_i \text{ or } x = u_{i+1(\text{mod } n)}, \text{ with } i \text{ odd} \\ 6 & , \text{ if } x = u_0 \\ 7 & , \text{ if } x = w_i \text{ or } x = u_{i+1}, \text{ with } i \text{ even} \end{cases}$$

It can be shown that  $f_n$  is a total rainbow 7-coloring of  $SF_n$  and a total rainbow coloring of  $SF_7$  cannot have a fewer than 7 colors. Hence, we must have,  $rtc(SF_n) = 7$  if  $n \geq 5$ .

#### IV. RECOMMENDATION

We recommend the rainbow total connection number of other cycle related graphs mentioned in Ponraj *et al.* (2015) be determined also.

#### V. ACKNOWLEDGEMENT

The researcher would like to thank Samar State University for supporting this paper.

#### VI. REFERENCES

1. Yuefang S., On rainbow total-coloring of a graph, *Discrete Applied Mathematics* 5 (3) (2015) 260-276. <http://dx.doi.org/10.1016/j.dam.2015.05.012>.
2. Chakraborty S., Fischer E., Matsliah A. & Yuster R. (2009), Hardness and algorithms for rainbow connectivity, 26th International Symposium on Theoretical Aspects of Computer Science STACS 2009 (2009), 243-254.
3. Chartrand G., Johns G.I., Mc Keon K.A. & Zhang P. (2008), Rainbow connection in graphs. *Math. Bohem.* 133 (2008) 85-98.
4. Li X., Shi Y. & Sun Y. (2013), Rainbow connection in graphs – A survey, *Graphs Combin.* 29 (2013) 1-38.
5. Uchizawa K., Aoki T., Ito T., Suzuki A. & Zhou X. (2011), On the Rainbow Connectivity of Graphs: Complexity and FPT Algorithms, in: LNCS, vol. 6842.
6. Sun Y. (2013), On two variants of rainbow connection, *WSEAS Trans. Math.* 12 (3) (2013) 266-276.
7. J. Gross, J. Yellen, *Handbook of Graph Theory*, CRC Press LLC, N. W., Corporate Blvd, Boca Raton, Florida.
8. R. Ponraj, S. Sathish Narayanan, R. Kala, Radio mean labeling of a graph, *AKCE International Journal of Graphs and Combinatorics*, 12 (2015) 224-228.

**Source of support: Nil, Conflict of interest: None Declared.**

***[Copy right © 2019. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]***