

DECOMPOSITION OF RECURRENT AND H-PROJECTIVE CURVATURE TENSOR FIELDS
 IN A KAEHLERIAN SPACE OF FIRST ORDER

TRISHNA DEVI¹, U. S. NEGI*² AND MAHENDER SINGH POONIA³

¹Research Scholar,
 Department of Mathematics,
 Shri Jagdish Prasad Jhabarmal Tibrewala University,
 Vidyanagari, Jhunjhunu, Rajasthan-333001, India.

²Department of Mathematics,
 H.N.B. Garhwal University, SRT Campus Badshahi Thaul,
 Tehri Garhwal- 249 199, Uttarakhand, India.

³Associate Professor,
 Department of Mathematics,
 Shri Jagdish Prasad Jhabarmal Tibrewala University,
 Vidyanagari, Jhunjhunu, Rajasthan-333001, India.

(Received On: 30-01-19; Revised & Accepted On: 11-03-19)

ABSTRACT

In this paper, we have studied decomposition of recurrent and H-Projective curvature tensor fields in a Kaehlerian space of first order by considering the decomposition of curvature tensor field in terms of a non- zero vector and tensor field. Also, several theorems have been derived.

Key Words: Kaehlerian space, Projective, Recurrent, Curvature tensor.

2010 MSC: 32C15, 46A13, 53B35, 53C55.

1. INTRODUCTION

A 2n-dimensional Kaehlerian space K_n^c is a Riemannian space which admits a tensor field an almost complex structure F_i^h satisfying the relation (Yano 1965).

$$F_j^i F_i^h = -A_j^h, \tag{1.1}$$

$$F_s^t F_i^s g_{ts} = g_{ji} \quad \text{and} \quad F_i^h{}_{,j} = 0 \tag{1.2}$$

$$F_{ji} = -F_{ij} \tag{1.3}$$

$$F_{ji} = F_j^t g_{ti} \tag{1.4}$$

And finally has the property that the skew-symmetric tensor F_{ih} is a killing tensor, then

$$F_{ih,j} + F_{jh,i} = 0 \tag{1.5}$$

$$F_i^h{}_{,j} + F_j^h{}_{,i} = 0 \tag{1.6}$$

$$F_i = -F_{i,j}^j \tag{1.7}$$

Where the comma (,) followed by an index denotes the operator of covariant differentiation with respect to the metric tensor g_{ji} of the Riemannian space.

The Riemannian curvature tensor field is defined by

$$R_{ijk}^h = \partial_i \{^h_{jk}\} - \partial_j \{^h_{ik}\} + \{^h_{ia}\} \{^a_{jk}\} - \{^h_{ja}\} \{^a_{ik}\} \tag{1.8}$$

Corresponding Author: U. S. Negi*²,
²Department of Mathematics, H.N.B. Garhwal University,
 SRT Campus Badshahi Thaul, Tehri Garhwal- 249 199, Uttarakhand, India.

The Ricci tensor and scalar curvature are respectively given by

$$R_{ij} = R_{aj}^a \quad \text{and} \quad R = g^{ij} R_{ij} \tag{1.9}$$

It is well known that these tensors satisfy the following identities

$$R_{ijk}^a = R_{jk,i}^a - R_{ik,j}^a \tag{1.10}$$

$$R_{,i} = 2 R_{i,a}^a \tag{1.11}$$

$$F_i^a R_{aj} = - R_{ia} F_j^a \tag{1.12}$$

$$F_i^a R_a^j = R_i^a F_a^j \tag{1.13}$$

The holomorphically projective curvature tensor P_{ijk}^h is defined by (Sinha, 1973)

$$P_{ijk}^h = R_{ijk}^h + \frac{1}{(n+2)} (R_{ik} \delta_j^h - R_{jk} \delta_i^h + S_{ik} F_j^h - S_{jk} F_i^h + 2 S_{ij} F_k^h) \tag{1.14}$$

Where $S_{ij} = F_i^a R_{aj}$

The Bianchi identities are given by (Takano,1967).

$$R_{ijk}^h + R_{jki}^h + R_{kij}^h = 0 \tag{1.15}$$

$$R_{ijk,a}^h + R_{ika,j}^h + R_{iaj,k}^h = 0 \tag{1.16}$$

The Commutative formulae for the curvature tensor fields are given as follows:

$$N_{,jk}^i - N_{kj}^i = N^a R_{ajk}^i \tag{1.17}$$

$$N_{i,ml}^h - N_{i,lm}^h = N_i^a R_{aml}^h - N_a^h R_{iml}^a \tag{1.18}$$

Definition (1.1): A Kaehlerian space is said to be recurrent, if we have (Singh 1971)

$$R_{ijk,a}^h = \lambda_a R_{ijk}^h, \tag{1.19}$$

for some non-zero recurrence vector λ_a , and is called semi-recurrent (or Ricci-recurrent), if it satisfies

$$R_{ij,a} = \lambda_a R_{ij}, \tag{1.20}$$

Multiplying the above equation by g^{ij} , we get

$$R_{,a} = \lambda_a R. \tag{1.21}$$

Remark (1.1): From (1.2) it follows that every Kaehlerian recurrent space is Kaehlerian Ricci-recurrent space but the converse is not necessarily true.

2. DECOMPOSITION OF RECURRENT CURVATURE TENSOR FIELDS IN A KAEHLERIAN SPACE OF FIRST ORDER.

We Consider the decomposition of recurrent curvature tensor field R_{ijk}^h in the following form:

$$R_{ijk}^h = X^h Y_{ij,k} \tag{2.1}$$

Where two vectors X^h and a tensor field $Y_{ij,k}$ such that

$$\lambda_h X^h = 1 \tag{2.2}$$

Theorem 2.1: Under the decomposition (2.1), the Bianchi identity for R_{ijk}^h take the forms

$$Y_{ij,k} + Y_{jk,i} + Y_{ki,j} = 0 \tag{2.3}$$

and $\lambda_a Y_{ij,k} + \lambda_j Y_{ik,a} + \lambda_k Y_{ia,j} = 0$ (2.4)

Proof: From (1.15) and (2.1), we have

$$X^h Y_{ij,k} + X^h Y_{jk,i} + X^h Y_{ki,j} = 0 \tag{2.5}$$

Multiplying (2.5) by λ_h and using (2.2), we obtain the required result (2.3)

Again, using (1.16), (1.19) and (2.1) we have

$$X^h (\lambda_a Y_{ij,k} + \lambda_j Y_{ik,a} + \lambda_k Y_{ia,j}) = 0 \tag{2.6}$$

Multiplying (2.6) by λ_h and using (2.2), we get the required result (2.4).

Theorem 2.2: Under the decomposition (2.1), the tensor fields R_{ijk}^h , R_{ij} and $Y_{ij,k}$ satisfy relation

$$\lambda_a R_{ijk}^a = \lambda_i R_{jk} - \lambda_j R_{ik} = Y_{ij,k} \quad (2.7)$$

Proof: With the help of (1.10), (1.19) and (1.20), we have

$$\lambda_a R_{ijk}^a = \lambda_i R_{jk} - \lambda_j R_{ik} \quad (2.8)$$

Multiplying (2.1) by λ_h and using relation (2.2), we have

$$\lambda_h R_{ijk}^h = Y_{ij,k} \quad (2.9)$$

From (2.8) and (2.9), we get the required relation (2.7).

Theorem 2.3: Under the decomposition (2.1), the quantities λ_a and X^h behave like the recurrent vectors. The recurrent form of these quantities are given by

$$\lambda_{a,m} = \mu_m \lambda_a \quad (2.10)$$

and $X_{,m}^h = -\mu_m X^h \quad (2.11)$

Proof: Differentiating (2.8) covariantly w.r.t. x^m and using (2.1) and (2.7), we have

$$\lambda_{a,m} X^a Y_{ij,k} = \lambda_{i,m} R_{jk} - \lambda_{j,m} R_{ik} \quad (2.12)$$

Multiplying (2.13) by λ_a and using (2.1) and (2.9), we have

$$\lambda_{a,m} (\lambda_i R_{jk} - \lambda_j R_{ik}) = \lambda_a (\lambda_{i,m} R_{jk} - \lambda_{j,m} R_{ik}) \quad (2.13)$$

Now multiplying (2.13) by λ_h , we have

$$\lambda_{a,m} (\lambda_i R_{jk} - \lambda_j R_{ik}) \lambda_a = \lambda_a \lambda_h (\lambda_{i,m} R_{jk} - \lambda_{j,m} R_{ik}) \quad (2.14)$$

Since the expression on the right hand side of the above equation is symmetric in a and h, therefore

$$\lambda_{a,m} \lambda_h = \lambda_{h,m} \lambda_a \quad (2.15)$$

Provided $\lambda_i R_{jk} - \lambda_j R_{ik} \neq 0$

The vector field λ_a being a non-zero, we can choose a proportional vector field μ_m such that

$$\lambda_{a,m} = \mu_m \lambda_a \quad (2.16)$$

Further, differentiating (2.2) covariantly w.r.t. x^m and using (2.16), we have

$$X_{,m}^h = -\mu_m X^h$$

Theorem 2.4: Under the decomposition (2.1), the vector X^h and the tensor $Y_{ij,k}$ satisfy the relation

$$(\lambda_m + \mu_m) Y_{ij,k} = Y_{ij,km} \quad (2.17)$$

Proof: Differentiating (2.1) covariantly w.r.t. x^m and using (1.19), (2.1) and (2.11), we get the required result (2.17).

3. DECOMPOSITION OF H-PROJECTIVE CURVATURE TENSOR FIELDS IN A KAEHLERIAN SPACE OF FIRST ORDER.

Theorem 2.5: Under the decomposition (2.1), the curvature tensor and holomorphically projective curvature tensor are equal iff

$$(Y_{ik,m} \delta_j^h - Y_{jk,m} \delta_i^h) + Y_{ik,m} (F_i^l F_j^h - F_j^l F_i^h) + 2 F_i^l Y_{ij,m} F_k^h = 0 \quad (2.18)$$

Proof: The equation (1.14) may be written in the form

$$P_{ijk}^h = R_{ijk}^h + D_{ijk}^h \quad (2.19)$$

Where

$$D_{ijk}^h = \frac{1}{(n+2)} (R_{ik} \delta_j^h - R_{jk} \delta_i^h - S_{ik} F_j^h - S_{jk} F_i^h + 2S_{ij} F_k^h) \quad (2.20)$$

Contracting indices h and k in (2.1), we have

$$R_{ij} = X^k Y_{ij,k} \quad (2.21)$$

In view of (2.21), we have

$$S_{ij} = F_i^l X^m Y_{ij,m} \quad (2.22)$$

Making use of (2.21) and (2.22) in (2.20), we obtain

$$D_{ijk}^h = \frac{1}{(n+2)} [X^m (Y_{ij,m} \delta_j^h - Y_{jk,m} \delta_i^h) + X^m Y_{lk,m} (F_i^l F_j^h - F_j^l F_i^h) + 2 F_i^l Y_{ij,m} F_k^h] \quad (2.23)$$

From equation (2.19), it is clear that

$$P_{ijk}^h = R_{ijk}^h \text{ iff } D_{ijk}^h = 0, \text{ which in view of (2.23) becomes}$$

$$X_m \{ (Y_{ij,m} \delta_j^h - Y_{jk,m} \delta_i^h) + X^m Y_{lk,m} (F_i^l F_j^h - F_j^l F_i^h) \} + 2 F_i^l Y_{ij,m} F_k^h = 0 \quad (2.24)$$

Multiplying (2.24) by λ_m and using (2.2), we obtain the required result (2.18).

Theorem 2.6: Under the decomposition (2.1), the scalar curvature R, satisfy the relation

$$\lambda_k R = g^{ij} Y_{ij,k} \quad (2.25)$$

Proof: Contracting indices h and k in (2.1), we get

$$R_{ij} = X^k Y_{ij,k} \quad (2.26)$$

Multiplying (2.26) by g^{ij} on both sides, we have

$$g^{ij} R_{ij} = g^{ij} X^k Y_{ij,k} \text{ or } R = g^{ij} X^k Y_{ij,k} \quad (2.27)$$

Now, multiplying (2.27) by λ_k , then using (2.2), we have

$$\lambda_k R = g^{ij} Y_{ij,k} \text{ or } R_{,K} = g^{ij} Y_{ij,k}$$

Which completes the proof of the theorem.

REFERENCES

1. Walker A.G. (1950). On Ruse's space of recurrent curvature. Proc. Lond. Math. Soc., 2(52), 36-64.
2. Yano K. (1965). Differential Geometry on Complex and Almost Complex Spaces. Pergamon press.
3. Takano K. (1967). Decomposition of curvature tensor in a recurrent space. Tensor (N.S.), 18(3), 343-347.
4. Sinha B.B. and Singh S.P. (1970). On decomposition of recurrent curvature tensor fields in a Finsler space. Bull. Cal. Math. Soc., 62, 91-96.
5. Singh, S.S. (1971): On Kaehlerian spaces with recurrent Bochner curvature. Acc. Naz. Dei Lincei, Rend, Vol. 51, No. (3, 4), pp. 213-220.
6. Sinha, B.B. (1973): On H-curvature tensors in Kaehler manifold. Kyungpook Math. J., 13, No.2, pp. 185-189.
7. Negi, U.S. and Kailash Gairola (2012): Admitting a conformal transformation group on Kaehlerian recurrent spaces, International Journal of Mathematical Archive-3(4), pp.1584-1589.
8. Negi, U.S. (2018): Geometrical interpretation of an analytic HP-transformations in almost Kaehlerian spaces, International Journal of Mathematical Archive- 9(1), pp.18-22.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2019. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]