

**COMPUTING CERTAIN TOPOLOGICAL INDICES
 FOR LINE GRAPHS OF TADPOLE GRAPHS, WHEEL GRAPH AND LADDER GRAPH**

PRASHANT V. PATIL* AND GIRISH G. YATTINAHALLI**

*Department of Mathematics, Jain College of Engineering, Belagavi, Karnataka, India.

**Department of Mathematics, SKSVMACET, Laxmeshwar, Karnataka, India.

(Received On: 13-09-19; Revised & Accepted On: 25-11-19)

ABSTRACT

Topological indices have been used as a special tools in mathematical chemistry to predict physico-chemical properties of certain chemical compounds. Recently various new topological indices have been conceived by researchers such as Sanskruti index $S(G)$, fifth M_1 and M_2 Zagreb indices, fifth M_1 and M_2 multiplicative Zagreb indices. In this paper, we give explicit formulae for tadpole graph, wheel graph and ladder graph and their corresponding line graphs for the above mentioned topological indices in terms of the order of the original graph G .

1. INTRODUCTION AND PRELIMINARIES

Let G be a simple graph. The order of a graph is $|V(G)|$, its number of vertices denoted by n . The size of a graph is $|E(G)|$, its number of edges denoted by m . The degree of a vertex v , denoted by $d_G(v)$. The subdivision graph $S_1(G)$ is the graph attained from G by replacing each of its edges by a path of length 2. The line graph $L(G)$ of a graph is the graph derived from G in such a way that the edges in G are replaced by vertices in $L(G)$ and two vertices in $L(G)$ are connected whenever the corresponding edges in G are adjacent [4]. For any number d , we define $V_d = \{u \in V(G) \mid s_G(u) = d\}$, in which $s_G(u) = \sum_{v \in N_G(u)} d_G(v)$ and

$$N_G(u) = \{v \in V(G) \mid uv \in E(G)\}.$$

In structural chemistry and biology, molecular structure descriptors are utilized for modeling information of molecules, which are known as topological indices. Many topological indices are introduced to explain the physical and chemical properties of molecules. Topological Index is a numerical value associated with the molecular graph and is obtained from various graph parameters. One of them is degree based topological indices. A large number of such indices depend only on vertex degree of the molecular graph. One of them is the Sanskruti index, proposed by Hosamani [8]. Which is defined as follows:

$$S(G) = \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u) + s_G(v) - 2} \right)^3. \tag{1}$$

Recently, the first neighborhood Zagreb index was introduced and studied by Basavangoud *et.al*, [1]

$$NM_1(G) = \sum_{u \in V(G)} S_G(u)^2. \tag{2}$$

Corresponding Author: Prashant V. Patil*

***Department of Mathematics, Jain College of Engineering, Belagavi, Karnataka, India.**

The fifth M_1 and M_2 Zagreb indices were introduced by Graovac *et.al.*, [2], defined as:

$$M_1 G_5(G) = \sum_{uv \in E(G)} [S_G(u) + S_G(v)]. \tag{3}$$

$$M_2 G_5(G) = \sum_{uv \in E(G)} [S_G(u) \cdot S_G(v)]. \tag{4}$$

The fifth multiplicative M_1 and M_2 Zagreb indices were proposed by Kulli in [9], defined as

$$M_1 G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u) + S_G(v)]. \tag{5}$$

$$M_2 G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u) \cdot S_G(v)]. \tag{6}$$

For more information on topological indices we refer the articles [3,5,6,10].

Now we define some notions of the graph theory. The subdivision graph $S(G)$ is the graph obtained from G by replacing each of its edge by a path of length 2. The line graph $L(G)$ of graph G is the graph whose vertices are the edges of G , two vertices e and f are incident if and only if they have a common end vertex in G . The tadpole graph $T_{n,k}$ is the graph obtained by joining a cycle of n vertices with a path of length k . A ladder graph L_n is obtained by taking cartesian product of two paths $P_n \times P_2$. A wheel graph W_n of order n composed of a vertex, which will be called the hub, adjacent to all vertices of a cycle of order n , i.e $W_n = C_{n-1} + K_1$.

2. TOPOLOGICAL INDICES OF THE TADPOLE GRAPH AND ITS LINE GRAPH

In 2011, Ranjini *et.al.* calculated the explicit expressions for the Shultz index of the subdivision graphs of the tadpole graph, wheel, helm and ladder graphs [13]. They also studied the Zagreb indices of the line graph of tadpole, wheel and ladder graphs with subdivision in [12]. In 2015, Su and Xu calculated the general sum-connectivity index and co-index for the line graph of tadpole, wheel and ladder graphs with subdivision [14]. In the same year (2015) M. F. Nadeem *et.al.* computed the ABC_4 and GA_5 indices for the same graphs [11].




Fig.1. (a) The tadpole graph $T_{n,k}$; (b) the line graph $L(T_{n,k})$ of tadpole graph

Table-1: The edge partition of tadpole graph $T_{n,k}$.

(S_u, S_v) where $uv \in E(G)$	(4,4)	(5,6)	(3,2)
Number of edges	$n + k - 4$	3	1

Table-2: The edge partition of line graph of tadpole graph $T_{n,k}$.

(S_u, S_v) where $uv \in E(G)$	(4,4)	(5,8)	(8,8)	(2,3)
Number of edges	$n+k-6$	3	3	1

Theorem 1:1 Let $G = T_{n,k}$ be the tadpole graph. Then

1. $S(G) = (n+k-4)(18.9572) + 119.1077$
2. $M_1G_5(G) = 8(n+k) + 6$
3. $M_2G_5(G) = 16(n+k) + 32$
4. $M_1G_5 \prod(G) = 1320(n+k) - 5280$
5. $M_2G_5 \prod(G) = 8640(n+k) - 34560$

Proof: Let G be a tadpole graph $T_{n,k}$ with $|V(G) = n+k| = E(G)$. Since G is unicyclic therefore G has same order and size. Now consider the following cases:

Case-1: Consider the Sanskruti index $S(G) = \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u) + s_G(v) - 2} \right)^3$. By using the information in Table 1,

we have

$$\begin{aligned} S(G) &= \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u) + s_G(v) - 2} \right)^3 \\ &= (n+k-4) \left(\frac{4 \times 4}{4+4-2} \right)^3 + 3 \left(\frac{5 \times 6}{5+6-2} \right)^3 + 1 \left(\frac{3 \times 2}{3+2-2} \right)^3 \\ &= (n+k-4)(2.6666)^3 + 3(3.3333)^3 + 1(2)^3 \\ &= (n+k-4)(18.9572) + 119.1077. \end{aligned}$$

Case-2: Consider the fifth M_1 Zagreb Index, $M_1G_5(G) = \sum_{uv \in E(G)} [S_G(u) + S_G(v)]$. Again by using the information in Table 1, we have

$$\begin{aligned} M_1G_5(G) &= \sum_{uv \in E(G)} [S_G(u) + S_G(v)] \\ &= (n+k-4)(4+4) + 3(5+6) + 1(3+2) \\ &= 8(n+k) + 6. \end{aligned}$$

Case-3: Consider the fifth M_2 Zagreb Index, $M_2G_5(G) = \sum_{uv \in E(G)} [S_G(u)S_G(v)]$. Again by using the information in Table 1, we have

$$\begin{aligned} M_2G_5(G) &= \sum_{uv \in E(G)} [S_G(u)S_G(v)] \\ &= (n+k-4)(4 \times 4) + 3(5 \times 6) + 1(3 \times 2) \\ &= 16(n+k) + 32. \end{aligned}$$

Case-4: Consider the The fifth multiplicative M_1 Zagreb index $M_1G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u) + S_G(v)]$. Using Table 1, we have

$$\begin{aligned} M_1G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u) + S_G(v)] \\ &= (n+k-4)(4+4) \times 3(5+6) \times 1(3+2) \\ &= 1320(n+k) - 5280. \end{aligned}$$

Consider the The fifth multiplicative M_2 Zagreb index $M_2G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u)S_G(v)]$. Using Table 1, we have

$$\begin{aligned} M_2G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u)S_G(v)] \\ &= (n+k-4)(4 \times 4) \times 3(5 \times 6) \times 1(3 \times 2) \\ &= 8640(n+k)(34560). \end{aligned}$$

Theorem 2:2 Let $H = L(T_{n,k})$ be the line graph of tadpole graph $T_{n,k}$. Then

1. $S(H) = (18.9615)(n+k-6) + 457.84$
2. $M_1G_5(H) = 8(n+k) + 44$
3. $M_2G_5(H) = 16(n+k) + 222$
4. $M_1G_5 \prod(H) = 74880(n+k-6)$
5. $M_2G_5 \prod(H) = (n+k-6)(2211840)$

Proof: Let G be a tadpole graph $T_{n,k}$ with $|V(G) = n+k| = E(G)$ and let $H = L(T_{n,k})$ be the line graph of tadpole graph. Now consider the following cases:

Case-1: Consider the Sanskruti index $S(G) = \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u) + s_G(v) - 2} \right)^3$. By using the information in Table 2,

we have

$$\begin{aligned} S(H) &= \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u) + s_G(v) - 2} \right)^3 \\ &= (n+k-6) \left(\frac{4 \times 4}{4+4-2} \right)^3 + 3 \left(\frac{5 \times 8}{5+8-2} \right)^3 + 3 \left(\frac{8 \times 8}{8+8-2} \right)^3 + 1 \left(\frac{2 \times 3}{2+3-2} \right)^3 \\ &= (n+k-6) \left(\frac{16}{6} \right)^3 + 3 \left(\frac{40}{11} \right)^3 + 3 \left(\frac{64}{14} \right)^3 + 1 \left(\frac{6}{3} \right)^3 \\ &= (n+k-6)(2.6666)^3 + 3(3.6363)^3 + 3(4.5714)^3 + 1(3)^3 \\ &= (n+k-6)(18.9615) + 457.84. \end{aligned}$$

Case-2: Consider the fifth M_1 Zagreb Index, $M_1G_5(G) = \sum_{uv \in E(G)} [S_G(u) + S_G(v)]$. Again by using the information in Table 2, we have

$$\begin{aligned} M_1G_5(G) &= \sum_{uv \in E(G)} [S_G(u) + S_G(v)] \\ &= (n+k-6)(4+4) + 3(5+8) + 3(8+8) + 1(2+3) \\ &= 8(n+k) + 44. \end{aligned}$$

Case-3: Consider the fifth M_2 Zagreb Index, $M_2G_5(G) = \sum_{uv \in E(G)} [S_G(u)S_G(v)]$. Again by using the information in Table 2, we have

$$\begin{aligned} M_1G_5(G) &= \sum_{uv \in E(G)} [S_G(u)S_G(v)] \\ &= (n+k-6)(4 \times 4) + 3(5 \times 8) + 3(8 \times 8) + 1(2 \times 3) \\ &= 16(n+k) + 222. \end{aligned}$$

Case-4: Consider the The fifth multiplicative M_1 Zagreb index $M_1G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u) + S_G(v)]$. Using

Table 2, we have

$$\begin{aligned} M_1G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u) + S_G(v)] \\ &= (n+k-4)(4+4) \times 3(5+8) \times 3(8+8) \times 1(2+3) \\ &= 74880(n+k-6) \end{aligned}$$

Case-5: Consider the The fifth multiplicative M_2 Zagreb index $M_2G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u)S_G(v)]$. Using

Table 2, we have

$$\begin{aligned} M_2G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u)S_G(v)] \\ &= (n+k-4)(4 \times 4) \times 3(5 \times 8) \times 3(8 \times 8) \times 1(2 \times 3) \\ &= (n+k-6)(2211840). \end{aligned}$$

3. TOPOLOGICAL INDICES OF THE WHEEL GRAPH AND ITS LINE GRAPH




Fig.2. The Wheel graph W_n and its line graph $L(W_n)$

Table-3: The edge partition wheel graph W_n .

(S_u, S_v) where $uv \in E(G)$	$(n+5, n+5)$	$(n+5, 3(n-1))$
Number of edges	$n-1$	$n-1$

Table-4: The edge partition of line graph of wheel graph W_n .

(S_u, S_v) where $uv \in E(G)$	$(2n+8, 2n+8)$	$(2n+8, n(n-2)+8)$	$(n(n-2)+8, n(n-2)+8)$
Number of edges	$n-1$	$2(n-1)$	$\frac{(n-1)(n-2)}{2}$

Theorem 3:3 Let $G = W_n$ be the wheel graph. Then

1. $(G) = (n-1) \left[\frac{(n+5)^6}{(2n+8)^3} + \frac{[(n+5)(3n-3)]^3}{64n^3} \right]$
2. $M_1G_5(G) = 6(n-1)(n+2)$
3. $M_2G_5(G) = (n-1)(n+5)(4n+3)$
4. $M_1G_5 \prod(G) = (n-1)^2(2n+10)(4n+2)$
5. $M_2G_5 \prod(G) = (n-1)^2(n+5)^3(3n-3)$

Proof: Let $G = W_n$ be a wheel graph of order n . Now consider the following cases:

Case-1: Consider the Sanskruti index $S(G) = \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u) + s_G(v) - 2} \right)^3$. By using the information in Table 3,

we have

$$\begin{aligned} S(G) &= \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u) + s_G(v) - 2} \right)^3 \\ &= (n-1) \left(\frac{(n+5)(n+5)}{n+5+n+5-2} \right)^3 + (n-1) \left(\frac{(n+5)(3n-3)}{n+5+3n-3-2} \right)^3 \\ &= (n-1) \left[\frac{(n+5)^6}{(2n+8)^3} + \frac{[(n+5)(3n-3)]^3}{64n^3} \right]. \end{aligned}$$

Case-2: Consider the fifth M_1 Zagreb Index, $M_1G_5(G) = \sum_{uv \in E(G)} [S_G(u) + S_G(v)]$. Again by using the information in Table 3, we have

$$\begin{aligned} M_1G_5(G) &= \sum_{uv \in E(G)} [S_G(u) + S_G(v)] \\ &= (n-1)(n+5+n+5) + (n-1)(n+5+3n-3) \\ &= 6(n-1)(n+2). \end{aligned}$$

Case-3: Consider the fifth M_2 Zagreb Index, $M_2G_5(G) = \sum_{uv \in E(G)} [S_G(u)S_G(v)]$. Again by using the information in Table 3, we have

$$\begin{aligned} M_2G_5(G) &= \sum_{uv \in E(G)} [S_G(u)S_G(v)] \\ &= (n+5)^2(n-1) + (n+5)(3n-2)(n-1) \\ &= (n-1)(n+5)(4n+3). \end{aligned}$$

Case-4: Consider the The fifth multiplicative M_1 Zagreb index $M_1G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u) + S_G(v)]$. Using Table 3, we have

$$\begin{aligned} M_1G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u) + S_G(v)] \\ &= (n-1)(n+5+n+5) \times (n-1)(n+5+3n-3) \\ &= (n-1)^2(4n+2)(2n+10). \end{aligned}$$

Case-5: Consider the The fifth multiplicative M_2 Zagreb index $M_2G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u)S_G(v)]$. Using Table 3, we have

$$\begin{aligned} M_2G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u)S_G(v)] \\ &= (n+5)(n+5)(n-1) \times (n-1)(n+5)(3n-3) \\ &= (3n-3)(n-1)^2(n+5)^3. \end{aligned}$$

Theorem 4:4 Let $H = L(W_n)$ be the line graph of wheel graph W_n . Then

1. $(H) = \frac{(2n+8)^6(n-1)}{4n+14} + \frac{2(2n+8)^3(n^2-2n+8)^3(n-1)}{n^2+14} + \frac{(n^2-2n+8)^6(n-1)(n-2)}{2[(n^2-2n+8)-2]}$
2. $M_1G_5(H) = (4n+16)(n-1) + 2(n^2+16)(n-1) + (n-1)(n-2)(n^2-2n+8)$
3. $M_2G_5(H) = (2n+8)^2(n-1) + 2(2n+8)(n^2-2n+8)(n-1) + \frac{(n-1)(n-2)}{2}(n^2-2n+8)^2$

$$4. \quad M_1 G_5 \prod(H) = (n-1)(n-2)(n^2 - 2n + 8)^2(n-1)^2(4n+16)(n^2 + 16)$$

$$5. \quad M_2 G_5 \prod(H) = (n^2 - 2n + 8)(n-1)(n-2)(n^2 - 2n + 8)(2n+8)(n-1)(2n+8)^2(n-1)$$

Proof: Let $G = W_n$ be a wheel graph of order n and let $H = L(W_n)$ be the line graph of wheel graph. Now consider the following cases:

Case-1: Consider the Sanskruti index $S(G) = \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u) + s_G(v) - 2} \right)^3$. By using the information in Table 4,

we have

$$\begin{aligned} S(G) &= \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u) + s_G(v) - 2} \right)^3 \\ &= (n-1) \left(\frac{(2n+8)^2}{2n+8+2n+8-2} \right)^3 + 2(n-1) \left(\frac{(2n+8)(n^2-2n+8)}{2n+8+n^2-2n+8-2} \right)^3 \\ &\quad + \frac{(n-1)(n-2)}{2} \left(\frac{(n^2-2n+8)^2}{n^2-2n+8+n^2-2n+8-2} \right)^3 \\ &= \frac{(2n+8)^6(n-1)}{4n+14} + \frac{2(2n+8)^3(n^2-2n+8)^3(n-1)}{n^2+14} + \frac{(n^2-2n+8)^6(n-1)(n-2)}{2[(n^2-2n+8)-2]}. \end{aligned}$$

Case-2: Consider the fifth M_1 Zagreb Index, $M_1 G_5(G) = \sum_{uv \in E(G)} [S_G(u) + S_G(v)]$. Again by using the information in Table 4, we have

$$\begin{aligned} M_1 G_5(G) &= \sum_{uv \in E(G)} [S_G(u) + S_G(v)] \\ &= (n-1)(2n+8+2n+8) + 2(n-1)(2n+8+n^2-2n+8) \\ &\quad + \frac{(n-1)(n-2)}{2}(n^2-2n+8+n^2-2n+8) \\ &= 4n+16)(n-1) + 2(n^2+16)(n-1) + (n-1)(n-2)(n^2-2n+8). \end{aligned}$$

Case-3: Consider the fifth M_2 Zagreb Index, $M_2 G_5(G) = \sum_{uv \in E(G)} [S_G(u)S_G(v)]$. Again by using the information in Table 4, we have

$$\begin{aligned} M_1 G_5(G) &= \sum_{uv \in E(G)} [S_G(u)S_G(v)] \\ &= (n-1)(2n+8)^2 + 2(n-1)(2n+8)(n^2-2n+8) + \frac{(n-1)(n-2)}{2}(n^2-2n+8)^2. \end{aligned}$$

Case-4: Consider the The fifth multiplicative M_1 Zagreb index $M_1 G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u) + S_G(v)]$. Using Table 4, we have

$$\begin{aligned} M_1 G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u) + S_G(v)] \\ &= (n-1)(2n+8+2n+8) \times 2(n-1)(2n+8+n^2-2n+8) \\ &\quad \times \frac{(n-1)(n-2)}{2}(n^2-2n+8+n^2-2n+8) \\ &= (n-1)(n-2)(n^2-2n+8)^2(n-1)^2(4n+16)(n^2+16). \end{aligned}$$

Case-5: Consider the The fifth multiplicative M_2 Zagreb index $M_2G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u)S_G(v)]$. Using

Table 4, we have

$$\begin{aligned}
 M_2G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u)S_G(v)] \\
 &= (n-1)(2n+8)^2 \times 2(n-1)(2n+8)(n^2-2n+8) \times \frac{(n-1)(n-2)}{2} (n^2-2n+8)^2 \\
 &= (n^2-2n+8)(n-1)(n-2)(n^2-2n+8)(2n+8)(n-1)(2n+8)^2(n-1).
 \end{aligned}$$

4. TOPOLOGICAL INDICES OF THE LADDER GRAPH AND ITS LINE GRAPH




Fig. 3. The ladder graph $P_n \times P_2$ and its line graph $L(P_n \times P_2)$

Table-5:The edge partition of the ladder graph $P_n \times P_2; n \geq 4$.

(S_u, S_v) where $uv \in E(G)$	(5,5)	(5,8)	(8,9)	(8,8)	(9,9)
Number of edges	2	4	4	2	$3n-14$

Table-6: The edge partition of the line graph of ladder graph $P_n \times P_2; n \geq 6$.

(S_u, S_v) where $uv \in E(G)$	(6,10)	(10,15)	(10,14)	(14,15)	(15,16)	(16,16)
Number of edges	4	4	4	4	8	$6n-32$

Theorem 5:5 Let $G = P_n \times P_2$ be the ladder graph. Then

1. $S(G) = (129.7463)(3n-14) + 886.7929$
2. $M_1G_5(G) = 54n-80$
3. $M_2G_5(G) = 243n-80$
4. $M_1G_5 \prod(G) = 18(3n-14)(2263040)$
5. $M_2G_5 \prod(G) = 81(3n-4)(294912000)$

Proof: Let $G = P_n \times P_2$ be a ladder graph with $|V(G)| = 2n$ and $|E(G)| = 3n-2$. Since it is easy to calculate the topological indices value for G up to $n=3$. Therefore we consider $n \geq 4$. Now consider the following cases:

Case-1: Consider the Sanskruti index $S(G) = \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u)+s_G(v)-2} \right)^3$. By using the information in Table 5,

we have

$$\begin{aligned}
 S(G) &= \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u)+s_G(v)-2} \right)^3 \\
 &= 2 \left(\frac{5 \times 5}{5+5-2} \right)^3 + 4 \left(\frac{5 \times 8}{5+8-2} \right)^3 + 4 \left(\frac{8 \times 9}{8+9-2} \right)^3 + 2 \left(\frac{8 \times 8}{8+8-2} \right)^3 + (3n-4) \left(\frac{9 \times 9}{9+9-2} \right)^3 \\
 &= (129.7463)(3n-14) + 886.7929.
 \end{aligned}$$

Case-2: Consider the fifth M_1 Zagreb Index, $M_1G_5(G) = \sum_{uv \in E(G)} [S_G(u) + S_G(v)]$. Again by using the information in Table 5, we have

$$\begin{aligned} M_1G_5(G) &= \sum_{uv \in E(G)} [S_G(u) + S_G(v)] \\ &= 2(5+5) + 4(5+8) + 4(8+9) + 2(8+8) + (3n-4)(9+9) \\ &= 54n - 80. \end{aligned}$$

Case-3: Consider the fifth M_2 Zagreb Index, $M_2G_5(G) = \sum_{uv \in E(G)} [S_G(u)S_G(v)]$. Again by using the information in Table 5, we have

$$\begin{aligned} M_2G_5(G) &= \sum_{uv \in E(G)} [S_G(u)S_G(v)] \\ &= 2(5 \times 5) + 4(5 \times 8) + 4(8 \times 9) + 2(8 \times 8) + (3n-4)(9 \times 9) \\ &= 243n - 508. \end{aligned}$$

Case-4: Consider the The fifth multiplicative M_1 Zagreb index $M_1G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u) + S_G(v)]$. Using Table 5, we have

$$\begin{aligned} M_1G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u) + S_G(v)] \\ &= 2(5+5) \times 4(5+8) \times 4(8+9) \times 2(8+8) \times (3n-4)(9+9) \\ &= 18(3n-14)(2263040). \end{aligned}$$

Case-5: Consider the The fifth multiplicative M_2 Zagreb index $M_2G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u)S_G(v)]$. Using Table 5, we have

$$\begin{aligned} M_2G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u)S_G(v)] \\ &= 2(5 \times 5) \times 4(5 \times 8) \times 4(8 \times 9) \times 2(8 \times 8) \times (3n-4)(9 \times 9) \\ &= 81(3n-4)(294912000). \end{aligned}$$

Theorem 6:6 Let $H = L(P_n \times P_2)$ be the line graph of ladder graph $P_n \times P_2$. Then

1. $S(H) = (621.3735)(6n - 32) + 8871.0265$
2. $M_1G_5(H) = 192n - 375$
3. $M_2G_5(H) = 1536n - 3882$
4. $M_1G_5 \prod(H) = 248(6n - 32)(71270400)$
5. $M_2G_5 \prod(H) = (6n - 32)(80640000)(412876800)$

Proof: Let $G = P_n \times P_2$ be a ladder graph with $|V(G)| = 2n$ and $|E(G)| = 3n - 2$ and let $H = L(P_n \times P_2)$ be the line graph of $P_n \times P_2$. Since it is easy to calculate the topological indices value for G up to $n = 3$. Therefore we consider $n \geq 4$. Now consider the following cases:

Case-1: Consider the Sanskruti index $S(G) = \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u) + s_G(v) - 2} \right)^3$. By using the information in Table 6, we have

$$\begin{aligned}
S(G) &= \sum_{uv \in E(G)} \left(\frac{s_G(u)s_G(v)}{s_G(u) + s_G(v) - 2} \right)^3 \\
&= 4 \left(\frac{6 \times 10}{6 + 10 - 2} \right)^3 + 4 \left(\frac{10 \times 15}{10 + 15 - 2} \right)^3 + 4 \left(\frac{10 \times 14}{10 + 14 - 2} \right)^3 \\
&\quad + 4 \left(\frac{14 \times 15}{14 + 15 - 2} \right)^3 + 8 \left(\frac{15 \times 16}{15 + 16 - 2} \right)^3 + (6n - 32) \left(\frac{16 \times 16}{16 + 16 - 2} \right)^3 \\
&= (621.3735)(6n - 32) + 8871.0265.
\end{aligned}$$

Case-2: Consider the fifth M_1 Zagreb Index, $M_1G_5(G) = \sum_{uv \in E(G)} [S_G(u) + S_G(v)]$. Again by using the information in Table 6, we have

$$\begin{aligned}
M_1G_5(G) &= \sum_{uv \in E(G)} [S_G(u) + S_G(v)] \\
&= 4(6 + 10) + 5(10 + 15) + 4(10 + 14) + 4(14 + 15) + 8(15 + 16) + (6n - 32)(16 + 16) \\
&= 192n - 375.
\end{aligned}$$

Case-3: Consider the fifth M_2 Zagreb Index, $M_2G_5(G) = \sum_{uv \in E(G)} [S_G(u)S_G(v)]$. Again by using the information in Table 6, we have

$$\begin{aligned}
M_1G_5(G) &= \sum_{uv \in E(G)} [S_G(u)S_G(v)] \\
&= 4(6 \times 10) + 5(10 \times 15) + 4(10 \times 14) + 4(14 \times 15) + 8(15 \times 16) + (6n - 32)(16 \times 16) \\
&= 1536n - 3882.
\end{aligned}$$

Case-4: Consider the The fifth multiplicative M_1 Zagreb index $M_1G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u) + S_G(v)]$. Using Table 6, we have

$$\begin{aligned}
M_1G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u) + S_G(v)] \\
&= 4(6 + 10) \times 5(10 + 15) \times 4(10 + 14) \times 4(14 + 15) \\
&\quad \times 8(15 + 16) \times (6n - 32)(16 + 16) \\
&= (248)(6n - 32)(71270400).
\end{aligned}$$

Case-5: Consider the The fifth multiplicative M_2 Zagreb index $M_2G_5 \prod(G) = \prod_{uv \in E(G)} [S_G(u)S_G(v)]$. Using Table 5, we have

$$\begin{aligned}
M_2G_5 \prod(G) &= \prod_{uv \in E(G)} [S_G(u)S_G(v)] \\
&= 4(6 \times 10) \times 5(10 \times 15) \times 4(10 \times 14) \times 4(14 \times 15) \times 8(15 \times 16) \times (6n - 32)(16 \times 16) \\
&= (6n - 32)(80640000)(412876800).
\end{aligned}$$

REFERENCES

1. B.Basavanagoud, A.P.Barangi and S.M.Hosamani, First neighbourhood Zagreb index of some nanostructures, Proceedings IAM. 7(2) (2018) 178--193.
2. A.Graovac, M.Ghorbani and M.A.Hosseinzadeh, Computing fifth geometricarithmetic index of nanostar dendrimers, Journal of Mathematical Nanoscience, 1(1)(2011) 33-42.
3. Gutman, I., Trinajsti c' , N., (1972), Graph theory and molecular orbitals. Total π -electron energy of alternant hydrocarbons, Chem. Phys. Lett., 17, pp. 535--538.
4. Harary, F., (1969), Graph Theory, Addison--Wesely, Reading.
5. Hosamani, S. M., and Gutman, I., (2014), Zagreb indices of transformation graphs and total transformation graphs, Appl. Math. Comput., 247, pp. 1156--1160.

6. Hosamani, S. M., Basavanagoud, B., (2015), New upper bounds for the first Zagreb index, MATCH Commun. Math. Comput. Chem., 74(1), pp. 97--101.
7. Hosamani S. M, Malghan S. H and Cangul I. N, The first geometric-arithmetic index of graph operations, Advances and Applications in Mathematical Sciences, 14(6) (2015) 155--163.
8. Hosamani S. M, Computing Sanskruti index of certain nanostructures, J. Appl. Math. Comput. 1-9 (2016) DOI 10.1007/s12190-016-1016-9.
9. Kulli V.R, Some new multiplicative geometric-arithmetic index, Journal of Ultra Scientist of Physical Sciences, A, 29(2) (2017) 52-57.
10. Li, X., and Zhao, H., (2004), Trees with the first three smallest and largest generalized topological indices, MATCH Commun. Math. Comput. Chem., 50, pp. 57--62.
11. Nadeem, M.F., Zafar, S., Zahid, Z., (2015), Certain topological indices of the line graph of subdivision graphs, Appl. Math. Comput., (271), pp. 790--794.
12. Ranjini, P.S., Loksha, V., Cangul, I.N., (2011), On the Zagreb indices of the line graphs of the subdivision graphs, Appl. Math. Comput., 218, pp. 699--702.
13. Ranjini, P.S., Loksha, V., Rajan, M.A., (2011), On the Shultz index of the subdivision graphs, Adv. Stud. Contemp. Math., 21(3), pp. 279-290.
14. Su, G., Xu, L., (2015), Topological indices of the line graph of subdivision graphs and their Schur-bounds, Appl. Math. Comput., 253, pp. 395--401.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2019. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]