

## LOWER ANTI Q-FUZZY GROUP AND ITS LOWER LEVEL SUBGROUPS

R. Muthuraj\*

Department of Mathematics, H. H. The Rajah's College, Pudukkottai – 622 001, Tamilnadu, India

E-mail: [rmr1973@yahoo.co.in](mailto:rmr1973@yahoo.co.in)

&

M. S. Muthuraman , P. M. Sithar Selvam , K. H. Manikandan

Department of Mathematics, PSNA College of Engineering and Technology,  
Dindigul-624 622, Tamilnadu , India

E-mail : [msmraman@yahoo.com](mailto:msmraman@yahoo.com), [harisithar@rediffmail.com](mailto:harisithar@rediffmail.com)

(Received on: 20-09-11; Accepted on: 06-10-11)

### ABSTRACT

In this paper, we define the algebraic structures of a lower anti Q-fuzzy subgroup and some related properties are investigated. We establish the relation between upper Q-fuzzy group and lower anti Q-fuzzy group of a group. The purpose of this study is to implement the fuzzy set theory and group theory in upper Q-fuzzy groups and lower anti Q-fuzzy groups. Characterizations of lower level subsets of a lower anti Q-fuzzy group of a group are given. We also discussed the relation between a given a lower anti Q-fuzzy group and its lower level sub groups and investigate the conditions under which a given group has a properly inclusive chain of sub groups. In particular, we formulate how to structure a lower anti Q-fuzzy group by a given chain of sub groups.

**Keywords:** Fuzzy set, Q-fuzzy set, fuzzy subgroup, Q-fuzzy subgroup, anti-Q fuzzy subgroup, upper and lower anti-Q fuzzy subgroup, lower level subsets .

**AMS Subject Classification (2000):** 20N25, 03E72, 03F055, 06F35, 03G25.

## 1. INTRODUCTION

The notion of fuzzy sets was introduced by L.A. Zadeh [13]. Fuzzy set theory has been developed in many directions by many researchers and has evoked great interest among mathematicians working in different fields of mathematics. In 1971, Rosenfield [10] introduced the concept of fuzzy subgroup. R. Biswas [1] introduced the concept of anti- fuzzy subgroups of groups. K. H. Kim [5] introduced the concept of intuitionistic Q-fuzzy semi prime ideals in semi groups and Osman kazanci, sultan yamark and serife yilmaz [8] introduced the concept of intuitionistic Q-fuzzy R-subgroups of near rings. A.Solairaju and R.Nagarajan[12] introduced and defined a new algebraic structure of Q-fuzzy groups. R. Muthuraj, M.S. Muthuraman, P. M.Sithar selvam [7] introduced the concept of anti Q-fuzzy groups. In this paper we define a new algebraic structure of lower anti Q-fuzzy subgroups and study some their related properties.

## 2. PRELIMINARIES

In this section we site the fundamental definitions that will be used in the sequel.

**2.1 Definition: [13]** Let S be any non empty set. A fuzzy subset A of S is a function  $A: S \rightarrow [0, 1]$ .

**2.2 Definition: [10]** Let G be a group. A fuzzy subset A of G is called a fuzzy subgroup if for  $x, y \in G$ ,

- i.  $A(xy) \geq \min \{ A(x), A(y) \}$ ,
- ii.  $A(x^{-1}) = A(x)$ .

**2.3 Definition: [1]** Let G be a group. A fuzzy subset A of G is called an anti fuzzy subgroup if for  $x, y \in G$ ,

- i.  $A(xy) \leq \max \{ A(x), A(y) \}$ ,
- ii.  $A(x^{-1}) = A(x)$ .

---

**\*Corresponding author: R. Muthuraj\*, \*E-mail: [rmr1973@yahoo.co.in](mailto:rmr1973@yahoo.co.in)**

**2.4 Definition [12]** Let Q and G be any two sets. A mapping  $A: G \times Q \rightarrow [0, 1]$  is called a Q-fuzzy set in G.

**2.5 Definition [12]** A Q-fuzzy set 'A' is called Q-fuzzy group of a group G if for  $x, y \in G, q \in Q$ ,

- i.  $A(xy, q) \geq \min \{A(x, q), A(y, q)\}$
- ii.  $A(x^{-1}, q) = A(x, q)$ .

**2.6 Definition:** A Q-fuzzy set 'A' is called an upper Q-fuzzy group of a group G if for  $x, y \in G, q \in Q$ ,

- i.  $A(xy, q) \geq \min \{A(x, q), A(y, q)\}$ ,
- ii.  $A(x^{-1}, q) = A(x, q)$ ,
- iii.  $A(e, q) = 1$ .

**2.7 Definition: [7]** A Q-fuzzy set 'A' is called an anti Q-fuzzy group of a group G if for  $x, y \in G, q \in Q$ ,

- i.  $A(xy, q) \leq \max \{A(x, q), A(y, q)\}$
- ii.  $A(x^{-1}, q) = A(x, q)$ .

**2.8 Definition:** A Q-fuzzy set 'A' is called a lower anti Q-fuzzy group of a group G if for  $x, y \in G, q \in Q$ ,

- i.  $A(xy, q) \leq \max \{A(x, q), A(y, q)\}$ ,
- ii.  $A(x^{-1}, q) = A(x, q)$ ,
- iii.  $A(e, q) = 0$ .

**2.9 Definition:** An lower anti Q-fuzzy group of a group G is said to be Normal if  $A(xy, q) = A(yx, q)$  for  $x, y \in G, q \in Q$ .

### 3. PROPERTIES OF LOWER ANTI Q-FUZZY SUBGROUPS

In this section, we discuss some of the properties of lower anti Q-fuzzy subgroups.

**3.1 Theorem:** Let 'A' be an lower anti Q-fuzzy subgroup of a group G then

- i.  $A(x, q) \geq 0$  for all  $x \in G, q \in Q$ .
- ii. The subset  $H = \{x \in G / A(x, q) = 0\}$  is a subgroup of G.

**Proof:**

- i. Let  $x \in G$  and  $q \in Q$ .

$$\begin{aligned} A(x, q) &= \max \{A(x, q), A(x, q)\} \\ &= \max \{A(x, q), A(x^{-1}, q)\} \\ &\geq A(xx^{-1}, q) \\ &= A(e, q). \\ A(x, q) &\geq 0. \end{aligned}$$

- ii. Let  $H = \{x \in G / A(x, q) = 0\}$ .

Clearly H is non-empty as  $e \in H$ . Let  $x, y \in H$ .

Then,  $A(x, q) = A(y, q) = 0$ .

$$\begin{aligned} A(xy^{-1}, q) &\leq \max \{A(x, q), A(y^{-1}, q)\} \\ &= \max \{A(x, q), A(y, q)\} \\ &= \max \{0, 0\} \\ &= 0. \end{aligned}$$

That is,  $A(xy^{-1}, q) \leq 0$  and obviously  $A(xy^{-1}, q) \geq A(e, q) = 0$ .

Hence,  $A(xy^{-1}, q) = 0$  and  $xy^{-1} \in H$ .

Clearly, H is a subgroup of G.

**3.2 Theorem:** If 'A' is an upper Q-fuzzy subgroup of G, iff  $A^c$  is a lower anti Q-fuzzy subgroup of G.

**Proof:** Suppose A is an upper Q-fuzzy subgroup of G. Then for all  $x, y \in G$  and  $q \in Q$ ,

$$\begin{aligned} A(xy, q) &\geq \min \{A(x, q), A(y, q)\} \\ \Leftrightarrow 1 - A^c(xy, q) &\geq \min \{(1 - A^c(x, q)), (1 - A^c(y, q))\} \\ \Leftrightarrow A^c(xy, q) &\leq 1 - \min \{(1 - A^c(x, q)), (1 - A^c(y, q))\} \end{aligned}$$

$$\Leftrightarrow A^c(xy, q) \leq \max \{ A^c(x, q), A^c(y, q) \}.$$

We have,  $A(x, q) = A(x^{-1}, q)$  for all  $x$  in  $G$  and  $q \in Q$ ,  
 $\Leftrightarrow 1 - A^c(x, q) = 1 - A^c(x^{-1}, q)$ .

Therefore,  $A^c(x, q) = A^c(x^{-1}, q)$ .

Also,  $A^c(e, q) = 1 - A(e, q) = 0$

Hence  $A^c$  is a lower anti Q-fuzzy subgroup of  $G$ .

**3.3 Theorem:** Let  $A$  be any lower anti Q-fuzzy subgroup of a group  $G$  with identity  $e$ . Then  $A(xy^{-1}, q) = 0 \Rightarrow A(x, q) = A(y, q)$  for all  $x, y$  in  $G$  and  $q \in Q$ .

**Proof:** Given  $A$  is a lower anti Q-fuzzy subgroup of  $G$  and  $A(xy^{-1}, q) = 0$ .  
 Then for all  $x, y$  in  $G$  and  $q \in Q$ ,

$$\begin{aligned} A(x, q) &= A(x(y^{-1}y), q) \\ &= A((xy^{-1})y, q) \\ &\leq \max \{ A(xy^{-1}, q), A(y, q) \} \\ &= \max \{ 0, A(y, q) \} \\ &= A(y, q). \end{aligned}$$

That is,  $A(x, q) \leq A(y, q)$ .

Now,  $A(y, q) = A(y^{-1}, q)$ , since  $A$  is a lower anti Q-fuzzy subgroup of  $G$ .

$$\begin{aligned} &= A(ey^{-1}, q) \\ &= A((x^{-1}x)y^{-1}, q) \\ &= A(x^{-1}(xy^{-1}), q) \\ &\leq \max \{ A(x^{-1}, q), A(xy^{-1}, q) \} \\ &= \max \{ A(x, q), 0 \} \\ &= A(x, q). \end{aligned}$$

That is,  $A(y, q) \leq A(x, q)$ .

Hence,  $A(x, q) = A(y, q)$ .

**3.4 Theorem:**  $A$  is a lower anti Q-fuzzy subgroup of a group  $G$  if and only if  $A(xy^{-1}, q) \leq \max \{ A(x, q), A(y, q) \}$ , for all  $x, y$  in  $G$  and  $q \in Q$ .

**Proof:** Let  $A$  be a lower anti Q-fuzzy subgroup of a group  $G$ . Then for all  $x, y$  in  $G$  and  $q \in Q$ ,

$$A(xy, q) \leq \max \{ A(x, q), A(y, q) \} \text{ and } A(x, q) = A(x^{-1}, q) \text{ with } A(e, q) = 0.$$

Now,  $A(xy^{-1}, q) \leq \max \{ A(x, q), A(y^{-1}, q) \}$   
 $= \max \{ A(x, q), A(y, q) \}$   
 $\Leftrightarrow A(xy^{-1}, q) \leq \max \{ A(x, q), A(y, q) \}.$

#### 4. PROPERTIES OF LOWER LEVEL SUBSETS OF A LOWER ANTI Q-FUZZY SUBGROUP

In this section, we introduce the concept of lower level subset of a lower anti Q-fuzzy subgroup and discuss some of its properties.

**4.1 Definition:** Let  $A$  be a lower anti Q-fuzzy group of a group  $G$ . For any  $t \in [0, 1]$ , we define the lower level subset of  $A$  is the set,  $L(A; t) = \{x \in G / A(x, q) \leq t \text{ and } q \in Q\}$ .

**4.1 Theorem:** Let  $A$  be a lower anti Q-fuzzy subgroup of a group  $G$ . Then for  $t \in [0, 1]$  such that  $t \geq 0$ ,  $L(A; t)$  is a subgroup of  $G$ .

**Proof:**  $L(A; 0)$  is non empty as  $e \in L(A; 0)$ .

For all  $x, y \in L(A; t)$  and  $q \in Q$ , we have,  $A(x, q) \leq t$ ;  $A(y, q) \leq t$ .

Now,  $A(xy^{-1}, q) \leq \max \{ A(x, q), A(y, q) \}$ .  
 $A(xy^{-1}, q) \leq \max \{ t, t \}$ .  
 $A(xy^{-1}, q) \leq t$ .

$$xy^{-1} \in L(A; t).$$

Hence  $L(A; t)$  is a subgroup of  $G$ .

**4.2 Theorem:** Let  $G$  be a group and  $A$  be a  $Q$ -fuzzy subset of  $G$  such that  $L(A; t)$  is a subgroup of  $G$ . For  $t \in [0, 1]$  such that  $t \geq 0$ ,  $A$  is a lower anti  $Q$ -fuzzy subgroup of  $G$ .

**Proof:** Let  $x, y$  in  $G$  and  $q \in Q$ , let  $A(x, q) = t_1$  and  $A(y, q) = t_2$ .

Suppose  $t_1 < t_2$ , then  $x, y \in L(A; t_2)$ .

As  $L(A; t_2)$  is a subgroup of  $G$ ,  $xy^{-1} \in L(A; t_2)$ .

$$\begin{aligned} \text{Hence, } A(xy^{-1}, q) &\leq t_2 = \max\{t_1, t_2\} \\ &\leq \max\{A(x, q), A(y, q)\} \end{aligned}$$

That is,  $A(xy^{-1}, q) \leq \max\{A(x, q), A(y, q)\}$ .

By Theorem 3.4,  $A$  is a lower anti  $Q$ -fuzzy subgroup of  $G$ .

**4.2 Definition:** Let  $A$  be a lower anti  $Q$ -fuzzy subgroup of a group  $G$ . The subgroups  $L(A; t)$  for  $t \in [0, 1]$  and  $t \geq 0$ , are called lower level subgroups of  $A$ .

**4.3 Theorem:** Let  $A$  be a lower anti  $Q$ -fuzzy subgroup of a group  $G$ . If two lower level subgroups  $L(A; t_1), L(A; t_2)$ , for,  $t_1, t_2 \in [0, 1]$  and  $t_1, t_2 \geq 0$  with  $t_1 < t_2$  of  $A$  are equal then there is no  $x$  in  $G$  such that  $t_1 < A(x, q) \leq t_2$ .

**Proof:** Let  $L(A; t_1) = L(A; t_2)$ .

Suppose there exists a  $x \in G$  such that  $t_1 < A(x, q) \leq t_2$  then  $L(A; t_1) \subsetneq L(A; t_2)$ .

Then  $x \in L(A; t_2)$ , but  $x \notin L(A; t_1)$ , which contradicts the assumption that,  $L(A; t_1) = L(A; t_2)$ . Hence there is no  $x$  in  $G$  such that  $t_1 < A(x, q) \leq t_2$ .

Conversely, suppose that there is no  $x$  in  $G$  such that  $t_1 < A(x, q) \leq t_2$ .

Then, by definition,  $L(A; t_1) \subseteq L(A; t_2)$ .

Let  $x \in L(A; t_2)$  and there is no  $x$  in  $G$  such that  $t_1 < A(x, q) \leq t_2$ .

Hence  $x \in L(A; t_1)$  and  $L(A; t_2) \subseteq L(A; t_1)$ .

Hence  $L(A; t_1) = L(A; t_2)$ .

**4.4 Theorem:** A  $Q$ -fuzzy subset  $A$  of  $G$  is a lower anti  $Q$ -fuzzy subgroup of a group  $G$  if and only if the lower level subsets  $L(A; t)$ ,  $t \in \text{Image } A$ , are subgroups of  $G$ .

**Proof:** It is clear.

**4.5 Theorem:** Any subgroup  $H$  of a group  $G$  can be realized as a lower level subgroup of some lower anti  $Q$ -fuzzy subgroup of  $G$ .

**Proof:** Let  $A$  be a  $Q$ -fuzzy subset and  $x \in G$  and  $q \in Q$ .

$$\text{Define, } A(x, q) = \begin{cases} 0 & \text{if } x \in H \\ t & \text{if } x \notin H, \text{ where } t \in (0, 1]. \end{cases}$$

We shall prove that  $A$  is a lower anti  $Q$ -fuzzy subgroup of  $G$ .

Let  $x, y \in G$  and  $q \in Q$ .

(i) Suppose  $x, y \in H$ , then  $xy \in H$  and  $xy^{-1} \in H$ .  $A(x, q) = 0, A(y, q) = 0, A$  and  $A(xy^{-1}, q) = 0$ .  
 Hence  $A(xy^{-1}, q) \leq \max\{A(x, q), A(y, q)\}$ .

(i) Suppose  $x \in H$  and  $y \notin H$ , then  $xy \notin H$  and  $xy^{-1} \notin H$ .  $A(x, q) = 0$ ,  $A(y, q) = t$  and  $A(xy^{-1}, q) = t$ .  
Hence  $A(xy^{-1}, q) \leq \max \{ A(x, q), A(y, q) \}$ .

(iii) Suppose  $x, y \notin H$ , then  $xy^{-1} \in H$  or  $xy^{-1} \notin H$ .  $A(x, q) = t$ ,  $A(y, q) = t$  and  $A(xy^{-1}, q) = 0$  or  $t$ .  
Hence  $A(xy^{-1}, q) \leq \max \{ A(x, q), A(y, q) \}$ .

Since  $H$  is a subgroup and  $e \in H$ , then  $A(e, q) = 0$ .

Thus in all cases,  $A$  is a lower anti Q-fuzzy subgroup of  $G$ .

For this lower anti Q-fuzzy subgroup,  $L(A; t) = H$ .

**Remark:** As a consequence of the Theorem 4.3, the lower level subgroups of a lower anti Q-fuzzy subgroup  $A$  of a group  $G$  form a chain. Since  $0 \leq A(x, q)$  for all  $x$  in  $G$  and  $q \in Q$ , therefore  $L(A; 0)$  is the smallest and we have the chain :

$\{e\} \subset L(A; 0) \subset L(A; t_1) \subset L(A; t_2) \subset \dots \subset L(A; t_n) = G$ , where  $0 < t_1 < t_2 < \dots < t_n$ .

## REFERENCES

- [1] Biswas .R, Fuzzy subgroups and Anti Fuzzy subgroups, Fuzzy sets and Systems, 35(1990) 121-124.
- [2] Das. P. S, Fuzzy groups and level subgroups, J. Math. Anal. Appl, 84 (1981), 264-269.
- [3] K. H. Kim, Y. B. Yun, on fuzzy R- subgroups of near rings, J.fuzzy math 8 (3) (2000), 549-558.
- [4] K. H. Kim, Y. B. Jun. Normal fuzzy R- subgroups in near rings, Fuzzy sets systems 121 (2001), 341-345.
- [5] K. H. Kim, on intuitionistic Q- fuzzy semi prime ideals in semi groups, Advances in fuzzy mathematics, 1 (1) (2006) 15-21.
- [6] Mohamed Asaad, Groups and Fuzzy subgroups Fuzzy sets and systems 39(1991), 323-328.
- [7] R. Muthuraj, M. S. Muthuraman, P. M. Sithar selvam, Anti Q-fuzzy group and its lower level subgroups, International Journal of Computer Applications (0975 – 8887),Volume 3 – No.3, 16-20, June 2010
- [8] Osman kazanci, sultan yamark and serife yilmaz “On intuitionistic Q- fuzzy R-subgroups of near rings” International mathematical forum, 2, 2007 no. 59, 2899-2910.
- [9] Palaniappan N, Muthuraj .R , Anti fuzzy group and Lower level subgroups, Antartica J. Math., 1 (1) (2004) , 71-76.
- [10] Rosenfeld. A., fuzzy groups, J. math. Anal. Appl. 35 (1971), 512-517.
- [11] Solairaju. A, Nagarajan. R, Q- fuzzy left R- subgroups of near rings w.r.t T- norms, Antarctica journal of mathematics.5, (1-2), 2008.
- [12] Solairaju.A, Nagarajan.R, A New Structure and Construction of Q-Fuzzy Groups, Advances in fuzzy mathematics, Volume 4, Number 1 (2009) pp.23-29.
- [13] Zadeh. L. A., Fuzzy sets, Information and control, 8, (1965) 338-353.

\*\*\*\*\*