

SECOND ZAGREB-K-BANHATTI INDEX OF A GRAPH

V. R. KULLI*

Department of Mathematics, Gulbarga University, Gulbarga - 585106, India.

(Received On: 11-09-20; Revised & Accepted On: 18-09-20)

ABSTRACT

The *K*-Banhatti indices were introduced by Kulli in 2016. The Zagreb indices were proposed by Gutman and Trinajstić in 1972. The Zagreb indices and *K*-Banhatti indices are closely related. In this paper, we establish some relations between Zagreb, *K*-Banhatti and second Zagreb-*K*-Banhatti indices. Also we obtain lower and upper bounds for the second Zagreb-*K*-Banhatti index of a graph in terms of Zagreb and *K*-Banhatti indices.

Keywords: Zagreb index, *K*-Banhatti index, Zagreb-*K*-Banhatti index, second Zagreb-*K*-Banhatti index, graph.

Mathematics Subject Classification: 05C05, 05C07, 05C35.

1. INTRODUCTION

A molecular graph is a finite, simple graph, representing the carbon-atom skeleton of an organic molecule of a hydrocarbon. The vertices of a molecular graph represent the carbon atoms and its undirected edges the carbon-carbon bonds. Chemical Graph Theory is a branch of Mathematical Chemistry. This branch of Mathematics has an important effect on the development of Chemical Sciences. In Chemical Science, physicochemical properties of chemical compounds often modeled by means of molecular based structure descriptors, which are also referred as topological indices or graph indices. For graph indices, see [1]. A topological index or a graph index is a numerical parameter mathematically derived from the graph structure.

These indices are useful for establishing correlation between the structure a molecular compound and its physicochemical properties. Several such topological indices have been considered in Theoretical Chemistry and have obtained some applications, especially in QSPR/QSAR research, see [2, 3, 4].

Let G be a simple, connected graph with n vertices and m edges with vertex set $V(G)$ and edge set $E(G)$. The degree $d_G(u)$ of a vertex u is the number of vertices adjacent to u . The edge connecting the vertices u and v will be denoted by uv . Let $d_G(e)$ denote the degree of an edge $e = uv$ in a graph G , which is defined by $d_G(e) = d_G(u) + d_G(v) - 2$. The vertices and edges of a graph G are called its elements. Any undefined term in this paper may be found in [5].

The first and second Zagreb indices take into account the contributions of pairs of adjacent vertices. The first and second Zagreb indices were introduced by Gutman et al. in [6] and they are defined as

$$M_1(G) = \sum_{u \in V(G)} d_G(u)^2 = \sum_{uv \in E(G)} [d_G(u) + d_G(v)], \quad M_2(G) = \sum_{uv \in E(G)} d_G(u)d_G(v).$$

Zagreb indices have been studied extensively. For their history, applications and mathematical properties, see [7, 8, 9, 10, 11].

Followed by the first Zagreb index of a graph G , Furtula *et al.* [12] introduced the forgotten topological index and defined it as

$$F(G) = \sum_{u \in V(G)} d_G(u)^3 = \sum_{uv \in E(G)} [d_G(u)^2 + d_G(v)^2].$$

This index was studied, for example, in [13, 14, 15].

Corresponding Author: V. R. Kulli*

Department of Mathematics, Gulbarga University, Gulbarga - 585106, India.

The first and second K -Banhatti indices take into account the contributions of pairs of incident elements. The first and second K -Banhatti indices were introduced by Kulli in [16] and they are defined as

$$B_1(G) = \sum_{ue} [d_G(u) + d_G(e)], \quad B_2(G) = \sum_{ue} d_G(u) d_G(e).$$

where ue means that the vertex u and edge e are incident in G .

The first and second hyper Bhanhatti indices of a graph G are defined by Kulli in [17] as

$$HB_1(G) = \sum_{ue} [d_G(u) + d_G(e)]^2, \quad HB_2(G) = \sum_{ue} d_G(u) d_G(e).$$

These indices have been studied extensively. For their history, applications and mathematical properties, see [18, 19, 20, 21, 22, 23, 24, 25, 26].

In [27], Miličević *et al.* introduced the first and second reformulated Zagreb indices of a graph G in terms of edges instead of vertex degrees and defined as

$$EM_1(G) = \sum_{e \in E(G)} d_G(e)^2, \quad EM_2(G) = \sum_{e \sim f} d_G(e) d_G(f)$$

where $e \sim f$ means that the edges e and f are adjacent.

More on these indices can be found in [28, 29, 30, 31, 32, 33, 34].

Motivated by the work on Zagreb and K -Banhatti indices, Kulli *et al.* introduced the Zagreb- K -Banhatti index of a graph G and defined as [35]

$$MB(G) = \sum_{\substack{a \text{ is either adjacent} \\ \text{or incident to } b}} d_G(a) + d_G(b)$$

where a and b are elements of G .

In [36], Kulli introduced the first and second hyper Zagreb- K -Banhatti indices of a graph G and they are defined as

$$HMB_1(G) = \sum_{\substack{a \text{ is either adjacent} \\ \text{or incident to } b}} [d_G(a) + d_G(b)]^2,$$

$$HMB_2(G) = \sum_{\substack{a \text{ is either adjacent} \\ \text{or incident to } b}} [d_G(a) d_G(b)]^2.$$

where a and b are elements of G .

Based on the successful consideration of Zagreb- K -Banhatti indices, we consider the second Zagreb- K -Banhatti index of a graph G as

$$MB_2(G) = \sum_{\substack{a \text{ is either adjacent} \\ \text{or incident to } b}} d_G(a) d_G(b)$$

where a and b are elements of G .

In this paper, we establish some relations between the second Zagreb- K -Banhatti index, Zagreb and K -Banhatti indices. We also provide lower and upper bounds for $MB_2(G)$ of a connected graph G in terms of Zagreb and K -Banhatti indices.

2. Comparison of Second Zagreb- K -Banhatti, Zagreb, K -Banhatti-type indices

Theorem 1: Let G be a graph with $n \geq 3$ vertices and m edges. Then

$$MB_2(G) = M_2(G) + EM_2(G) + B_2(G).$$

Proof: Let G be a graph with $n \geq 3$ vertices and m edges. Then

$$MB_2(G) = \sum_{\substack{a \text{ is either adjacent} \\ \text{or incident to } b}} d_G(a) d_G(b)$$

$$= \sum_{ab \in E(G)} d_G(a) d_G(b) + \sum_{e, f \in E(G), e \sim f} d_G(a) d_G(b) + \sum_{a(ab)} d_G(a) d_G(b)$$

$$= M_2(G) + EM_2(G) + B_2(G).$$

In order to prove our next result, we use the following result.

Theorem 2 [26]: Let G be a graph with $n \geq 3$ vertices and m edges. Then

$$B_2(G) = HM_1(G) - 2M_1(G).$$

Theorem 3: Let G be a graph with $n \geq 3$ vertices and m edges. Then

$$MB_2(G) = M_2(G) + EM_2(G) + HM_1(G) - 2M_1(G).$$

Proof: From Theorem 1, we have

$$MB_2(G) = M_2(G) + EM_2(G) + B_2(G).$$

Using Theorem 2, we obtain

$$MB_2(G) = M_2(G) + EM_2(G) + HM_1(G) - 2M_1(G).$$

We use the following result to establish our next result.

Theorem 4 [26]: Let G be a graph with $n \geq 3$ vertices and m edges. Then

$$B_2(G) = EM_1(G) + 2M_1(G) - 4m.$$

Theorem 5: Let G be a graph with $n \geq 3$ vertices and m edges. Then

$$MB_2(G) = M_2(G) + EM_1(G) + EM_2(G) + 2M_1(G) - 4m.$$

Proof: From Theorem 1, we have

$$MB_2(G) = M_2(G) + EM_2(G) + B_2(G).$$

From Theorems 4, we obtain

$$MB_2(G) = M_2(G) + EM_1(G) + EM_2(G) + 2M_1(G) - 4m.$$

We use the following result to prove Theorem 7.

Theorem 6 [26]: Let G be a graph with $n \geq 3$ vertices and m edges. Then

$$B_2(G) = F(G) + 2M_2(G) - 2M_1(G).$$

Theorem 7: Let G be a graph with $n \geq 3$ vertices and m edges. Then

$$MB_2(G) = 3M_2(G) + EM_2(G) + F(G) - 2M_1(G).$$

Proof: From Theorem 1, we have

$$MB_2(G) = M_2(G) + EM_2(G) + B_2(G).$$

Using Theorem 6, we get

$$MB_2(G) = 3M_2(G) + EM_2(G) + F(G) - 2M_1(G).$$

We use the following result to establish our next result.

Theorem 8 [26]: Let G be a graph with $n \geq 3$ vertices and m edges. Then

$$B_2(G) = \frac{1}{2}HB_1(G) - 12m.$$

Theorem 9: Let G be a graph with $n \geq 3$ vertices and m edges. Then

$$MB_2(G) = EM_1(G) + EM_2(G) + M_2(G) - M_1(G) + B_1(G).$$

Proof: From Theorem 1, we have

$$MB_2(G) = M_2(G) + EM_2(G) + B_2(G).$$

Thus by using Theorem 8, we get

$$MB_2(G) = M_2(G) + EM_2(G) + \frac{1}{2}HB_1(G) - 12m.$$

We have from [26],

$$HB_1(G) = 2B_1(G) + 2EM_1(G) - 2M_1(G) + 24m.$$

Using this result, we obtain

$$MB_2(G) = EM_1(G) + EM_2(G) + M_2(G) - M_1(G) + B_1(G).$$

3. Bounds on Second Zagreb-K-Banhatti, Zagreb, K-Banhatti-type indices

Theorem 10 [26]: For any connected graph G .

$$4M_2(G) - 2M_1(G) \leq B_2(G).$$

Theorem 11: Let G be a connected graph with $n \geq 3$ vertices and m edges. Then

$$5M_2(G) - 2M_1(G) + EM_2(G) \leq MB_2(G).$$

Proof: From Theorem 1, we have

$$MB_2(G) = M_2(G) + EM_2(G) + B_2(G).$$

From Theorem 10, we obtain

$$5M_2(G) - 2M_1(G) + EM_2(G) \leq MB_2(G).$$

To prove the next result, we use the earlier established:

Theorem 12 [34]: Let G be a graph with $n \geq 3$ vertices and m edges. Then

$$EM_2(G) \leq \left(\frac{1}{2}M_1(G) - m\right) \cdot \left(\sqrt{M_1(G) - 2m + \frac{1}{4} - \frac{1}{2}}\right)^2.$$

Theorem 13: Let G be a connected graph with $n \geq 3$ vertices and m edges. Then

$$MB_2(G) \leq F(G) + 3M_2(G) - 2M_1(G) + \left(\frac{1}{2}M_1(G) - m\right) \left(\sqrt{M_1(G) - 2m + \frac{1}{4} - \frac{1}{2}}\right)^2.$$

Proof: From Theorem 7, we have

$$MB_2(G) = 3M_2(G) - 2M_1(G) + EM_2(G) + F(G).$$

From Theorem 12, we get

$$MB_2(G) \leq F(G) + 3M_2(G) - 2M_1(G) + \left(\frac{1}{2}M_1(G) - m\right) \left(\sqrt{M_1(G) - 2m + \frac{1}{4} - \frac{1}{2}}\right)^2.$$

To prove our next result, we use the following result.

Theorem 14 [26]: For any graph G ,

$$M_1(G) \leq B_1(G).$$

Theorem 15: Let G be a graph with n vertices and m edges. Then

$$EM_1(G) + EM_2(G) + M_2(G) \leq MB_2(G).$$

Proof: From Theorem 9, we have

$$MB_2(G) = EM_1(G) + EM_2(G) + M_2(G) - M_1(G) + B_1(G).$$

Using Theorem 14, we get

$$EM_1(G) + EM_2(G) + M_2(G) \leq MB_2(G).$$

To prove the next result, we use the following result.

Theorem 16 [26]: Let G be a connected graph with $n \geq 3$ vertices and m edges. Then

$$B_2(G) \leq \frac{\delta(G) + \Delta(G)}{4m\delta(G)\Delta(G)} M_1(G)^2 - 2M_1(G).$$

Theorem 17: For any connected graph G with $n \geq 3$ vertices and m edges,

$$MB_2(G) \leq EM_2(G) + M_2(G) + \frac{\delta(G) + \Delta(G)}{4m\delta(G)\Delta(G)} M_1(G)^2 - 2M_1(G).$$

Proof: From Theorem 1, we have

$$MB_2(G) = EM_2(G) + M_2(G) + B_2(G).$$

Then from Theorem 16, we get the desired result.

We obtain lower and upper bounds on $MB_2(G)$ in terms of $\delta(G)$, $\Delta(G)$, $M_1(G)$, $M_2(G)$, m of a graph G .

Theorem 18: For any connected graph G with $n \geq 3$ vertices and m edges,

$$MB_2(G) \leq \left(\frac{1}{2}M_1(G) - m\right) \left(\sqrt{M_1(G) - 2m + \frac{1}{4} - \frac{1}{2}}\right)^2 + M_2(G) + \frac{\delta(G) + \Delta(G)}{4m\delta(G)\Delta(G)} M_1(G)^2 - 2M_1(G).$$

Proof: From Theorem 1, we have

$$MB_2(G) = EM_2(G) + M_2(G) + B_2(G).$$

Then from Theorems 12 and 16, we obtain

$$MB_2(G) \leq \left(\frac{1}{2}M_1(G) - m\right) \left(\sqrt{M_1(G) - 2m + \frac{1}{4} - \frac{1}{2}}\right)^2 + M_2(G) + \frac{\delta(G) + \Delta(G)}{4m\delta(G)\Delta(G)} M_1(G)^2 - 2M_1(G).$$

We use the following result to prove our next result.

Theorem 19 [26]: For any connected graph G with $n \geq 3$ vertices and m edges,

$$[\delta(G) - 2]M_1(G) + 2M_2(G) \leq B_2(G) \leq [\Delta(G) - 2]M_1(G) + 2M_2(G)$$

Further, equality in both lower and upper bounds will hold if and only if G is regular.

Theorem 20: For any connected graph G with $n \geq 3$ vertices and m edges,

$$[\delta(G) - 2]M_1(G) + 3M_2(G) + EM_2(G) \leq MB_2(G) \leq [\Delta(G) - 2]M_1(G) + 3M_2(G) + EM_2(G).$$

On both sides, equality holds if and only if G is regular.

Proof: From Theorem 1, we have

$$MB_2(G) = M_2(G) + EM_2(G) + B_2(G).$$

Using Theorem 19, we obtain

$$[\delta(G) - 2]M_1(G) + 3M_2(G) + EM_2(G) \leq MB_2(G) \leq [\Delta(G) - 2]M_1(G) + 3M_2(G) + EM_2(G).$$

Second part is obvious.

We use the following result, to establish our next result.

Theorem 21 [26]: For any connected graph G with $n \geq 3$ vertices and m edges,

$$4[\delta(G) - 2]^2 + 2M_1(G) - 4m \leq B_2(G) \leq [2M_1(G) - 4m]\Delta(G).$$

Theorem 22: For any connected graph G with $n \geq 3$ vertices and m edges,

$$4[\delta(G) - 1]^2 + 2M_1(G) - 4m + M_2(G) + EM_2(G) \leq MB_2(G) \leq [2M_1(G) - 4m]\Delta(G) + M_2(G) + EM_2(G).$$

Proof: From Theorem 1, we have

$$MB_2(G) = M_2(G) + EM_2(G) + B_2(G).$$

Therefore using Theorem 21, we get

$$4[\delta(G)-1]^2 + 2M_1(G) - 4m + M_2(G) + EM_2(G) \leq MB_2(G) \\ \leq [2M_1(G) - 4m]\Delta(G) + M_2(G) + EM_2(G).$$

The following result of $EM_2(G)$ to prove our next two results in terms of $\delta(G)$, $\Delta(G)$, $M_1(G)$, $M_2(G)$ and m of a graph G .

Theorem 23 [28]: Let G be a connected graph with n vertices and m edges. Then

$$[\delta(G)-1]^2 2(M_1(G) - 2m) \leq EB_2(G) \leq [\Delta(G)-1]^2 2(M_1(G) - 2m).$$

Theorem 24: Let G be a connected graph with $n \geq 3$ vertices and m edges. Then

$$[\delta(G)-2]M_1(G) + [\delta(G)-1]^2 2(M_1(G) - 2m) + 3M_2(G) \leq MB_2(G) \leq \\ [\Delta(G)-2]M_1(G) + [\Delta(G)-1]^2 2(M_1(G) - 2m) + 3M_2(G).$$

Proof: From Theorem 20, we have

$$[\delta(G)-2]M_1(G) + 3M_2(G) + EM_2(G) \leq MB_2(G) \leq [\Delta(G)-2]M_1(G) + 3M_2(G) + EM_2(G).$$

Using Theorem 23, we obtain

$$[\delta(G)-2]M_1(G) + [\delta(G)-1]^2 2(M_1(G) - 2m) + 3M_2(G) \leq MB_2(G) \leq \\ [\Delta(G)-2]M_1(G) + [\Delta(G)-1]^2 2(M_1(G) - 2m) + 3M_2(G).$$

Theorem 25: Let G be a connected graph with $n \geq 3$ vertices and m edges. Then

$$4[\delta(G)-1]^2 + [1 + (\delta(G)-1)^2] 2(M_1(G) - 2m) + M_2(G) \leq MB_2(G) \leq \\ [\Delta(G) + (\Delta(G)-1)^2] 2(M_1(G) - 2m) + M_2(G).$$

Proof: From Theorem 22, we have

$$4[\delta(G)-1]^2 + 2M_1(G) - 4m + M_2(G) + EM_2(G) \leq MB_2(G) \leq \\ [2M_1(G) - 4m]\Delta(G) + M_2(G) + EM_2(G).$$

Using Theorem 23, we obtain

$$4[\delta(G)-1]^2 + 2M_1(G) - 4m + M_2(G) + [\delta(G)-1]^2 2(M_1(G) - 2m) \leq MB_2(G) \leq \\ [2M_1(G) - 4m]\Delta(G) + M_2(G) + [\Delta(G)-1]^2 2(M_1(G) - 2m).$$

Therefore

$$4[\delta(G)-1]^2 + [1 + (\delta(G)-1)^2] 2(M_1(G) - 2m) + M_2(G) \leq MB_2(G) \leq \\ [\Delta(G) + (\Delta(G)-1)^2] 2(M_1(G) - 2m) + M_2(G).$$

We use the following result to establish our next result.

Theorem 26 [28]: Let G be a connected graph with n vertices and m edges. Then

$$[\delta(G)-1]^2 4m \leq EM_1(G) \leq [\Delta(G)-1]^2 4m.$$

Theorem 27: Let G be a connected graph with $n \geq 3$ vertices and m edges. Then

$$[(\delta(G)-1)^2 + 1] 2M_1(G) + M_2(G) - 4m \leq MB_2(G) \leq \\ [(\Delta(G)-1)^2 + 1] 2M_1(G) + M_2(G) - 4m.$$

Proof: From Theorems 22 and 26, we obtain

$$[\delta(G)-1]^2 4m + [\delta(G)-1]^2 (2M_1(G) - 4m) \leq EM_1(G) + EM_2(G) \leq \\ [\Delta(G)-1]^2 4m + [\Delta(G)-1]^2 (2M_1(G) - 4m).$$

$$\text{Therefore } [\delta(G)-1]^2 2M_1(G) \leq EM_1(G) + EM_2(G) \leq [\Delta(G)-1]^2 2M_1(G). \quad (1)$$

From Theorem 5, we have

$$MB_2(G) = M_2(G) + 2M_1(G) - 4m + EM_1(G) + EM_2(G).$$

Then from inequality (1), we obtain

$$\begin{aligned} [\delta(G)-1]^2 2M_1(G) + 2M_1(G) - 4m &\leq MB_2(G) \leq \\ [\Delta(G)-1]^2 2M_1(G) + M_2(G) + 2M_1(G) - 4m. \end{aligned}$$

Therefore

$$\begin{aligned} [(\delta(G)-1)^2 + 1] 2M_1(G) + M_2(G) - 4m &\leq MB_2(G) \leq \\ [(\Delta(G)-1)^2 + 1] 2M_1(G) + M_2(G) - 4m. \end{aligned}$$

Theorem 28: Let G be a connected graph with $n \geq 3$ vertices and m edges. Then

$$\begin{aligned} [\delta(G)-1]^2 2M_1(G) + M_2(G) - M_1(G) + B_1(G) &\leq MB_2(G) \leq \\ [\Delta(G)-1]^2 2M_1(G) + M_2(G) - M_1(G) + B_1(G). \end{aligned}$$

Proof: From Theorems 22 and 26, we obtain

$$MB_2(G) = EM_1(G) + EM_2(G) + M_2(G) - M_1(G) + B_1(G).$$

Then from inequality (1), we get

$$\begin{aligned} [\delta(G)-1]^2 2M_1(G) + M_2(G) - M_1(G) + B_1(G) &\leq MB_2(G) \leq \\ [\Delta(G)-1]^2 2M_1(G) + M_2(G) - M_1(G) + B_1(G). \end{aligned}$$

Theorem 29: Let G be a connected graph with $n \geq 3$ vertices and m edges. Then

$$\begin{aligned} [\delta(G)-1]^2 (2M_1(G) - 4m) + 3M_2(G) - 2M_1(G) + F(G) &\leq MB_2(G) \leq \\ [\Delta(G)-1]^2 (2M_1(G) - 4m) + 3M_2(G) - 2M_1(G) + F(G). \end{aligned}$$

Proof: From Theorem 7, we have

$$MB_2(G) = EM_2(G) + 3M_2(G) - 2M_1(G) + F(G).$$

Then from Theorem 22, we obtain

$$\begin{aligned} [\delta(G)-1]^2 (2M_1(G) - 4m) + 3M_2(G) - 2M_1(G) + F(G) &\leq MB_2(G) \leq \\ [\Delta(G)-1]^2 (2M_1(G) - 4m) + 3M_2(G) - 2M_1(G) + F(G). \end{aligned}$$

To prove the next two results, we use the earlier established:

Theorem 30 [28]: Let G be a connected graph with n vertices and m edges. Then

$$[\delta(G)-1]EM_1(G) \leq EM_2(G) \leq [\Delta(G)-1]EM_1(G).$$

Theorem 31: Let G be a connected graph with $n \geq 3$ vertices and m edges. Then

$$\delta(G)EM_1(G) + M_2(G) + 2M_1(G) - 4m \leq MB_2(G) \leq \Delta(G)EM_1(G) + M_2(G) + 2M_1(G) - 4m.$$

Proof: From Theorem 5, we have

$$MB_2(G) = EM_1(G) + EM_2(G) + M_2(G) + 2M_1(G) - 4m.$$

Using Theorem 30, we obtain

$$\delta(G)EM_1(G) + M_2(G) + 2M_1(G) - 4m \leq MB_2(G) \leq \Delta(G)EM_1(G) + M_2(G) + 2M_1(G) - 4m.$$

Theorem 32: Let G be a connected graph with $n \geq 3$ vertices and m edges. Then

$$\delta(G)EM_1(G) + M_2(G) - M_1(G) + B_1(G) \leq MB_2(G) \leq \Delta(G)EM_1(G) + M_2(G) - M_1(G) + B_1(G).$$

Proof: From Theorem 9, we have

$$MB_2(G) = EM_1(G) + EM_2(G) + M_2(G) - M_1(G) + B_1(G).$$

Using Theorem 30, we get

$$\delta(G)EM_1(G) + M_2(G) - M_1(G) + B_1(G) \leq MB_2(G) \leq \Delta(G)EM_1(G) + M_2(G) - M_1(G) + B_1(G).$$

CONCLUSION

In this paper, we have established some relations between second Zagreb-K-Banhatti index with Zagreb and K-Banhatti indices. We also provided lower and upper bounds for $MB_2(G)$ of a connected graph G in terms of other topological indices.

REFERENCES

1. V.R.Kulli, Graph indices, in *Hand Book of Research on Advanced Applications of Application Graph Theory in Modern Society*, M. Pal. S. Samanta and A. Pal (eds.) IGI Global, USA (2020) 66-91.
2. I.Gutman and O.E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer, Berlin (1986).
3. V.R.Kulli, *Multiplicative Connectivity Indices of Nanostructures*, LAP LEMBERT Academic Publishing (2018).
4. R.Todeschini and V. Consonni, *Molecular Descriptors for Chemoinformatics*, Wiley-VCH, Weinheim, (2009).
5. V.R.Kulli, *College Graph Theory*, Vishwa International Publications, Gulbarga, India (2012).
6. I.Gutman and N. Trinajstić, Graph theory and molecular orbitals. Total π -electron energy of alternant hydrocarbons, *Chem. Phys. Lett.* 17 (1972) 535-538.
7. K.C.Das, I. Gutman and B. Zhou, New upper bounds on Zagreb indices, *J. Math. Chem.* 46(2) (2009) 514-521.
8. M.H.Khalifeh, H. Yousefi-Azari and A.R. Ashrafi, The first and second Zagreb indices of some graph operations, *Discrete Appl. Math.* 157(4) (2009) 804-811.
9. P.S.Ranjini, V. Loksha and I.N. Cangül, On the Zagreb indices of the line graphs of the subdivision graphs. *Applied Mathematics and Computation* 218(3) (2011) 699-702.
10. T.Retí, On the relationships between the first and second Zagreb indices, *MATCH commun. Math. Comput. Chem.* 68 (2012) 169-188.
11. K.Xu, K. Tang, H. Liu and J. Wang, The Zagreb indices of bipartite graphs with more edges, *J. Appl. Math. and Informatics*, 33(3) (2015) 365-377.
12. B.Furtula and I. Gutman, A forgotten topological index, *J. Math. Chem.* 53 (2015), 1184-1190.
13. B.Basavanagoud, V.R.Desai, Forgotten topological index and hyper Zagreb index of generalized transformation graphs, *Bull Math. Sci. Appl.* 14 (2016) 1-6.
14. I.Gutman, A Ghalavand, T. Dehghan-Zadeh and A.R. Ashrafi, Graphs with smallest forgotten index, *Iran. J. Math. Chem.* 8 (2017) 259-273.
15. H. S.Ramane, R.B. Jummannaver, Note on forgotten topological index of chemical structure in drugs, *Applied Mathematics and Nonlinear Sciences*, 1(2) (2016) 369-374. doi.10.21042/AMNS. 2016.2.00032
16. V.R.Kulli, On K Banhatti indices of graphs, *Journal of Computer and Mathematical Sciences*, 7 (2016) 213-218.
17. V.R.Kulli, On K hyper Banhatti indices and coindices of graphs, *International Research Journal of Pure Algebra*, 6(5) (2016) 300-304.
18. A.Asghar, M. Rafaqat, W. Nazeer and W. Gao, K Banhatti and K hyper Banhatti indices of circulant graphs, *International Journal of Advanced and Applied Sciences*, 5(5) (2018) 107-109.
19. F.Dayan. M. Javaid, M. Zulqarnain, M.T. Ali and B. Ahmad, Computing Banhatti indices of hexagonal, honeycomb and derived graphs, *American Journal of Mathematical and Computer Modeling*, 3(2) (2018) 38-45.
20. W.Gao, B. Muzaffer and W. Nazeer, K. Banhatti and K hyper Banhatti indices of dominating David derived network, *Open J. Math. Anal.* 1(1) (2019) 13-24.
21. V.R.Kulli, Computing Banhatti indices of networks, *International Journal of Advances in Mathematics*, 2018(1) 2018) 31-40.
22. V.R.Kulli, K Banhatti indices of chloroquine and hydroxychloroquine: Research Applied for the treatment and prevention of COVID-19, *SSRG International Journal of Applied Chemistry*, 7(1) (2020) 63-68.
23. V.R.Kulli, K Banhatti polynomials of remdesivir, chloroquine and hydroxychloroquine: Research Advances for the prevention and treatment of COVID-19, *SSRG International Journal of Applied Chemistry*, 7(2) (2020) 48-55.

24. V.R.Kulli, B.Chaluvaraju and H.S. Baregowda, K -Banhatti and K hyper-Banhatti indices of windmill graphs, *South East Asian J. of Math. and Math. Sci*, 13(1) (2017) 11-18.
25. V.R.Kulli, B.Chaluvaraju and H.S.Baregowda, Some bounds of sum connectivity Banhatti index of graphs, *Palestine Journal of Mathematics*, 8(2) (2019) 355-364.
26. I. Gutman, V.R. Kulli, B. Chaluvaraju and H. S. Boregowda, On Banhatti and Zagreb indices, *Journal of the International Mathematical Virtual Institute*, 7 (2017) 53-67.
27. A.Milićević, S. Nikolić and N. Trinajstić, On reformulated Zagreb indices, *Molecular Diversity*, 8, (2004) 393-399.
28. N.De, Some bounds of reformulated Zagreb indices, *Appl. Mathematical Sciences*, 6 (2012) 5005-5012.
29. A.Ilić and B. Zhou, On reformulated Zagreb indices, *Discrete Appl. Math.* 160 (2012) 204-209.
30. V.R.Kulli, F -index and reformulated Zagreb index of certain nanostructures, *International Research Journal of Pure Algebra*, 7(1) (2017) 489-495.
31. T.Mansour, M.A. Rostami, E. Suresh and G.B.A. Xavier, On the bounds of the first reformulated Zagreb index, *Turkish Journal of Analysis and Number Theory*, 4(1) (2016) 8-15.
32. P.S.Ranjini, V. Loksha, M Bindusree and M. Phani Raju, New bounds on Zagreb indices and the Zagreb coindices, *Bol. Soc. Paran. Mat.* 31(1) (2013) 51-55.
33. A.Selvakumar and K.Agilarasan, Bounds on general Zagreb indices, *International Journal of Mathematics Trends and Technology*, 56(7) (2018) 524-526.
34. B.Zhou and N. Trinajstić, Some properties of the reformulated Zagreb indices, *J. Math. Chem.* 48 (2010) 714-719.
35. V.R.Kulli and B. Chaluvaraju, Zagreb-K-Banhatti index of a graph, *Journal of Ultra Scientist of Physical Sciences – A*, 32(5) (2020) 29-36.
36. V.R.Kulli, Hyper Zagreb-K-Banhatti indices of a graphs, *International Journal of Mathematics Trends and Technology*, 66(8) (2020) 123-130.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2020. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]