

ON A COMMON FIXED POINT RESULT IN INTUITIONISTIC FUZZY METRIC SPACE

VED PRAKASH BHARDWAJ^{1*}, KAILASH NAMDEO² AND KAMAL WADHWA³

^{1*}Department of Mathematics,
Govt. Arts, Commerce and Science College Sukhtawa,
Hoshangabad (M.P.), India 461001.

²Govt. P.G. College Gadawada, Narshingpur, (M.P.), India.

³Govt. P.G. College Pipariya, Hoshangabad (M.P.), India 461001.

(Received On: 30-11-20; Revised & Accepted On: 28-12-20)

ABSTRACT

In this paper, we prove a common fixed point theorem for new type of common limit range property under functionally rationalized contractive condition in intuitionistic fuzzy metric space.

2010 Mathematics Subject Classification: 47H10, 54H25.

Keywords: Intuitionistic fuzzy metric space (IFMS), weakly compatible mapping, new type of common limit in the range property.

1. INTRODUCTION

Kramosil and Michalek [10] introduced the notion of fuzzy metric space using the idea of [17], which opened the way for further development of analysis. IFMS is defined by Alaca *et al.* [3] using continuous t-norms and t-conorms with the help of intuitionistic fuzzy set as a generalization of fuzzy metric space, as defined by Kramosil and Michalek [10]. The consequence of fixed point theory is marked from the fact that it has its applications in different disciplines of Science, Engineering, and Economics in dealing with problems arising in: Approximation theory, potential theory, game theory, mathematical economics, etc. Intuitionistic fuzzy fixed point theory has turn into a topic of enormous attention for expert in fixed point theory for the reason that this branch of mathematics has cover novel possibilities for fixed point theorists. Some recent important results using IFMS are [5, 8, 14].

There is a diversity of different mappings used by different authors to take a range of fixed point theorems in various fuzzy spaces. Aamri and Moutawakil [1] gave the idea of property E.A. for a pair of self mappings which contains the class of non-compatible mappings. On the other hand, the notion of common limit range property is given by Sintunavarat and Kumam [13] as an improvement of property E.A., which relaxes the condition of closedness of the underlying subspace. Recently, Popa and Patriciu [11] introduced a new type of common limit in the range property in metric space and proved some fixed point results. Some recent important results using common limit in the range property are [4, 5, 6, 7, 15, 16].

The aim of this paper is to prove a common fixed point result in IFMS under functionally rationalized contractive condition using new type of common limit range property motivated by [11].

2. PRELIMINARIES

In this section, we have recalled some definitions and useful results which are already in the literature.

Definition 2.1 [12]: A binary operation $*$: $[0, 1] \times [0, 1] \rightarrow [0, 1]$ is called a continuous t-norm if $([0, 1], *)$ is an abelian topological monoid with the unit 1 such that $a * b \leq c * d$, whenever $a \leq c$ and $b \leq d$, $\forall a, b, c, d \in [0, 1]$.

Corresponding Author: Ved Prakash Bhardwaj^{1},*

^{1}Department of Mathematics, Govt. Arts, Commerce and Science College Sukhtawa,
Hoshangabad (M.P.), India 461001.*

Definition 2.2 [12]: A binary operation $\diamond : [0, 1] \times [0, 1] \rightarrow [0, 1]$ is continuous t-conorm if \diamond satisfies the following conditions: (i) \diamond is commutative and associative; (ii) \diamond is continuous; (iii) $a \diamond 0 = a$ for all $a \in [0, 1]$; (iv) $a \diamond b \leq c \diamond d$ whenever $a \leq c$ and $b \leq d$ for all $a, b, c, d \in [0, 1]$.

Alaca et al. [3] defined the notion of IFMS as follows:

Definition 2.3 [3]: A 5-tuple $(X, M, N, *, \diamond)$ is said to be an intuitionistic fuzzy metric space if X is an arbitrary set, $*$ is a continuous t-norm, \diamond is a continuous t-conorm and M, N are fuzzy sets on $X^2 \times [0, \infty)$ satisfying

- (i) $M(x, y, t) + N(x, y, t) \leq 1 \quad \forall x, y \in X$ and $t > 0$;
- (ii) $M(x, y, 0) = 0 \quad \forall x, y \in X$;
- (iii) $M(x, y, t) = 1 \quad \forall x, y \in X$ and $t > 0$ if and only if $x = y$;
- (iv) $M(x, y, t) = M(y, x, t)$ for all $x, y \in X$ and $t > 0$;
- (v) $M(x, y, t) * M(y, z, s) \leq M(x, z, t + s) \quad \forall x, y, z \in X$ and $s, t > 0$;
- (vi) $\forall x, y \in X, M(x, y, \cdot) : [0, \infty) \rightarrow [0, 1]$ is left continuous;
- (vii) $\lim_{t \rightarrow \infty} M(x, y, t) = 1 \quad \forall x, y \in X$ and $t > 0$;
- (viii) $N(x, y, 0) = 1 \quad \forall x, y \in X$;
- (ix) $N(x, y, t) = 0 \quad \forall x, y \in X$ and $t > 0$ if and only if $x = y$;
- (x) $N(x, y, t) = N(y, x, t)$ for all $x, y \in X$ and $t > 0$;
- (xi) $N(x, y, t) \diamond N(y, z, s) \geq N(x, z, t + s) \quad \forall x, y, z \in X$ and $s, t > 0$;
- (xii) $N(x, y, \cdot) : [0, \infty) \rightarrow [0, 1]$ is right continuous;
- (xiii) $\lim_{t \rightarrow \infty} N(x, y, t) = 0 \quad \forall x, y \in X$

Then (M, N) is called an intuitionistic fuzzy metric on X . The functions $M(x, y, t)$ and $N(x, y, t)$ denote the degree of nearness and the degree of nonnearness between x and y with respect to t , respectively.

Remark 2.4 [3]. Every fuzzy metric space $(X, M, *)$ is an IFMS of the form $(X, M, 1 - M, *, \diamond)$ such that t-norm $*$ and t-conorm \diamond are associated as $x \diamond y = 1 - ((1 - x) * (1 - y)) \quad \forall x, y \in X$.

Remark 2.5 [3]. In an IFMS $(X, M, N, *, \diamond)$, $M(x, y, \cdot)$ is non-decreasing and $N(x, y, \cdot)$ is non-increasing $\forall x, y \in X$.

Definition 2.6 [9]: Two self-maps P and Q on set X are said to be weakly compatible if they commute at their coincident point.

Definition 2.7 [1]: A pair of self mapping (f, g) on X said to satisfy the property (E.A.) if there exist a sequence x_n such that

$$\lim_{n \rightarrow \infty} f x_n = \lim_{n \rightarrow \infty} g x_n = t,$$

for some $t \in X$.

Definition 2.8 [13]: A pair (f, g) of self-mappings of a fuzzy metric space $(X, M, *)$ is said to satisfy the common limit in the range property with respect to mapping g (briefly CLR_g) property, if there exists a sequence x_n in X such that

$$\lim_{n \rightarrow \infty} f x_n = \lim_{n \rightarrow \infty} g x_n = g z, \text{ for some } z \in X.$$

Recently, Popa and Patriciu [11] introduced a new type of common limit range property for self mappings in metric space.

Definition 2.9 [11]: Let A, S and T be self mappings of a metric space (X, d) . The pair (A, S) is said to satisfy common limit range property with respect to T (shortly CLR_{(A,S)T} property), if there exists a sequence x_n in X such that

$$\lim_{n \rightarrow \infty} A x_n = \lim_{n \rightarrow \infty} S x_n = t, \text{ for some } t \in S(X) \cap T(X).$$

Lemma 2.10 [2]: Let $(X, M, N, *, \diamond)$ be an IFMS and if for a number $k \in (0, 1)$,

$$M(x, y, kt) \geq M(x, y, t) \text{ and } N(x, y, kt) \leq N(x, y, t)$$

for all $x, y \in X, t > 0$, then $x = y$.

Lemma 2.11 [11]: Let f, g be two weakly compatible self mappings of a nonempty set X . If f and g have a unique point of coincidence $w = fx = gx$ for some $x \in X$, then w is the unique common fixed point of f and g .

We use the following relations in our results:

(δ) $\alpha_1, \alpha_2, \beta_1, \beta_2, \gamma_1, \gamma_2 : [0, 1] \rightarrow [0, 1]$ are continuous functions such that $\alpha_1 \& \alpha_2, \beta_1 \& \beta_2$ and $\gamma_1 \& \gamma_2$ cannot be simultaneously 0, and for $h \in [0, 1]$

$$\alpha_1(h) + \alpha_2(h) = 1, \beta_1(h) + \beta_2(h) = 1, \gamma_1(h) + \gamma_2(h) = 1.$$

(δ^*) Let Φ be the set of all real valued continuous function $\phi, \phi' : [0, \infty)^5 \rightarrow [0, \infty)$ such that

($\delta_{2.1}$) ϕ is non increasing and ϕ' is non decreasing,

$(\delta_{2.2})$ for all $s, s' \geq 0$,

$$\begin{aligned}\phi(1, 1, s, 1, s) &\geq s, & \phi'(1, 1, s', 1, s') &\leq s' \text{ and} \\ \phi(1, s, 1, s, 1) &\geq s, & \phi'(1, s', 1, s', 1) &\leq s' \text{ and} \\ \phi(s, 1, 1, s, s) &\geq s, & \phi'(s', 1, 1, s', s') &\leq s'.\end{aligned}$$

Motivated from Definition 2.9, we can have following definition in intuitionistic fuzzy metric space:

Definition 2.12: Let A, S and T be self mappings of an intuitionistic fuzzy metric space (IFMS) $(X, M, N, *, \phi)$. The pair (A, S) is said to satisfy common limit range property with respect to T (shortly $CLR_{(A,S)T}$ property), if there exists a sequence x_n in X such that

$$\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Sx_n = x,$$

i.e.,

$$\lim_{n \rightarrow \infty} M(Ax_n, x, t) = \lim_{n \rightarrow \infty} M(Sx_n, x, t) = 1$$

and

$$\lim_{n \rightarrow \infty} N(Ax_n, x, t) = \lim_{n \rightarrow \infty} N(Sx_n, x, t) = 0,$$

for some $x \in S(X) \cap T(X)$.

Now, we prove our main result.

3. MAIN RESULT

Theorem 3.1: Let $(X, M, N, *, \phi)$ be a IFMS with continuous t -norm and t -conorm. Let A, B, S and T be self-mappings from $(X, M, N, *, \phi)$ into itself satisfying:

(3.1) for all $x, y \in X, t > 0$ and for a number $k \in (0, 1)$,

$$M(Ax, By, kt) \geq \phi \left(\frac{\alpha_2(h)M(Ty, Ax, t) + \alpha_1(h)M(Ty, Sx, t)}{\alpha_1(h) + \alpha_2(h)M(Ax, Sx, t)}, \frac{M(Ax, Sx, t), M(By, Ty, t), \beta_2(h)M(Ax, By, t) + \beta_1(h)M(Ax, Ty, t)}{\beta_1(h) + \beta_2(h)M(By, Ty, t)}, \frac{\gamma_2(h)M(By, Ax, t) + \gamma_1(h)M(By, Sx, t)}{\gamma_1(h) + \gamma_2(h)M(Ax, Sx, t)} \right),$$

and

$$N(Ax, By, kt) \leq \phi' \left(\frac{\alpha_2(h)N(Ty, Ax, t) + \alpha_1(h)N(Ty, Sx, t)}{\alpha_1(h) + \alpha_2(h)N(Ax, Sx, t)}, \frac{N(Ax, Sx, t), N(By, Ty, t), \beta_2(h)M(Ax, By, t) + \beta_1(h)N(Ax, Ty, t)}{\beta_1(h) + \beta_2(h)N(By, Ty, t)}, \frac{\gamma_2(h)N(By, Ax, t) + \gamma_1(h)N(By, Sx, t)}{\gamma_1(h) + \gamma_2(h)N(Ax, Sx, t)} \right),$$

where $\phi, \phi' \in \Phi$ and $\alpha_1, \alpha_2, \beta_1, \beta_2, \gamma_1, \gamma_2$ are defined as above and if (A, S) and T enjoys the $CLR_{(A,S)T}$ property, then $C(A, S) \neq \emptyset$ and $C(B, T) \neq \emptyset$.

Moreover, if (A, S) and (B, T) are weakly compatible then A, B, S and T have a unique common fixed point in X .

Proof: Since (A, S) and T enjoys $CLR_{(A,S)T}$ property, then there exists a sequence x_n in X such that

$$\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Sx_n = z, \text{ for some } z \in S(X) \cap T(X).$$

Since $z \in T(X)$, there exists $u \in X$ such that $z = Tu$.

Using (3.1), we have

$$M(Ax_n, Bu, kt) \geq \phi \left(\frac{\alpha_2(h)M(Tu, Ax_n, t) + \alpha_1(h)M(Tu, Sx_n, t)}{\alpha_1(h) + \alpha_2(h)M(Ax_n, Sx_n, t)}, \frac{M(Ax_n, Sx_n, t), M(Bu, Tu, t), \beta_2(h)M(Ax_n, Bu, t) + \beta_1(h)M(Ax_n, Tu, t)}{\beta_1(h) + \beta_2(h)M(Bu, Tu, t)}, \frac{\gamma_2(h)M(Bu, Ax_n, t) + \gamma_1(h)M(Bu, Sx_n, t)}{\gamma_1(h) + \gamma_2(h)M(Ax_n, Sx_n, t)} \right).$$

Take the limit as $n \rightarrow \infty$, we get,

$$M(z, Bu, kt) \geq \phi \left(\frac{\alpha_2(h)M(z, z, t) + \alpha_1(h)M(z, z, t)}{\alpha_1(h) + \alpha_2(h)M(z, z, t)}, \frac{M(z, z, t), M(Bu, z, t), \beta_2(h)M(z, Bu, t) + \beta_1(h)M(z, z, t)}{\beta_1(h) + \beta_2(h)M(Bu, z, t)}, \frac{\gamma_2(h)M(Bu, z, t) + \gamma_1(h)M(Bu, z, t)}{\gamma_1(h) + \gamma_2(h)M(z, z, t)} \right).$$

$$M(z, Bu, kt) \geq \phi(1, 1, M(Bu, z, t), 1, M(Bu, z, t)).$$

Using $(\delta_{2.2})$, we have,

$$M(Bu, z, kt) \geq M(Bu, z, t) \tag{1}$$

and

$$N(Ax_n, Bu, kt) \leq \phi' \left(\frac{\alpha_2(h)N(Tu, Ax_n, t) + \alpha_1(h)N(Tu, Sx_n, t)}{\alpha_1(h) + \alpha_2(h)N(Ax_n, Sx_n, t)}, \frac{N(Ax_n, Sx_n, t), N(Bu, Tu, t), \beta_2(h)N(Ax_n, Bu, t) + \beta_1(h)N(Ax_n, Tu, t)}{\beta_1(h) + \beta_2(h)N(Bu, Tu, t)}, \frac{\gamma_2(h)N(Bu, Ax_n, t) + \gamma_1(h)N(Bu, Sx_n, t)}{\gamma_1(h) + \gamma_2(h)N(Ax_n, Sx_n, t)} \right).$$

Take the limit as $n \rightarrow \infty$, we get,

$$N(z, Bu, kt) \leq \phi' \left(\frac{\alpha_2(h)N(z, z, t) + \alpha_1(h)N(z, z, t)}{\alpha_1(h) + \alpha_2(h)N(z, z, t)}, \frac{N(z, z, t), N(Bu, z, t), \beta_2(h)N(z, Bu, t) + \beta_1(h)N(z, z, t)}{\beta_1(h) + \beta_2(h)M(Bu, z, t)}, \frac{\gamma_2(h)N(Bu, z, t) + \gamma_1(h)N(Bu, z, t)}{\gamma_1(h) + \gamma_2(h)N(z, z, t)} \right).$$

$$N(z, Bu, kt) \leq \phi'(1, 1, N(Bu, z, t), 1, N(Bu, z, t)).$$

Using $(\delta'_{2.2})$, we have,

$$N(Bu, z, kt) \leq N(Bu, z, t) \tag{2}$$

Using (1), (2) and Lemma 2.10, we have, $Bu = z = Tu$. Therefore $C(B, T) \neq \emptyset$.

Since $z \in S(X)$, there exists $v \in X$ such that $z = Sv$. Using (3.1), we have

$$M(Av, Bu, kt) \geq \phi \left(\frac{\alpha_2(h)M(Tu, Av, t) + \alpha_1(h)M(Tu, Sv, t)}{\alpha_1(h) + \alpha_2(h)M(Av, Sv, t)}, \frac{M(Av, Sv, t), M(Bu, Tu, t), \beta_2(h)M(Av, Bu, t) + \beta_1(h)M(Av, Tu, t)}{\beta_1(h) + \beta_2(h)M(Bu, Tu, t)}, \frac{\gamma_2(h)M(Bu, Av, t) + \gamma_1(h)M(Bu, Sv, t)}{\gamma_1(h) + \gamma_2(h)M(Av, Sv, t)} \right).$$

$$M(Av, z, kt) \geq \phi \left(\frac{\alpha_2(h)M(z, Av, t) + \alpha_1(h)M(z, z, t)}{\alpha_1(h) + \alpha_2(h)M(Av, z, t)}, \frac{M(Av, z, t), M(z, z, t), \beta_2(h)M(Av, z, t) + \beta_1(h)M(Av, z, t)}{\beta_1(h) + \beta_2(h)M(z, z, t)}, \frac{\gamma_2(h)M(z, Av, t) + \gamma_1(h)M(z, z, t)}{\gamma_1(h) + \gamma_2(h)M(Av, z, t)} \right).$$

$$M(Av, z, kt) \geq \phi(1, M(Av, z, t), 1, M(Av, z, t), 1).$$

Using $(\delta_{2.2})$, we have,

$$M(Av, z, kt) \geq M(Av, z, t) \quad (3)$$

and

$$N(Av, Bu, kt) \leq \phi' \left(\frac{\alpha_2(h)N(Tu, Av, t) + \alpha_1(h)N(Tu, Sv, t)}{\alpha_1(h) + \alpha_2(h)N(Av, Sv, t)}, \frac{N(Av, Sv, t), N(Bu, Tu, t), \beta_2(h)N(Av, Bu, t) + \beta_1(h)N(Av, Tu, t)}{\beta_1(h) + \beta_2(h)N(Bu, Tu, t)}, \frac{\gamma_2(h)N(Bu, Av, t) + \gamma_1(h)N(Bu, Sv, t)}{\gamma_1(h) + \gamma_2(h)N(Av, Sv, t)} \right).$$

$$N(Av, z, kt) \leq \phi' \left(\frac{\alpha_2(h)N(z, Av, t) + \alpha_1(h)N(z, z, t)}{\alpha_1(h) + \alpha_2(h)N(Av, z, t)}, \frac{N(Av, z, t), N(z, z, t), \beta_2(h)N(Av, z, t) + \beta_1(h)N(Av, z, t)}{\beta_1(h) + \beta_2(h)N(z, z, t)}, \frac{\gamma_2(h)N(z, Av, t) + \gamma_1(h)N(z, z, t)}{\gamma_1(h) + \gamma_2(h)N(Av, z, t)} \right).$$

$$N(Av, z, kt) \leq \phi'(1, N(Av, z, t), 1, N(Av, z, t), 1).$$

Using $(\delta'_{2.2})$, we have,

$$N(Av, z, kt) \leq N(Av, z, t) \quad (4)$$

Using (3), (4) and Lemma 2.10, we have, $Av = z = Sv$. Therefore $C(A, S) \neq \emptyset$.

Hence, $z = Av = Sv = Bu = Tu$ and z is a point of coincidence of (A, S) and of (B, T) .

We show that z is the unique point of coincidence of (A, S) and (B, T) .

Let p be another point of coincidence of (A, S) , i.e., $p = Aw = Sw$. Using (3.1), we get

$$M(Aw, Bu, kt) \geq \phi \left(\frac{\alpha_2(h)M(Tu, Aw, t) + \alpha_1(h)M(Tu, Sw, t)}{\alpha_1(h) + \alpha_2(h)M(Aw, Sw, t)}, \frac{M(Aw, Sw, t), M(Bu, Tu, t), \beta_2(h)M(Aw, Bu, t) + \beta_1(h)M(Aw, Tu, t)}{\beta_1(h) + \beta_2(h)M(Bu, Tu, t)}, \frac{\gamma_2(h)M(Bu, Aw, t) + \gamma_1(h)M(Bu, Sw, t)}{\gamma_1(h) + \gamma_2(h)M(Aw, Sw, t)} \right),$$

$$M(p, z, kt) \geq \phi \left(\frac{\alpha_2(h)M(z, p, t) + \alpha_1(h)M(z, p, t)}{\alpha_1(h) + \alpha_2(h)M(p, p, t)}, \frac{M(p, p, t), M(p, p, t), \beta_2(h)M(p, z, t) + \beta_1(h)M(p, z, t)}{\beta_1(h) + \beta_2(h)M(z, z, t)}, \frac{\gamma_2(h)M(z, p, t) + \gamma_1(h)M(z, p, t)}{\gamma_1(h) + \gamma_2(h)M(p, p, t)} \right),$$

$$M(p, z, kt) \geq \phi(M(z, p, t), 1, 1, M(p, z, t), M(p, z, t)),$$

Using $(\delta_{2.2})$, we have,

$$M(p, z, kt) \geq M(p, z, t) \quad (5)$$

and

$$N(Aw, Bu, kt) \leq \phi' \left(\frac{\alpha_2(h)N(Tu, Aw, t) + \alpha_1(h)N(Tu, Sw, t)}{\alpha_1(h) + \alpha_2(h)N(Aw, Sw, t)}, \frac{N(Aw, Sw, t), N(Bu, Tu, t), \beta_2(h)N(Aw, Bu, t) + \beta_1(h)N(Aw, Tu, t)}{\beta_1(h) + \beta_2(h)N(Bu, Tu, t)}, \frac{\gamma_2(h)N(Bu, Aw, t) + \gamma_1(h)N(Bu, Sw, t)}{\gamma_1(h) + \gamma_2(h)N(Aw, Sw, t)} \right),$$

$$N(p, z, kt) \leq \phi' \left(\frac{\alpha_2(h)N(z, p, t) + \alpha_1(h)N(z, p, t)}{\alpha_1(h) + \alpha_2(h)N(p, p, t)}, \frac{N(p, p, t), N(p, p, t), \beta_2(h)N(p, z, t) + \beta_1(h)N(p, z, t)}{\beta_1(h) + \beta_2(h)N(z, z, t)}, \frac{\gamma_2(h)N(z, p, t) + \gamma_1(h)N(z, p, t)}{\gamma_1(h) + \gamma_2(h)N(p, p, t)} \right),$$

$$N(p, z, kt) \leq \phi'(N(z, p, t), 1, 1, N(p, z, t), N(p, z, t)),$$

Using $(\delta'_{2.2})$, we have,

$$N(p, z, kt) \leq N(p, z, t) \quad (6)$$

Using (5), (6) and Lemma 2.10, we have, $p = z$. Thus, z is the unique point of coincidence of (A, S) . Similarly, using (3.1) it is easy to see that z is the unique point of coincidence of (B, T) .

Further, if (A, S) and (B, T) are weakly compatible, then by Lemma 2.11, z is the unique common fixed point of A, B, S and T . This completes the proof of the theorem.

If we take $\alpha_2(h) = \beta_2(h) = \gamma_2(h) = 0$, for all $h \in [0, 1]$ in Theorem 3.1, then we have following:

Corollary 3.2: Let $(X, M, N, *, \phi)$ be a IFMS with continuous t - norm and t -conorm. Let A, B, S and T be self-mappings from $(X, M, N, *, \phi)$ into itself satisfying:

(3.2) for all $x, y \in X$, $t > 0$ and for a number $k \in (0, 1)$,

$$M(Ax, By, kt) \geq \phi \left(\frac{M(Ty, Sx, t), M(Ax, Sx, t)}{M(By, Ty, t), M(Ax, Ty, t), M(By, Sx, t)} \right),$$

and

$$N(Ax, By, kt) \geq \phi' \left(\frac{N(Ty, Sx, t), N(Ax, Sx, t)}{N(By, Ty, t), N(Ax, Ty, t), N(By, Sx, t)} \right),$$

where $\phi, \phi' \in \Phi$ and $\alpha_1, \alpha_2, \beta_1, \beta_2, \gamma_1, \gamma_2$ are defined above and if (A, S) and T enjoys the $CLR_{(A,S)T}$ property, then $C(A, S) \neq \emptyset$ and $C(B, T) \neq \emptyset$.

Moreover, if (A, S) and (B, T) are weakly compatible then A, B, S and T have a unique common fixed point in X .

Proof: Proof follows from Theorem 3.1 by taking $\alpha_2(h) = \beta_2(h) = \gamma_2(h) = 0$, for all $h \in [0, 1]$.

CONCLUSION

In this paper, we proved a common fixed point theorem in intuitionistic fuzzy metric space using new type of common limit in the range property under functionally rationalized contractive condition for four mappings.

REFERENCES

- [1] M. Aamri and D. El Moutawakil, Some new common fixed point theorems under strict contractive conditions, J. Math. Anal. Appl., 270 (2002), 181-188.
- [2] C. Alaca, I. Altun & D. Turkoglu On compatible mappings of type (I) and (II) in intuitionistic fuzzy metric spaces. Communications of the Korean Mathematical Society, 23 (2008), 427-446.
- [3] C. Alaca, D. Turkoglu and C. Yildiz, Fixed points in intuitionistic fuzzy metric spaces, Chaos Solitons Fractals, 29(5), 2006, 1073-1078.

- [4] S. Chauhan, M.A. Khan and S. Kumar, Unified fixed point theorems in fuzzy metric spaces via common limit range property, J. of Inequalities and Appl., 2013, 2013:182, 1-17.
- [5] S. Chauhan, S. Kutukcu, N. Dhiman and S. Kumar, Variants of R-weakly commuting mappings and common fixed point theorems in intuitionistic fuzzy metric spaces, Kuwait J. Sci. 41 (2) (2014) 49-64.
- [6] B. S. Choudhary, P. Das and P. Saha, Some ψ -contraction results using CLR_g property in probabilistic and fuzzy metric spaces, Ann. Fuzzy Math. Inform. 12 (3) (2016) 387-395.
- [7] S. Chauhan, W. Sintunavarat, P. Kumam, common fixed point theorems for weakly compatible mappings in fuzzy metric spaces using (JCLR) property, Applied Mathematics, 2012, 3, 976-982.
- [8] V. Gupta and A. Kanwar, Some new fixed point results on intuitionistic fuzzy metric spaces, Cogent Mathematics (2016), 3: 1142839, pages 12.
- [9] G. Jungck, & B.E. Rhoades, Fixed point for set valued functions without continuity. Indian Journal of Pure and Applied Mathematics, 29 (1998), 227-238.
- [10] O. Kramosil and J. Michalek, Fuzzy metric and statistical metric spaces, Kybernetika, 11(1975), 326-334.
- [11] V. Popa, A. Patriciu, Fixed point theorems for two pairs of mappings satisfying a new type of common limit range property in G_p metric spaces, Annales Mathematicae Silesianae 32 (2018), 295-312.
- [12] B. Schweizer and A. Sklar, Statistical metric spaces, Pacific J. of Math., 10(1960), 314-334.
- [13] W. Sintunavarat and P. Kumam, Common fixed point theorems for a pair of weakly compatible mappings in fuzzy metric spaces, J. Appl. Math., 2011 (2011) pages 14. Article ID 637958.
- [14] K. Wadhwa and V.P. Bhardwaj, Fixed point theorems for faintly compatible mappings in intuitionistic fuzzy metric space, Journal of the Indian Mathematical Society, Vol. 84, No.1-2, (2017), 130-142.
- [15] K. Wadhwa and V.P. Bhardwaj, Some remarks on results of Pant and Pant along with generalization for hybrid mappings, Annals of Fuzzy Mathematics and Informatics, Vol. 14, No. 3, (2017), 303-313.
- [16] S. Wang, Common fixed point theorems for weakly compatible mappings in fuzzy metric spaces using CLR_g property, J. Nonlinear Sci. Appl. 9 (2016) 1043-1051.
- [17] L.A. Zadeh, Fuzzy sets, Infor. and Control, 8 (1965), 338-353.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2020. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]