

MATRIX REPRESENTATION OF FINITE SOFT TOPOLOGIES

DR. R. ANGELINE CHELLA RAJATHI¹, DR. U. KUMARAN*² AND DR. M. KIRUTHIKA³

¹Assistant Professor, PG and Research Department of Mathematics,
Thiagarajar College, Madurai, India.

²Assistant Professor (Senior Grade) Department of Mathematics,
Ramco Institute of Technology, Rajapalayam, India.

³PG – Teacher, Department of Mathematics, Vedanta Academy, Coimbatore, India.

(Received On: 06-02-21; Revised & Accepted On: 13-03-21)

ABSTRACT

In this paper, matrix representation of a finite soft topology is discussed. Muhammad Shabir and MunazzaNaz have shown that every soft topology gives a parametrized family of topologies on a set X . However they have an example to show that the converse is not true. The converse is discussed here and it is established that every finite soft topology with finite parameter space is represented by a matrix whose elements are sets and each row represents a finite topology on X and conversely every such matrix represents a finite soft topology.

Keywords: Soft sets, Soft Topology, Soft open, Soft closed, Parametrized family.

1. INTRODUCTION

Probability theory, Fuzzy sets [22], intuitionistic fuzzy sets [3], interval mathematics [7] and rough sets [15] deal with uncertainties, but they have their own limitations due to the inadequacy of the parameterization of these theories. To overcome these difficulties Molodtsov [12] initiated the notion of soft sets associated with parameters in the year 1999. Soft set theory is free from the difficulties and soft sets have applications to data reduction [23], data mining [18, 20], medical image processing [17], parameter reduction [6, 21]. In the soft set theory we can use any convenient parameterization strategy. The problem of setting the membership function does not arise in soft set theory, which makes soft set theory very convenient and practicable. Following this, Maji *et.al.* [10, 11] defined several operations on soft set theory and he described an application of soft set theory to decision making problems [9]. Muhammad Shabir introduced the concept of soft topology [13]. Following this several Mathematicians concentrated their studies on applications of soft topology [9]. Both topology and soft topology have application to image processing [17]. The purpose of this paper is to discuss the matrix representation of a finite soft topology.

2. PRELIMINARIES

Let X be an initial universe and E be a set of parameters. A pair (F, E) is called a soft set [4] over X , where $F : E \rightarrow 2^X$ is a mapping. $S(X, E)$ denotes the collection of all soft sets over X with parameter space E . We denote (F, E) by \tilde{F} . Moreover we can denote \tilde{F} by $\tilde{F} = \{(e, F(e)) : e \in E\}$. Soft set theoretic operations are discussed in [10]. Let \tilde{F} and \tilde{G} be any two soft sets over a common universe X with a common parameter space E . \tilde{F} is a soft subset of \tilde{G} if $F(e) \subseteq G(e)$ for all $e \in E$. If \tilde{F} is a soft subset of \tilde{G} then we write $\tilde{F} \subseteq \tilde{G}$. Also $\tilde{F} = \tilde{G}$ if and only if $F(e) = G(e)$ for all $e \in E$.

A soft set $\tilde{\phi}$ over X is said to be a NULL soft set if for all $e \in E$, $\phi(e) = \phi$ (empty set). A soft set \tilde{X} over X is said to be an absolute soft set if for all $e \in E$, $X(e) = X$.

Corresponding Author: Dr. U. Kumaran*²,

²Assistant Professor (Senior Grade) Department of Mathematics,
Ramco Institute of Technology, Rajapalayam, India.

The union of two soft sets \tilde{F} and \tilde{G} over X is defined as $\tilde{F} \cup \tilde{G} = (F \cup G, E)$ where $(F \cup G)(e) = F(e) \cup G(e)$ for all $e \in E$. The intersection of two soft sets \tilde{F} and \tilde{G} over X is defined as $\tilde{F} \cap \tilde{G} = (F \cap G, E)$ where $(F \cap G)(e) = F(e) \cap G(e)$ for all $e \in E$. The arbitrary union and arbitrary intersection of soft sets $(F_\alpha: E)$, $\alpha \in \Delta$ are defined as

$$\cup \left\{ \widetilde{F_\alpha} : \alpha \in \Delta \right\} = \left(\cup \left\{ F_\alpha : \alpha \in \Delta \right\}, E \right) \text{ and } \cap \left\{ \widetilde{F_\alpha} : \alpha \in \Delta \right\} = \left(\cap \left\{ F_\alpha : \alpha \in \Delta \right\}, E \right)$$

where $\left(\cup \left\{ F_\alpha : \alpha \in \Delta \right\} \right)(e) = \cup \left\{ F_\alpha(e) : \alpha \in \Delta \right\}$ and $\left(\cap \left\{ F_\alpha : \alpha \in \Delta \right\} \right)(e) = \cap \left\{ F_\alpha(e) : \alpha \in \Delta \right\}$, for all $e \in E$.

The complement of a soft set \tilde{F} is denoted by $(\tilde{F})' = (F', E)$ where $F' : E \rightarrow 2^X$ is a mapping given by $F'(e) = X - F(e)$ for all $e \in E$.

Let \tilde{F} be a soft set over X and $x \in X$, Then $x \in \tilde{F}$ if $x \in F(e)$ for all $e \in E$.

Definition 2.1: Let X be an initial universal set. If $\tilde{\tau}$ is a collection of soft sets over X , then $\tilde{\tau}$ is said to be a soft topology [10] on X if

- (1) $\tilde{\phi}, \tilde{X}$ belong to $\tilde{\tau}$
- (2) Arbitrary union of soft sets in $\tilde{\tau}$ belongs to $\tilde{\tau}$,
- (3) The intersection of any two soft sets in $\tilde{\tau}$ belongs to $\tilde{\tau}$.

If $\tilde{\tau}$ is a soft topology over X then $(X, \tilde{\tau}, E)$ is a soft space over X .

Lemma 2.2[13]: Let $(X, \tilde{\tau}, E)$ be a soft space over X . Then the collection $(\tilde{\tau})_e = \{ \tilde{F}(e) : \tilde{F} \in \tilde{\tau} \}$ defines a topology on X , for each $e \in E$.

In the next section, $\text{PFT}(\tilde{\tau})$ denotes the parameterized family of topologies on X induced by $\tilde{\tau}$. That is $\text{PFT}(\tilde{\tau}) = \{ (\tilde{\tau})_e : e \in E \}$. $|\text{PFT}(\tilde{\tau})|$ denotes the cardinality of $\text{PFT}(\tilde{\tau})$.

3. PARAMETRIZED FAMILY OF TOPOLOGIES INDUCED BY A SOFT TOPOLOGY

Proposition 3.1: Let $\tilde{\tau}$ be a soft topology over X with parameter space E . Then $|\text{PFT}(\tilde{\tau})| \leq |E|$ and $|(\tilde{\tau})_e| \leq |\tilde{\tau}|$ for every $e \in E$.

Proof: Define $\varphi : E \rightarrow \text{PFT}(\tilde{\tau})$ by $\varphi(e) = (\tilde{\tau})_e$. Clearly φ is onto. Therefore $|\text{PFT}(\tilde{\tau})| \leq |E|$. Now define $\theta : \tilde{\tau} \rightarrow (\tilde{\tau})_e$ by $\theta(\tilde{F}) = \tilde{F}(e)$. Obviously θ is onto. Therefore $|(\tilde{\tau})_e| \leq |\tilde{\tau}|$.

It is note worthy to see that there is a soft topological space $(X, \tilde{\tau}, E)$ for which $|(\tilde{\tau})_e| < |\tilde{\tau}|$ and $|\text{PFT}(\tilde{\tau})| = |E|$ as shown in the next example.

Example 3.2: Let $X = \{a_1, a_2, a_3\}$, $E = \{e_1, e_2\}$ and $\tilde{\tau} = \{ \tilde{\phi}, \tilde{X}, \tilde{F}_1, \tilde{F}_2, \tilde{F}_3, \tilde{F}_4, \tilde{F}_5, \tilde{F}_6, \tilde{F}_7, \tilde{F}_8, \tilde{F}_9 \}$ where

$$\begin{aligned} \tilde{F}_1 &= \{(e_1, \{a_2\}), (e_2, \{a_1\})\} \\ \tilde{F}_2 &= \{(e_1, \{a_2, a_3\}), (e_2, \{a_1, a_2\})\} \quad \tilde{F}_3 = \{(e_1, \{a_1, a_2\}), (e_2, \{a_1, a_2\})\} \\ \tilde{F}_4 &= \{(e_1, \{a_1, a_2\}), (e_2, \{a_1, a_3\})\} \quad \tilde{F}_5 = \{(e_1, \{\tilde{X}\}), (e_2, \{a_1, a_2\})\} \\ \tilde{F}_6 &= \{(e_1, \{a_2\}), (e_2, \{a_1, a_2\})\} \quad \tilde{F}_7 = \{(e_1, \{a_2, a_3\}), (e_2, \{\tilde{X}\})\}, \\ \tilde{F}_8 &= \{(e_1, \{a_1, a_2\}), (e_2, \{\tilde{X}\})\}, \quad \tilde{F}_9 = \{(e_1, a_2), (e_2, \{\tilde{X}\})\} \end{aligned}$$

Then $\tilde{\tau}$ defines a soft topology on X and hence $(X, \tilde{\tau}, E)$ is a soft topological space over X . It can be easily seen that $(\tilde{\tau})_{e_1} = \{ \{\phi, X\}, \{a_2\}, \{a_1, a_2\}, \{a_2, a_3\} \}$

and $(\tilde{\tau})_{e_2} = \{ \{\phi, X\}, \{a_1\}, \{a_1, a_3\}, \{a_2, a_3\} \}$ are topologies on X .

Here $(\tilde{\tau})_{e_1} \neq (\tilde{\tau})_{e_2}$. $\text{PFT}(\tilde{\tau}) = \{(\tilde{\tau})_{e_1}, (\tilde{\tau})_{e_2}\}$. $|\text{PFT}(\tilde{\tau})|=2$ and $|E| = 2$

This shows that $\text{PFT}(\tilde{\tau}) = 2 = |E|$ and $|(\tilde{\tau})_{e_1}| = 5 < 11 = |\tilde{\tau}|$.

The next example shows that $|\text{PFT}(\tilde{\tau})| < |E|$ for some soft topological space $(X, \tilde{\tau}, E)$.

Example 3.3: Let $X = \{b_1, b_2, b_3\}$, $E = \{k_1, k_2\}$ and $\tilde{\tau} = \{\tilde{\phi}, \tilde{X}, \tilde{G}_1, \tilde{G}_2, \tilde{G}_3, \tilde{G}_4, \tilde{G}_5, \tilde{G}_6\}$

where $\tilde{G}_1 = \{(k_1, \{b_2\}), (k_2, \{b_2\})\}$, $\tilde{G}_2 = \{(k_1, \{X\}), (k_2, \{b_2, b_3\})\}$,

$\tilde{G}_3 = \{(k_1, \{b_2\}), (k_2, \{X\})\}$, $\tilde{G}_4 = \{(k_1, \{b_2\}), (k_2, \{b_2, b_3\})\}$

$\tilde{G}_5 = \{(k_1, \{b_2, b_3\}), (k_2, \{X\})\}$, $\tilde{G}_6 = \{(k_1, \{b_2, b_3\}), (k_2, \{b_2, b_3\})\}$

Then $\tilde{\tau}$ defines a soft topology on X and hence $(X, \tilde{\tau}, E)$ is a soft topological space over X . It can be easily seen that

$(\tilde{\tau})_{k_1} = \{\phi, X, \{b_2\}, \{b_2, b_3\}\}$ and $(\tilde{\tau})_{k_2} = \{\phi, X, \{b_2\}, \{b_2, b_3\}\}$ are topologies on X .

Here $(\tilde{\tau})_{k_1} = (\tilde{\tau})_{k_2}$ and $|\text{PFT}(\tilde{\tau})| = 1$ and $|E| = 2$

This proves $|\text{PFT}(\tilde{\tau})| < |E|$.

It is interesting to see that there is a soft topology for which

$$|(\tilde{\tau})_e| = |\tilde{\tau}|$$

Example 3.4: Let $X = \{d_1, d_2, d_3\}$, $E = \{\beta_1, \beta_2\}$ and $\tilde{\tau} = \{\tilde{\phi}, \tilde{X}, \tilde{L}_1, \tilde{L}_2\}$

where $\tilde{L}_1 = \{(\beta_1, \{d_2, d_3\}), (\beta_2, \{d_2, d_3\})\}$, $\tilde{L}_2 = \{(\beta_1, \{d_2\}), (\beta_2, \{d_2\})\}$

Then $\tilde{\tau}$ defines a soft topology on X and hence $(X, \tilde{\tau}, E)$ is a soft topological space over X . For every parameter of E . It can be easily seen that

$(\tilde{\tau})_{\beta_1} = \{\phi, X, \{d_2\}, \{d_2, d_3\}\}$ and $(\tilde{\tau})_{\beta_2} = \{\phi, X, \{d_2\}, \{d_2, d_3\}\}$ are topologies on X .

Here $(\tilde{\tau})_{\beta_1} = (\tilde{\tau})_{\beta_2}$. $|\text{PFT}(\tilde{\tau})| = 4$ and $|\tilde{\tau}| = 4$. Hence $|\text{PFT}(\tilde{\tau})| = |\tilde{\tau}|$ and $|\text{PFT}(\tilde{\tau})| = 1 < 2 = |E|$.

4. MATRIX REPRESENTATION OF A SOFT TOPOLOGY

Muhammad Shabir *et al.* established that every soft topology induces a parameterized family of topological spaces and further gave an example to show that the converse is not true. (Computers and Mathematics with Applications 61(2011) 1786-1799). Then the following question will arise.

Given a collection $\{\tau_e : e \in E\}$ of topologies on X , does there exist a soft topology $\tilde{\tau}$ over X with parameter space E such that $(\tilde{\tau})_e = \tau_e$, for all $e \in E$. This question is discussed in this section.

Let $(X, \tilde{\tau}, E)$ be a soft topological space with finite parameter space $E = \{e_1, e_2, \dots, e_n\}$. According to Shabir, $\tilde{\tau}$ induces a parameterized collection of topologies $(\tilde{\tau})_e, e \in E$. That is for each e_i , let $(\tilde{\tau})_{e_i} = \{G_{i1}, G_{i2}, \dots, G_{in}\}$ be the topology over X , for each $e_i \in E$. Let

$$\text{Mat}(\tilde{\tau}) = \begin{bmatrix} (\tilde{\tau})_{e_1} \\ (\tilde{\tau})_{e_2} \\ \vdots \\ (\tilde{\tau})_{e_n} \end{bmatrix} \dots \dots \dots (*)$$

$$= \begin{bmatrix} G_{11} & G_{12} & G_{13} & \cdots & G_{1n} \\ G_{21} & G_{22} & G_{23} & \cdots & G_{2n} \\ G_{31} & G_{32} & G_{33} & \cdots & G_{3n} \\ \cdots & \cdots & \cdots & \cdots & \vdots \\ \cdots & \cdots & \cdots & \cdots & \vdots \\ G_{n1} & G_{n2} & G_{n3} & \cdots & G_{nn} \end{bmatrix} \text{ where } G_{i1} = \phi \text{ and } G_{in} = X .$$

Therefore every soft topology $\tilde{\tau}$ over X can be identified by a matrix of the form (*)

Remark 4.1: The soft topologies in Example 3.3, Example 3.4 and Example 3.5 can be represented in the matrix forms.

$$\text{Example 3.3: } \text{Mat}(\tilde{\tau}) = \begin{bmatrix} (\tilde{\tau})_{e_1} \\ (\tilde{\tau})_{e_2} \end{bmatrix} = \begin{bmatrix} \phi & \{a_2\} & \{a_1, a_2\} & \{a_2, a_3\} & X \\ \phi & \{a_1\} & \{a_1, a_3\} & \{a_2, a_3\} & X \end{bmatrix} .$$

$$\text{Example 3.4: } \text{Mat}(\tilde{\tau}) = \begin{bmatrix} (\tilde{\tau})_{k_1} \\ (\tilde{\tau})_{k_2} \end{bmatrix} = \begin{bmatrix} \phi & \{b_2\} & \{b_2, b_3\} & X \\ \phi & \{b_2\} & \{b_2, b_3\} & X \end{bmatrix} .$$

$$\text{Example 3.5: } \text{Mat}(\tilde{\tau}) = \begin{bmatrix} (\tilde{\tau})_{\beta_1} \\ (\tilde{\tau})_{\beta_2} \end{bmatrix} = \begin{bmatrix} \phi & \{d_2\} & \{d_2, d_3\} & X \\ \phi & \{d_2\} & \{d_2, d_3\} & X \end{bmatrix} .$$

Theorem 4.2: Let $E = \{e_1, e_2, e_3, \dots, e_n\}$ be a finite parametric space.

$$\text{Let } M = \begin{bmatrix} G_{11} & G_{12} & G_{13} & \cdots & G_{1n} \\ G_{21} & G_{22} & G_{23} & \cdots & G_{2n} \\ G_{31} & G_{32} & G_{33} & \cdots & G_{3n} \\ \cdots & \cdots & \cdots & \cdots & \vdots \\ \cdots & \cdots & \cdots & \cdots & \vdots \\ G_{n1} & G_{n2} & G_{n3} & \cdots & G_{nn} \end{bmatrix} \text{ where } G_{i1} = \phi \text{ and } G_{in} = X \text{ such that}$$

- 1) i^{th} row represents a topology τ_{e_i} on X,
- 2) For each j, k there exists l such that $G_{ij} \cup G_{ik} = G_{il}$ for all i
- 3) For each j, k there exists r such that $G_{ij} \cap G_{ik} = G_{ir}$ for all i.

Then there is a soft topology $\tilde{\tau}$ over X with the parameter space E such that $M = \text{Mat}(\tilde{\tau})$

Proof: Define $\tilde{F}_j : E \rightarrow 2^X$ by $\tilde{F}_j(e_i) = G_{ij}$ for $i=1,2,3, \dots, n$. Then $\{\tilde{F}_j : j = 1, 2, \dots, n\}$ is a collection of soft sets over X with parameter space E.

Claim : $\tilde{\tau} = \{\tilde{F}_1, \tilde{F}_2, \dots, \tilde{F}_n\}$ is a soft topology on X

$$\tilde{F}_1(e_i) = G_{i1} = \phi \text{ that implies } \tilde{\phi} = \tilde{F}_1 \in \tilde{\tau}$$

$$\tilde{F}_n(e_i) = G_{in} = X \text{ so that } \tilde{X} = \tilde{F}_n \in \tilde{\tau}$$

$$(\tilde{F}_j \cup \tilde{F}_k)(e_i) = F_j(e_i) \cup F_k(e_i) = G_{ij} \cup G_{ik} = G_{il} = \tilde{F}_l(e_i). \text{ That is } \tilde{F}_j \cup \tilde{F}_k = \tilde{F}_l \in \tilde{\tau}.$$

$$(\tilde{F}_j \cap \tilde{F}_k)(e_i) = F_j(e_i) \cap F_k(e_i) = G_{ij} \cap G_{ik} = G_{ir} = \tilde{F}_r(e_i)$$

$$\text{i.e. } \tilde{F}_j \cap \tilde{F}_k = \tilde{F}_r \in \tilde{\tau}$$

Then $\tilde{\tau}$ is a soft topology over X with a parameter space E.

Claim: $(\tilde{\tau})_{e_i} = \tau_{e_i}$ for $i = 1, 2, \dots, n$

$$(\tilde{\tau})_{e_i} = \{\tilde{F}_j(e_i) : \tilde{F}_j \in \tilde{\tau}\} = \{G_{ij} : j = 1, 2, \dots, n\} = (\tau_{e_i})_{e_i} \text{ for } i = 1, 2, \dots, n$$

$$\therefore \text{Mat}(\tilde{\tau}) = \begin{bmatrix} (\tilde{\tau})_{e_1} \\ (\tilde{\tau})_{e_2} \\ \vdots \\ (\tilde{\tau})_{e_n} \end{bmatrix} = \begin{bmatrix} \tau_{e_1} \\ \tau_{e_2} \\ \vdots \\ \tau_{e_n} \end{bmatrix} = \begin{bmatrix} G_{11} & G_{12} & G_{13} & \cdots & G_{1n} \\ G_{21} & G_{22} & G_{23} & \cdots & G_{2n} \\ G_{31} & G_{32} & G_{33} & \cdots & G_{3n} \\ \cdots & \cdots & \cdots & \cdots & \vdots \\ \cdots & \cdots & \cdots & \cdots & \vdots \\ G_{n1} & G_{n2} & G_{n3} & \cdots & G_{nm} \end{bmatrix} = M$$

The above theorem is justified in the following example.

Example 4.3: Let $X = \{a, b, c\}$, $E = \{e_1, e_2, e_3\}$ and

$$\tau_{e_1} = \{ \phi, \{a\}, \{a, b\}, X \} \quad \tau_{e_2} = \{ \phi, \{b\}, \{b, c\}, X \} \quad \tau_{e_3} = \{ \phi, \{c\}, \{a, c\}, X \}$$

Then matrix representation of the above family of topologies is

$$M = \begin{bmatrix} \tau_{e_1} \\ \tau_{e_2} \\ \tau_{e_3} \end{bmatrix} = \begin{bmatrix} \phi & \{a\} & \{a, b\} & X \\ \phi & \{b\} & \{b, c\} & X \\ \phi & \{c\} & \{a, c\} & X \end{bmatrix} = \begin{bmatrix} G_{11} & G_{12} & G_{13} & G_{14} \\ G_{21} & G_{22} & G_{23} & G_{24} \\ G_{31} & G_{32} & G_{33} & G_{34} \end{bmatrix}$$

It is easy to verify the conditions 1, 2, and 3 of the above theorem. It can be easily verified.

Each column represents a soft set over X . The first column represents $\tilde{\phi}$, Second column represents \tilde{F}_1 , Third column represents \tilde{F}_2 and Fourth column represents \tilde{X} where

$$\tilde{F}_1(e_1) = \{a\}, \tilde{F}_1(e_2) = \{b\} \text{ and } \tilde{F}_1(e_3) = \{c\}$$

$$\tilde{F}_2(e_1) = \{a, b\}, \tilde{F}_2(e_2) = \{b, c\} \text{ and } \tilde{F}_2(e_3) = \{a, c\}$$

$$\tilde{F}_3(e_1) = \{a, b\}, \tilde{F}_3(e_2) = \{b\} \text{ and } \tilde{F}_3(e_3) = \{c\}$$

$$\tilde{F}_4(e_1) = \{a\}, \tilde{F}_4(e_2) = \{b, c\} \text{ and } \tilde{F}_4(e_3) = \{c\} \text{ and}$$

$$\tilde{F}_5(e_1) = \{a, b\}, \tilde{F}_5(e_2) = \{b, c\} \text{ and } \tilde{F}_5(e_3) = \{c\}$$

Therefore $\tilde{F}_1 = \{(e_1, \{a\}), (e_2, \{b\}), (e_3, \{c\})\}$, $\tilde{F}_2 = \{(e_1, \{a, b\}), (e_2, \{b, c\}), (e_3, \{a, c\})\}$,

$$\tilde{F}_3 = \{(e_1, \{a, b\}), (e_2, \{b\}), (e_3, \{c\})\}, \tilde{F}_4 = \{(e_1, \{a\}), (e_2, \{b, c\}), (e_3, \{c\})\}, \text{ and}$$

$$\tilde{F}_5 = \{(e_1, \{a, b\}), (e_2, \{b, c\}), (e_3, \{c\})\}$$

Then $\tilde{\tau} = \{\tilde{F}_1, \tilde{F}_2, \tilde{F}_3, \tilde{F}_4, \tilde{F}_5\}$ is a soft topology over X with a parameter space E .

Here $(\tilde{\tau})_{e_1} = \{ \phi, \{a\}, \{a, b\}, X \} = \tau_{e_1}$ $(\tilde{\tau})_{e_2} = \{ \phi, \{b\}, \{b, c\}, X \} = \tau_{e_2}$ and $(\tilde{\tau})_{e_3} = \{ \phi, \{c\}, \{a, c\}, X \} = \tau_{e_3}$

$$\text{Therefore Mat}(\tilde{\tau}) = \begin{bmatrix} \tau_{e_1} \\ \tau_{e_2} \\ \tau_{e_3} \end{bmatrix} = \begin{bmatrix} \phi & \{a\} & \{a, b\} & X \\ \phi & \{b\} & \{b, c\} & X \\ \phi & \{c\} & \{a, c\} & X \end{bmatrix} = \begin{bmatrix} G_{11} & G_{12} & G_{13} & G_{14} \\ G_{21} & G_{22} & G_{23} & G_{24} \\ G_{31} & G_{32} & G_{33} & G_{34} \end{bmatrix} = M$$

Thus the theorem is justified by this example.

5. CONCLUSION

Soft topology is characterized using matrix under suitable conditions. Every finite family of topologies induces a soft topology with finite parameter space.

REFERENCES

1. Akta.H, Naman, Soft sets and soft groups, Inf. Sci. 177(2007)27262735.
2. Ali.M.I, F. Feng, X.Y. Liu, W.K. Min, M. Shabir, On some new operations in soft set theory, Computers and Math. with Appl.57(2009)1547 - 1553.
3. Atanassov.K, Intuitionistic fuzzy sets, Fuzzy Sets and Systems 20(1986), 87 - 96.
4. Atanassov.K, Operators over interval valued intuitionistic fuzzy sets, Fuzzy Sets and Systems 64(1994) 159 - 174.

5. BanuPazarVarol and HalisAygün, Fuzzy Soft Topology, Hacettepe Journal of Mathematics and Statistics, 41(3)(2012); 407 - 419.
6. Chen.D, The parametrization reduction of soft sets and its applications, Computers and Math. with Appl. 49(2005),757 - 763.
7. Gorzalzany.M.B, A method of inference in approximate reasoning basedon interval-valued fuzzy sets, Fuzzy Sets and Systems 21(1987), 1 - 17.
8. Kong.Z, L. Gao, L. Wong, S. Li, The normal parameter reduction of softsets and its algorithm, J. Comp. Appl. Math. 56(2008),3029 - 3037.
9. Maji.P.K, R. Biswas, R. Roy, An application of soft sets in a decision making problem, Comput. Math.Appl. 44(2002)10771083.
10. Maji.P.K, R. Biswas, R. Roy, Soft set theory, Comput. Math.Appl.45 (2003), 555 - 562.
11. Maji, P.K. Biswas, R.and Roy, A.R. Fuzzy soft sets, J.Fuzzy Math.9(3); 589 - 602; 2001.
12. Molodtsov.D, Soft set theory first results, Comput. Math.Appl.37 (1999)19 - 31.
13. Muhammad Shabir, MunazzaNaz, On Soft Topological Spaces, Computers and Math. With Appl. 61(2011),1786 - 1799.
14. Naim Cagman, SerkanKaratas, Serdar Enginoglu, Soft Topology, Computers and Mathematics with Applications, 62(2011)351 - 358.
15. Pawlak.Z, Rough sets, Int. J. Comput. Sci. 11(1982)341356.
16. SabirHussain, Bashir Ahmad, Soft Seperation axioms in soft topologicalspaces, Hacetepe journal of mathematics and statistics, 44(3) (2015); 559 -568.
17. Saima Anwar Lashari, Rosziati Ibrahim, A Framework for Medical Images Classification Using Soft Set, Procedia Technology,11(2013),548 - 556.
18. SatyaRanjan Dash, Satchidananda Dehuri, Soft set and Genetic Algorithms for Association Rule Mining: A road map and direction for Hybridization, Springer-Verlag Berlin Heidelberg 2013, AISC 199; 381-393.
19. Tanay, B. and Kandemir, M.B. Topological Structures of fuzzy soft sets, Computers and Mathematics with Applications 61, 412-418, 2011.
20. TututHerawan, Mustafa Mat Deris, A soft set approach for association rules mining, Knowledge-Based Systems, 24(2011),186 - 195.
21. Xiuqin Ma, Norrozila Sulaiman, Jansi Mohamad Zain, A new efficient normal parameter reduction algorithm of soft sets, Computers and Mathematics with Applications, 62(2011), 588 - 598.
22. Zadeh.L.A, Fuzzy sets, Inf. Control 8(1965)338353.
23. Zhi Kong, Liqun Gao, Lifu Wang, Steven Li, The normal parameter reduction of soft sets and its algorithm, Computers and Mathematics with Applications, 56(2008), 3029 – 3037.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2021. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]