

COMMON FIXED POINT THEOREM FOR SEMI - COMPATIBLE
AND SUB-SEQUENTIALLY CONTINUOUS MAPPINGS IN FUZZY METRIC SPACE

DR. PREETI SENGAR*

Department of Applied Mathematics and Computational Science, SGSITS, Indore (M.P.), India.

(Received On: 02-03-21; Revised & Accepted On: 10-03-21)

ABSTRACT

The purpose of this paper is to prove some common fixed point theorems in semi-compatible and sub-sequentially continuous mappings in fuzzy metric space.

Keywords: Compatible, Semi-compatible, Sontinuous maps, Sub-sequentially continuous maps.

INTRODUCTION

The concept of fuzzy sets was initially investigated by Zadeh [16] as a new way to represent vagueness in everyday life. Subsequently, it was developed by many authors and used in various fields. To use this concept in Topology and Analysis, several researchers have defined fuzzy metric space in various ways. In this chapter we deal with the fuzzy metric space defined by Kramosil and Michalek [10] and modified by George and Veeramani [4]. Recently, Grebiec [5] has proved fixed points result for Fuzzy metric space. In the sequel, Singh and Chouhan [14] introduced the compatible mappings of Fuzzy metric space and proved the common fixed point theorem.

Using the concept of R-weak commutativity of mappings, Vasuki [15] proved the fixed point theorem for Fuzzy metric space. Recently in 2009, using the concept of sub compatible maps, Bouhadjera *et.al* [1] proved common fixed point theorems. Using the concept of compatible maps of type (A), Jain *et.al.* [8] proved a fixed point theorem for six self maps in a fuzzy metric space. Using the concept compatible maps of type (β) Jain *et.al* [8] proved a fixed point theorem in fuzzy metric space and menger space using the concept of semi-compatibility, weak compatibility and compatibility of type (β) respectively. In this section, we introduced the new concept of semi-compatible and sub-sequentially continuous mapping in fuzzy metric space.

For the sake of completeness, we recall some definition and known results in fuzzy metric space.

Definition 2.1: [12] A triangular norm $*$ (shortly t- norm) is a binary operation on the unit interval $[0, 1]$ to $[0, 1]$ such that for all $a, b, c, d \in [0, 1]$ the following conditions are satisfied.

- (a) $a * 1 = a$,
- (b) $a * b = b * a$,
- (c) $a * b \leq c * d$ whenever $a \leq c$ and $b \leq d$,
- (d) $(a * b) * c = a * (b * c)$.

Example of t-norms are $a * b = \min \{a, b\}$ and $a * b = ab$.

Definition 2.2: [10] A 3-tuple $(X, M, *)$ is said to be a fuzzy metric space if X is an arbitrary set, $*$ is a continuous t-norm and M is a fuzzy set on $X^2 \times [0, \infty)$ satisfying the following conditions for all $x, y, z \in X, t, s > 0$,

- (FM-1) $M(x, y, t) = 0$,
- (FM-2) $M(x, y, t) = 1$ if and only if $x = y$,
- (FM-3) $M(x, y, t) = M(y, x, t)$,
- (FM-4) $M(x, y, t) * M(y, z, s) \leq M(x, z, t + s)$
- (FM-5) $M(x, y, \cdot): [0, \infty) \rightarrow [0, 1]$ is continuous.
- (FM-6) $\lim_{t \rightarrow \infty} M(x, y, t) = 1$.

Corresponding Author: Dr. Preeti Sengar*

Department of Applied Mathematics and Computational Science, SGSITS, Indore (M.P.), India.

In the definition of George and Veeramani [4], M is a fuzzy set on $X^2 \times [0, \infty)$ and (FM-1), (FM-2), (FM-3) are replaced respectively, with (GV-1), (GV-2), (GV-3) below the axiom (GV-2) is reformulated as in [7, Remark 1]:

(GV-1) $M(x, y, t) > 0$, for all $t > 0$;

(GV-2) $M(x, x, t) = 1$ for all $t > 0$ and $x \neq y \Rightarrow M(x, y, t) < 1$. for all $t > 0$;

(GV-5) $M(x, y, \cdot) : (0, \infty) \rightarrow [0, 1]$ is continuous for all $x, y \in X$.

Then M is called a fuzzy metric on X . Then $M(x, y, t)$ denotes the degree of nearness between x and y with respect to t . In the following example, it is showed that every metric induces a fuzzy metric:

Example 2.1: [4] Let (X, d) be a metric space. Define $a * b = ab$ (or $a * b = \min \{a, b\}$) for all $a, b \in X$ and let M be fuzzy set on $X^2 \times [0, \infty)$ defined as follows: $M(x, y, t) = \frac{t}{t+d(x,y)}$

Then $(X, M, *)$ is a fuzzy metric space and the fuzzy metric M induced by the metric d is the standard fuzzy metric.

Example 2.2: [3] Let $X = \mathbb{R}$, the set of all real numbers. Define $a * b = ab$ and

$M(x, y, t) = \left[\exp\left(\frac{x-y}{t}\right) \right]^{-1}$ for all x, y in \mathbb{R} and $t \in [0, \infty)$. then $(X, M, *)$ is a fuzzy metric space.

Lemma 2.1: [5] For all $x, y \in X$, (X, M, \cdot) is non decreasing function.

Definition 2.3: [4] Let $(X, M, *)$ be a fuzzy metric space. A sequence $\{x_n\}$ in X is said to be Convergent to a point $x \in X$ if $\lim_{n \rightarrow \infty} M(x_n, x, t) = 1$ for all $t > 0$. Further, the sequence $\{x_n\}$ is said to be a Cauchy sequence if $\lim_{n \rightarrow \infty} M(x_n, x_{n+p}, t) = 1$ for all $t > 0$ and for all p .

The space is said to be Complete if every Cauchy sequence in it converges to a point on it.

Proposition 2.1: [4] In a fuzzy metric space $(X, M, *)$ the limit of a sequence is unique.

Definition 2.4: [9] A pair (A, S) of self mappings defined on a fuzzy metric space $(X, M, *)$ is said to be compatible if and only if $M(ASx_n, SAx_n, t) \rightarrow 1$ for all $t > 0$ whenever $\{x_n\}$ is a sequence in X such that $Ax_n, Sx_n \rightarrow z$ for some $z \in X$ as $n \rightarrow \infty$.

Definition 2.5: [13] A pair (A, S) of self mappings defined on a fuzzy metric space $(X, M, *)$ is said to be semi-compatible if $\lim_{n \rightarrow \infty} ASx_n = Sx$ whenever there exists a sequence $\{x_n\} \in X$ such that $\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Sx_n = x$ for some $x \in X$.

Definition 2.6: [6] A pair (A, S) of self mappings defined on a fuzzy metric space $(X, M, *)$ is said to be Sub-compatible if and only if there exists a sequence $\{x_n\}$ such that

$$\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Sx_n = z \text{ for some } z \in X \text{ and } \lim_{n \rightarrow \infty} M(ASx_n, SAx_n, t) = 1 \text{ for all } t > 0.$$

Definition 2.7: [2] A pair (A, S) of self mappings defined on a fuzzy metric space $(X, M, *)$ is said to be reciprocally continuous if for a sequence $\{x_n\}$ in X ,

$$\begin{aligned} \lim_{n \rightarrow \infty} ASx_n &= Az \text{ and } \lim_{n \rightarrow \infty} SAx_n = Sz, \text{ whenever} \\ \lim_{n \rightarrow \infty} Ax_n &= \lim_{n \rightarrow \infty} Sx_n = z \text{ for some } z \in X. \end{aligned}$$

Definition 2.8: [11] Two self maps A and B of a fuzzy metric space $(X, M, *)$ are said to be weak compatible if they commute at their coincidence points, i.e. $Ax = Bx \Rightarrow ABx = BAx$.

In this section, we define the concept of sub-sequentially continuous and semi-compatible mapping and we introduced the common fixed point of pairs, coincident point of pair and unique common fixed point.

Definition 2.9: [6] A pair (A, S) of self mappings defined on a fuzzy metric space $(X, M, *)$ is said to be sub-sequentially continuous if and only if there exists sequence $\{x_n\}$ in X such that $\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Sx_n = z$ for some $z \in X$ and $\lim_{n \rightarrow \infty} ASx_n = Az$ and $\lim_{n \rightarrow \infty} SAx_n = Sz$

Theorem 2.1: Let A, B, S, T, P and Q be self mappings of fuzzy metric space $(X, M, *)$ with $a * b = \min \{a, b\}$ for all $a, b \in [0, 1]$. If the pairs (A, PQ) and (B, ST) are semi-compatible and sub-sequentially continuous mappings, then

(a) The pair (A, PQ) has a coincidence point.

(b) The pair (B, ST) has a coincidence point.

(c) Further, the mapping A, B, S, T, P and Q have a unique common fixed point in X provided the involved maps satisfy the inequality

$$M^2(Ax, By, t) * [M(PQx, Ax, t) * M(STy, By, t)] \geq [pM(PQx, Ax, t) + qM(PQx, STy, t)] * M(PQx, By, t) \quad (1.1)$$

for all $x, y \in X$ and $t > 0$, where $0 < p, q < 1$ and $p+q=1$.

Proof: The pairs (A, PQ) and (B,ST) are Semi-compatible and Sub-sequentially continuous mappings, there exists a sequence $\{x_n\}$ in X such that

$$\begin{aligned} \lim_{n \rightarrow \infty} Ax_n &= \lim_{n \rightarrow \infty} PQx_n = z \text{ for some } z \in X \\ \text{and } \lim_{n \rightarrow \infty} M(A(PQ)x_n, (PQ)Ax_n, t) &= 1, \text{ for all } t < 0. \\ \text{and } \lim_{n \rightarrow \infty} M(Az, PQz, t) &= 1 \end{aligned} \tag{1.2}$$

then we have $Az=PQz$,

$$\begin{aligned} \text{Similarly, } \lim_{n \rightarrow \infty} By_n &= \lim_{n \rightarrow \infty} STy_n = w \in X \\ \lim_{n \rightarrow \infty} M(B(ST)y_n, (ST)By_n, t) &= 1, \text{ for all } t < 0 \\ \text{and } \lim_{n \rightarrow \infty} M(Bw, STw, t) &= 1 \end{aligned} \tag{1.3}$$

Hence z is coincidence point of A, PQ and w is coincidence point of B, ST

$$\text{then we get } Az=PQz. \tag{1.4}$$

$$Bw=STw. \tag{1.5}$$

Step-1:- First we prove that $z = w$. Putting $x = x_n, y = y_n$ in inequality (1.1) we have

$$M^2(Ax_n, By_n, t) * [M(PQx_n, Ax_n, t).M(STy_n, By_n, t)] \geq [pM(PQx_n, Ax, t) + qM(PQx_n, STy_n, t)]M(PQx_n, By_n, t) \tag{1.6}$$

Now

$$\begin{aligned} M^2(z, w, t) * [M(z, z, t).M(w, w, t)] &\geq pM[(z, z, t) + qM(z, w, t)]M(z, w, t) \\ M^2(z, w, t) &\geq [p + qM(z, w, t)]M(z, w, t) \\ M(z, w, t) &\geq \frac{p}{1-q} \\ M(z, w, t) &= 1 \end{aligned} \tag{1.7}$$

Thus we have $z = w$

Step-2:- Now we prove that $Az = z$, Putting $x = z$ and $y = y_n$ in (1.1)

$$M^2(Az, By_n, t) * [M(PQz, Az, t).M(STy_n, By_n, t)] \geq [pM(PQz, Az, t) + qM(PQz, STy_n, t)]M(PQz, By_n, t)$$

$$M^2(Az, w, t) * [M(PQz, Az, t).M(w, w, t)] \geq [pM(PQz, Az, t) + qM(PQz, w, t)]M(PQz, w, t)$$

$$M^2(Az, w, t) * [M(Az, Az, t).M(w, w, t)] \geq [pM(Az, Az, t) + qM(Az, w, t)]M(Az, w, t)$$

$$M^2(Az, w, t) \geq [p + qM(Az, w, t)]M(Az, w, t)$$

$$M(Az, w, t) \geq \frac{p}{1-q}$$

$$M(Az, w, t) = 1$$

Hence $Az = w = z$

Step-3:- Again we prove that $Bz = z$

Then we have $x = x_n, y = z$ in (1.1)

$$M^2(Ax_n, Bz, t) * [M(PQx_n, Ax_n, t).M(STz, Bz, t)] \geq [pM(PQx_n, Ax_n, t) + qM(PQx_n, STz, t)] M(PQx_n, Bz, t)$$

$$M^2(z, Bz, t) * [M(Az, Az, t).M(z, z, t)] \geq [pM(Az, Az, t) + qM(z, Bz, t)]M(z, Bz, t)$$

$$M^2(z, Bz, t) \geq [p + qM(z, Bz, t)] M(z, Bz, t)$$

$$M(z, Bz, t) \geq \frac{p}{1-q}$$

$$M(z, Bz, t) = 1$$

we get $z = Bz$

(1.8)

Step-4:- Again we claim that $Tz = z$,

putting $x = Tz$ and $y = z$ in (1.1)

$$M^2(ATz, Bz, t) * [M(PQTz, ATz, t).M(STz, Bz, t)] \geq [pM(PQTz, ATz, t) + qM(PQTz, STz, t)] M(PQTz, Bz, t)$$

$$M^2(TAz, z, t) * [M(TPQz, TAz, t).M(Bz, Bz, t)] \geq [pM(TPQz, ATz, t) + qM(TPQz, Bz, t)] M(TPQz, Bz, t)$$

$$M^2(Tz, z, t) * [M(Tz, Tz, t).M(z, z, t)] \geq [pM(Tz, Tz, t) + qM(Tz, z, t)] M(Tz, z, t)$$

$$M^2(Tz, z, t) \geq [p + qM(Tz, z, t)] M(Tz, z, t)$$

$$M(Tz, z, t) \geq \frac{p}{1-q}$$

$$M(Tz, z, t) = 1$$

we get $Tz = z$

(1.9)

Step-5:- Again we show that $Sz = z$,

putting $x = Sz$ and $y = z$ in (1.1)

$$M^2(ASz, Bz, t) * [M(PQSz, ASz, t).M(STz, Bz, t)] \geq [pM(PQSz, ASz, t) + qM(PQSz, STz, t)] M(PQSz, Bz, t)$$

$$M^2(SAz, z, t) * [M(SPQz, SAz, t).M(Bz, Bz, t)] \geq [pM(SPQz, ASz, t) + qM(SPQz, Bz, t)] M(SPQz, Bz, t)$$

$$M^2(Sz, z, t) * [M(Sz, Sz, t).M(z, z, t)] \geq [pM(Sz, Sz, t) + qM(Sz, z, t)] M(Sz, z, t)$$

$$M^2(Sz, z, t) \geq [p + qM(Sz, z, t)] M(Sz, z, t)$$

$$M(Sz, z, t) \geq \frac{p}{1-q}$$

$$M(Sz, z, t) = 1$$

we get $Sz = z$

(1.10)

Step-6:- Again we prove that $Qz=z$,

putting $x=Qz$ and $y=z$ in (1.1)

$$M^2(AQz, Bz, t) * [M(PQz, AQz, t) * M(SQz, Bz, t)] \geq [pM(PQz, AQz, t) + qM(PQz, SQz, t)] M(PQz, Bz, t)$$

$$M^2(Az, z, t) * [M(QPz, Az, t) * M(Bz, Bz, t)] \geq [pM(QPz, Az, t) + qM(QPz, Bz, t)] M(QPz, Bz, t)$$

$$M^2(Qz, z, t) * [M(Qz, Qz, t) * M(z, z, t)] \geq pM(Qz, Qz, t) + qM(Qz, z, t) M(Qz, z, t)$$

$$M^2(Qz, z, t) \geq [p+qM(Qz, z, t)] M(Qz, z, t)$$

$$M(Qz, z, t) \geq \frac{p}{1-q}$$

$$M(Qz, z, t) = 1$$

we get $Qz = z$

(1.11)

Step-7:- Again we prove that $Pz = z$, putting $x = Pz$ and $y = z$ in (1.1)

$$M^2(APz, Bz, t) * [M(PQz, ATz, t) * M(SPz, Bz, t)]$$

$$[pM(PQz, APz, t) + qM(PQz, SPz, t)] M(PQz, Bz, t)$$

$$M^2(PAz, z, t) * [M(PPz, PAz, t) * M(Bz, Bz, t)] \geq [pM(PPz, APz, t) + qM(PPz, Bz, t)] M(PPz, Bz, t)$$

$$M^2(Pz, z, t) * [M(Pz, Pz, t) * M(z, z, t)] \geq [pM(Pz, Pz, t) + qM(Pz, z, t)] M(Pz, z, t)$$

$$M^2(Pz, z, t) \geq [p+qM(Pz, z, t)] M(Pz, z, t)$$

$$M(Pz, z, t) \geq \frac{p}{1-q}$$

$$M(Pz, z, t) = 1$$

we get $Pz = z$

(1.12)

i.e. $Az = Bz = Tz = Sz = Pz = Qz = z$

Hence z is a common fixed point of A, B, S, T, P , and Q .

REFERENCES

1. Bouhadjera, H. and Godet-Thobie, C., Common fixed point theorems for pairs of Sub-compatible maps, arxiv:0906.315w [math.FA] 17 June 2009.
2. Balasubramaniam, P., Muralisankar S., and Pant R.P., common fixed point of four mappings in a fuzzy metric, J. fuzzy Math. 10(2) (2002), 397-384.
3. Fukushima, H., on non-contractive mappings, Yakohama Math. Jour. 19(1) (1971), 205-230.
4. George A. and Veeramani, P. on some result in fuzzy metric space, fuzzy sets and system 64(1994), 395-399. MR1289545.
5. Grabiec M., fixed point in fuzzy metric space, fuzzy sets and system 27(3) (1988), 385-389 MR0956385 (89J:54007).
6. Gopal D. and Imdad, M. some new common fixed point theorems in fuzzy metric space, Ann. Univ. Ferrara Sez. VII Sci. Mat. 57(2) (2011), 303-316. MR2855364
7. Gregori, V. and Romaguera, S., Some properties of fuzzy metric spaces, Fuzzy Sets and System 115(2000), 485-489.
8. Jain, A. and Singh, B., fixed point theorem using semi-compatibility and weak-compatibility in fuzzy metric spaces, V.J.M.S.T (1), (2007), 139-147.
9. Kolmogorov, A.N. and Fromin, S.J., Element of theory of functions, Functional Analysis Vol. 1, Graylock Press. Rochester, 1957.
10. Kramosil, I. and Michalek, J., Fuzzy metric and statistical metric space, Kybernetika (Prague) 11(5) (1975), 336-344.
11. Mishra S.N., Sharma N. and Singh S.L., Common fixed point of maps on fuzzy metric space, Int. J. Math. Sci. 17(2) (1994), 253-258.
12. Schweizer B. and Sklar A., Probabilistic Metric Space, North-Holland Series in Probability and Applied Mathematics. North-Holland Publishing Co., New York, (1983). ISBN: 0-444-00666-4 MR0790314.
13. Singh, B. and Jain, S., Semi-compatible and fixed point theorems in fuzzy metric space, Chungcheong Math. Soc. 18 (2005), 1-22.
14. Singh, B. and Chouhan, M.S., Common fixed points of compatible maps in fuzzy metric spaces, fuzzy sets and systems 115(2000), 471-475.
15. Vasuki, R., Common fixed point for R-weakly commuting maps in fuzzy metric spaces, Indian J. pure Appl. Math. 30(4), (1999), 419-423.
16. Zadeh, L. A., Fuzzy sets, Inform. And control 89 (1965), 338-353.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2021. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]